

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE COHOMOLOGY OF COMPACT LIE GROUPS
Autor: Reeder, Mark
Kapitel: 6. The cohomology of a Lie group
DOI: <https://doi.org/10.5169/seals-61824>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 10.12.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

where, as in (3.3), ∂_i is the derivation of \mathcal{S} extending the functional $\lambda \mapsto \lambda(H_i)$. We have a perfect pairing

$$\mathcal{D} \otimes \mathcal{S} \rightarrow \mathbf{R}$$

given by $(D, f) = (Df)(0)$. Since the pairing is perfect, something in degree v must pair nontrivially with Π . Since an irreducible W -module can only pair nontrivially with its dual, and the self-dual character ε occurs with multiplicity one in \mathcal{D}^v , afforded by $\partial_1 \cdots \partial_v$, we must have $\partial_1 \cdots \partial_v \Pi \neq 0$, so $c(\Pi) \neq 0$.

Observe that $\partial_1 \cdots \partial_v$ is analogous to the fundamental class of G/T , and the pairing is essentially that between homology and cohomology. We further remark that in fact all irreducible representations of W are defined over the rational numbers, hence they are all self dual. This is a consequence of Springer's cohomological construction of W -modules [Sp].

Returning again to our task, we now inductively assume that $c: \mathcal{H}^k \rightarrow H^{2k}(G/T)$ is injective for $k \leq v$, and let $V = \mathcal{H}^{k-1} \cap \ker c$. Note that V is preserved by W since c is W -equivariant. The sign character does not occur in \mathcal{H}^{k-1} , so there is a positive root α whose corresponding reflection s_α does not act by $-I$ on V . Decompose $V = V_+ \oplus V_-$ according to the eigenspaces of s_α . If $V \neq 0$ then $V_+ \neq 0$, so take $f \in V_+$. Now $c(\alpha f) = c(\alpha)c(f) = 0$, and αf is in degree k , so we must have $\alpha f \in \mathcal{I}$ by the induction hypothesis. Let $h_1, \dots, h_{|W|}$ be a basis of \mathcal{H} with h_1, \dots, h_r s_α -skew and the rest s_α invariant. By Chevalley's theorem (3.2), we can write $\alpha f = \sum h_i \sigma_i$ with σ_i W -invariant of positive degree. Since αf is s_α -skew, the sum only goes up to r . Now for $i \leq r$, the polynomial h_i must vanish on $\ker \alpha$, hence can be written $h_i = \alpha h'_i$ for some $h'_i \in \mathcal{S}$. But then $f = \sum_{i=1}^r h'_i \sigma_i \in \mathcal{I}$. Since f is supposed to be harmonic, we must have $f = 0$. Hence c is injective on \mathcal{H} , and the proof of Borel's theorem is complete. \square

6. THE COHOMOLOGY OF A LIE GROUP

We now have all the ingredients for our proof. Consider the map $\psi: G/T \times T \rightarrow G$ given by $\psi(gT, t) = gtg^{-1}$. The Weyl group W acts on T by conjugation and on G/T by $w \cdot gT = gn^{-1}T$, where $w = nT$. Hence W acts on $H(G/T \times T) = H(G/T) \otimes H(T)$. Since $\psi(gn^{-1}T, wtw^{-1}) = \psi(gT, t)$, it follows that the induced map ψ^* on cohomology maps $H(G)$ to $[H(G/T) \otimes H(T)]^W$. Though we prefer to have it in this form, the latter group could be thought of as the cohomology of the quotient of $G/T \times T$

by the action of W , and this quotient has a natural interpretation. As in the introduction, let M be the set of pairs (g, T') where T' is a maximal torus in G containing $g \in G$. The map $G/T \times_W T \rightarrow M$ sending $(gT, t) \bmod(W)$ to (gtg^{-1}, gTg^{-1}) is a diffeomorphism.

PROPOSITION (6.1). *The map induced by ψ on cohomology is an isomorphism of graded rings*

$$\psi^*: H(G) \xrightarrow{\cong} [H(G/T) \otimes H(T)]^W.$$

Proof. We compute the derivative $(d\psi)_{(gT, t)}$ at a point $(gT, t) \in G/T \times T$. For each point $gT \in G/T$, we identify \mathfrak{m} with the tangent space $T_{gT}(G/T)$ by letting $X \in \mathfrak{m}$ correspond to the initial tangent vector X_{gT} of the path $s \mapsto g(\exp sX)T$ in G/T . Similarly, an element $X \in \mathfrak{g}$ (resp. $H \in \mathfrak{t}$) corresponds to a tangent vector $X_g \in T_g(G)$ (resp. $H_t \in T_t(T)$, for $t \in T$).

Then

$$\begin{aligned} (d\psi)_{gT, t}(X_{gT}, 0) &= \frac{d}{ds} g(\exp sX) t(\exp -sX) g^{-1} \Big|_{s=0} \\ &= \frac{d}{ds} gtg^{-1} [\exp sAd(gt^{-1})X] [\exp -sAd(g)X] \Big|_{s=0} \\ &= \frac{d}{ds} gtg^{-1} [X + sAd(g)(Ad(t^{-1}) - 1)X + O(s^2)] \Big|_{s=0} \\ &= [Ad(g)(Ad(t^{-1}))X]_{gtg^{-1}}. \end{aligned}$$

Similarly, we find, for $H \in \mathfrak{t}$, that

$$(d\psi)_{gT, t}(0, H_t) = [Ad(g)H]_{gtg^{-1}}.$$

Hence, under the identifications, $(d\psi)_{(gT, t)}$ is the map

$$(Ad(t^{-1}) - I)_{\mathfrak{m}} \oplus I: \mathfrak{m} \oplus \mathfrak{t} \rightarrow \mathfrak{m} \oplus \mathfrak{t} = \mathfrak{g}.$$

Here the subscript \mathfrak{m} means we view $Ad(t^{-1}) - I$ as a map from \mathfrak{m} to itself. Now G being compact and connected, we must have $\det Ad(t) = 1$, so

$$\det (d\psi)_{(gT, t)} = \det (I - Ad(t))_{\mathfrak{m}}.$$

(Actually, \mathfrak{m} is always even-dimensional as we have seen, so there is no need to reverse the subtraction).

We compute the degree of ψ by finding a regular value. Let t_0 be a generic element in T , as in (2.3). Consider $\psi^{-1}(t_0) = \{(gT, t): gtg^{-1} = t_0\}$. It turns out that any two elements of T conjugate in G must be conjugate by an element

of W . (In $U(n)$, two diagonal matrices with the same set of eigenvalues must be conjugate by a permutation matrix.) It follows easily then that

$$\psi^{-1}(t_0) = \{(wT, wt_0 w^{-1}) : w \in W\}.$$

We next show that ψ preserves orientation at each point in $\psi^{-1}(t_0)$. The eigenvalues of $Ad(t_0)$ in \mathfrak{m} are complex conjugate pairs z, \bar{z} , where $|z| = 1, z \neq 1$. Hence $|1 - z||1 - \bar{z}| = 2(1 - \operatorname{Re}(z)) > 0$, so $\det(I - Ad(t_0))_{\mathfrak{m}} > 0$.

At this point we know the degree of ψ is $\deg \psi = |W| \neq 0$. By Poincaré duality, any smooth map between compact manifolds of the same dimension is injective on cohomology as soon as it has nonzero degree. Hence $\psi^*: H(G) \rightarrow [H(G/T) \times H(T)]^W$ is injective. We finish the proof of (6.1) by showing that both sides have the same dimension.

For this we use, three times, the following basic principle. Let K be a compact group (here K will be G, T or W). Let dk be the left invariant Haar measure on K with total mass one. Let V be a finite dimensional real vector space, and $\rho: K \rightarrow GL(V)$ a continuous group homomorphism. Then the space V^K of vectors fixed by all $\rho(k), k \in K$, has dimension

$$\dim V^K = \int_K \operatorname{trace} \rho(k) dk.$$

To compute this integral over G , we must exploit further the computation of $d\psi$. Let $\omega_G, \omega_T, \omega_{G/T}$ be the unique invariant (under left translations by G, T , and G respectively) differential forms of top degree whose integral over the respective manifold is one. The pull-back formula for integration gives

$$\int_G f \omega_G = \frac{1}{\deg \psi} \int_{G/T \times T} f \circ \psi(gT, t) |\det(d\psi)_{(gT, t)}| \omega_{G/T} \wedge \omega_T,$$

where the determinant is computed with respect to bases spanning parallelograms of unit volume with respect to the appropriate forms. Taking f to be invariant under conjugation by G , we arrive at the famous Weyl integration formula:

$$\int_G f \omega_G = \frac{1}{|W|} \int_T f(t) \det(I - Ad(t))_{\mathfrak{m}} \omega_T.$$

Expand the function $t \mapsto \det(I - Ad(t))_{\mathfrak{m}}$ in a sum of characters of T : $n_0 \chi_0 + n_1 \chi_1 + \cdots + n_k \chi_k$. Here χ_0 is the trivial character of T ,

appearing n_0 times, and for $i > 0$ each χ_i is a nontrivial character appearing n_i times. Taking for f the constant function equal to one, and applying the basic principle of invariants to T , we find $n_0 = |W|$.

Taking for f the function $f(g) = \det(I + Ad(g))$, i.e., the trace of $Ad(g)$ acting on $\Lambda \mathfrak{g}$, we find, using the Cartan-de Rham isomorphism (4.3), that

$$\begin{aligned} \dim H(G) &= \dim(\Lambda \mathfrak{g})^G = \int_G \det(I + Ad(g)) \omega_G \\ &= \frac{1}{|W|} \int_T \det(I + Ad(t)) \det(I - Ad(t))_{\mathfrak{m}} \omega_T \\ &= \frac{2^{\dim T}}{|W|} \int_T \det(I - Ad(t^2))_{\mathfrak{m}} \omega_T. \end{aligned}$$

Now the squaring map on T is surjective, so the square of a nontrivial character of T is still nontrivial. Hence the trivial character again appears with multiplicity $|W|$ in the expansion of $\det(I - Ad(t^2))_{\mathfrak{m}}$. This multiplicity is the value of the integral, so $\dim H(G) = 2^{\dim T} = 2^l$.

On the other hand, we saw in (5.3) that the trace of $w \in W$ acting on $H(G/T)$ is $|W|$ if $w = 1$, zero otherwise. Applying the invariance formula one more time, we find that $\dim [H(G/T) \otimes H(T)]^W = 2^l$ as well, completing the proof of (6.1). \square

We now have the main result

(6.2) THEOREM. *The cohomology ring $H(G)$ with real coefficients is a bigraded exterior algebra with generators in bi-degrees $(2m_i, 1)$, for $1 \leq i \leq l$.*

Proof. By (6.1) and (5.4), we have

$$H(G) \simeq [H(G/T) \otimes H(T)]^W \simeq [\mathcal{H}_{(2)} \otimes \Lambda]^W,$$

and by (3.8), the latter space is an exterior algebra with generators in degrees $(2m_i, 1)$, for $1 \leq i \leq l$. \square

Moreover, from the multiplicity formula (3.8), the dimensions of the bi-graded pieces are given in terms of the exponents as follows

(6.3) COROLLARY. *For each $q \geq 0$, we have*

$$\sum_{n=0}^{\dim G} \dim [H^{n-q}(G/T) \otimes H^q(T)]^W u^n = u^q s_q(u^{2m_1}, \dots, u^{2m_l}).$$

(6.4) We give two interpretations of the bigrading. First, we follow [L] and consider the spectral sequence of the fibration $G \rightarrow G/T$, which has E_2 -term

$$E_2^{pq} = H^p(G/T) \otimes H^q(T),$$

and converges to $H(G)$. This spectral sequence does not degenerate at E_2 , but it has a spectral subsequence which does degenerate, and still converges to $H(G)$.

To see this we again consider the Weyl group action. More precisely, N acts by automorphisms of the fibration $G \rightarrow G/T$, which in turn induce automorphisms of each term in the spectral sequence, commuting with the differentials. On $E_2^{pq} = H^p(G/T) \otimes H^q(T)$, the action of N factors through W and is the same as that considered above. Thus we have representations of W on the spaces E_2^{pq} , hence on each E_r^{pq} for $r \geq 2$.

For each p, q, r we decompose $E_r^{pq} = (E_r^{pq})^W \oplus (E_r^{pq})_W$, where the subscript W indicates a W -stable complement to the invariants. Each of the latter two spaces is a spectral subsequence, and since E_∞^{pq} is a subquotient of $H^{p+q}(G)$ and N acts trivially on $H(G)$ (because G is connected), we must have $(E_\infty^{pq})_W = 0$. On the other hand, $(E_\infty^{pq})^W$ is a subquotient of $(E_2^{pq})^W = [H^p(G/T) \otimes H^q(T)]^W$, so we have

$$\begin{aligned} 2^l = \dim H(G) &= \sum_{p,q} \dim (E_\infty^{pq})^W \leq \sum_{p,q} \dim (E_2^{pq})^W \\ &= \sum_q \dim [H(G/T) \otimes \Lambda^q]^W = 2^l. \end{aligned}$$

It follows that $\dim (E_\infty^{pq})^W = \dim (E_2^{pq})^W$ for all p, q , so the spectral subsequence of W -invariants degenerates at $(E_2)^W$, and (6.1) is proved again.

(6.5) The significance of the bigrading on $H(G)$ can be seen in yet another way, inspired by [GHV]. We consider, for a fixed integer $k \neq 1$, the k^{th} -power maps $x \mapsto x^k$, denoted p_k and P_k , on T and G , respectively. It is shown in [GHV] that the Lefschetz number of P_k equals that of p_k , namely $(1-k)^l$. Let $H^n(G)_q$ be the k^q -eigenspace of P_k^* acting on $H^n(G)$. It is further shown in [GHV] that $\sum_n \dim H^n(G)_q = \binom{l}{q}$. We can refine this by computing each $\dim H^n(G)_q$ separately. Consider the commutative diagram

$$\begin{array}{ccc}
H(G) & \xrightarrow{\psi^*} & [H(G/T) \otimes H(T)]^W. \\
P_k^* \downarrow & & \downarrow 1 \otimes P_k^* \\
H(G) & \xrightarrow{\psi^*} & [H(G/T) \otimes H(T)]^W.
\end{array}$$

Since p_k^* acts by k^q on $H^q(T)$, (6.1) implies that $H^n(G)_q \simeq [H^{n-q}(G/T) \otimes H^q(T)]^W$, and (6.3) gives the dimension of the latter space.

(6.6) This last interpretation of the bigrading shows that it is natural in the following sense. Suppose $f: K \rightarrow G$ is a homomorphism between two compact connected Lie groups. Since f commutes with the power maps P_k on G and K , the cohomology map f^* sends $H^n(G)_q$ to $H^n(K)_q$. Suppose for example that K is a closed connected subgroup of G and f is the inclusion map. Choose, as we may, a maximal torus T of G such that $S := T \cap K$ is a maximal torus of K . The restriction map $H(G) \rightarrow H(K)$ becomes, via (6.1), the map $[H(G/T) \otimes H(T)]^W \rightarrow [H(K/S) \otimes H(S)]^{W_K}$ induced by restriction on each factor, where W_K is the Weyl group of S in K .

(6.7) We close with the homology interpretation of (6.1), which says the homology map ψ_* induced by ψ is surjective. It follows that the homology of G is spanned by the cycles $[\psi(\bar{X}_w, T_I)] = \{gtg^{-1}: gT \in \bar{X}_w, t \in T_I\}$. Here $w \in W$, X_w is the Schubert cell (see (5.2)) and $T_I = \prod_{i \in I} T_i$, where $T = T_1 \times \cdots \times T_l$, with each $T_i \simeq S^1$. Using the results in [BGG], one can explicitly write down the action of W on $H_*(G/T)$ in terms of the Schubert cell basis, and this leads, in principle, to the linear relations in $H_*(G)$ satisfied by the cycles $[\psi(\bar{X}_w, T_I)]$.

REFERENCES

- [A] ADAMS, J.F. *Lectures on Lie Groups*. W.A. Benjamin, 1969.
- [BGG] BERNSTEIN, I.N., I.M. GEL'FAND and S.I. GEL'FAND. Schubert cells and cohomology of the spaces G/P . *Representation theory — selected papers*. Cambridge University Press, 1982, 115-139.
- [Bo1] BOREL, A. Topology of Lie groups and characteristic classes. *Bull. Amer. Math. Soc.* (1955), 397-432.
- [Bo2] ——— Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts. *Ann. Math.* 57 (1953), 115-207.
- [Bk] BOURBAKI, N. *Groupes et algèbres de Lie*. Hermann, Paris, 1968.
- [BT] BOTT, R. and L. TU. *Differential Forms in Algebraic Topology*. Springer Verlag, 1982.