

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SYNTHETIC PROJECTIVE GEOMETRY AND POINCARÉ'S THEOREM ON AUTOMORPHISMS OF THE BALL

Autor: Shiffman, Bernard

Bibliographie

DOI: <https://doi.org/10.5169/seals-61825>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

sion $2n - 1$ in \mathcal{M}_{B_n} , it follows that $\psi \circ (f \times g)$ vanishes on the connected component of $\Omega \cap \mathcal{M}_{B_n}$ containing (z_0, \bar{z}_0) . After shrinking U if necessary, we can assume that $\psi \circ (f \times g)$ vanishes on $\Omega \cap \mathcal{M}_{B_n}$ and thus $(f \times g)(\Omega \cap \mathcal{M}_{B_n}) \subset \mathcal{M}_{B_n}$. We consider the embedding $\iota \times \iota: \mathbf{C}_n \times \mathbf{C}_n \hookrightarrow \mathbf{P}_{\mathbf{C}}^n \times \mathbf{P}_{\mathbf{C}}^n$ given by $\iota(z_1, \dots, z_n) = (\sqrt{-1}: z_1 : \dots : z_n)$, which maps \mathcal{M}_{B_n} onto a (dense open) subset of $\mathcal{M}_{\mathbf{C}}^n$. By Corollary 7 applied to the maps

$$\tilde{f} = \iota \circ f \circ \iota^{-1}: \iota(U) \rightarrow \iota(\hat{U}), \quad \tilde{g} = \iota \circ g \circ \iota^{-1}: \iota(V) \rightarrow \iota(\hat{V}),$$

there exists $A \in \mathrm{PGL}(n+1, \mathbf{C})$ such that $\tilde{f} = A|_{\iota(U)}$. Thus f extends to the fractional linear map $\iota^{-1} \circ A \circ \iota$, which gives an automorphism of B_n .

We now give a simplified form of Alexander's proof [Al, p. 250] that the Jacobian matrix of the map f must be nonsingular at some point of $U \cap \partial B_n$. We begin by observing that $f^{-1}(\partial B_n)$ is nowhere dense. Indeed, suppose on the contrary that $f^{-1}(\partial B_n)$ contains a connected open set U_0 and assume without loss of generality that $f(z_0) = (1, 0, \dots, 0)$ for some point $z_0 \in U_0$. Then by the maximum principle, $f_1 \equiv 1$ and hence $f \equiv (1, 0, \dots, 0)$ on U_0 and thus on U , contradicting the assumption that f is nonconstant. Now suppose on the contrary that the Jacobian determinant of f vanishes identically on $U \cap \partial B_n$. Since the zero of the Jacobian determinant is an analytic subvariety, the Jacobian determinant must vanish identically on U . As a consequence, the fibers of f contain no isolated points. Assume without loss of generality that $(1, 0, \dots, 0) \in U$ and choose $r < 1$ such that the spherical cap $W := \{z \in B_n : \operatorname{Re} z_1 > r\}$ is contained in U . Choose a point $p \in W$ such that $f(p) \notin \partial B_n$. Let A be the connected component of $f^{-1}(f(p)) \cap W$ that contains p ; A is an analytic subvariety of W of positive dimension. Furthermore $\bar{A} \setminus A \subset \{z \in \mathbf{C}^n : \operatorname{Re} z_1 = r\}$. By the maximum principle (see for example [Gu, Theorem H2]) applied to the holomorphic function $\varphi: A \rightarrow \mathbf{C}$ given by $\varphi(z) = \exp z_1$, we conclude that φ is constant and thus $\bar{A} \setminus A = \emptyset$ so that A is a compact subvariety of W of positive dimension, which is impossible. \square

REFERENCES

- [Al] ALEXANDER, H. Holomorphic mappings from the ball and polydisc. *Math. Ann.* 209 (1974), 249-256.
- [Ar] ARTIN, E. *Geometric Algebra*. Interscience Publishers, New York, 1957.
- [BB] BLASCHKE, W. and G. BOL. *Geometrie der Gewebe*. Springer, Berlin, 1938.
- [Ca] CARTAN, E. Sur le groupe de la géométrie hypersphérique. *Comm. Math. Helv.* 4 (1932), 158-171.

- [CG] CHERN, S.-S. and P. GRIFFITHS. Abel's theorem and webs. *Jahresber. Deutsch. Math.-Verein.* 80 (1978), 13-110. Corrections and addenda. *Jahresber. Deutsch. Math.-Verein.* 83 (1981), 78-83.
- [CJ] CHERN, S.-S. and S. JI. Projective geometry and Riemann's mapping problem. Preprint, 1994.
- [Co] COXETER, H.S.M. *Projective Geometry*. University of Toronto Press, Toronto, 1974.
- [Fo] FORSTNERIČ, F. Proper holomorphic mappings: a survey. *Several Complex Variables: Proceedings of the Mittag-Leffler Institute, 1987-1988* (J.E. Fornaess, ed.). Princeton Univ. Press, Princeton, 1993, 297-363.
- [Go] GOLDBERG, V. *Theory of Multicodimensional (n + 1)-Webs*. Kluwer, Dordrecht, 1988.
- [Gu] GUNNING, R.C. *Introduction to Holomorphic Functions of Several Variables, Volume II: Local Theory*. Brooks/Cole, Pacific Grove, CA, 1990.
- [MY] MOK, N. and S.K. YEUNG. Geometric realizations of uniformization of conjugates of hermitian locally symmetric manifolds. *Complex Analysis and Geometry* (V. Ancona and A. Silva, eds.). Plenum Press, New York, 1993, 253-270.
- [MM] MOLZON, R. and K.P. MORTENSEN. The Schwarzian derivative for maps between manifolds with complex projective connections. Preprint, 1994.
- [Pe] PELLES, D. Proper holomorphic self-maps of the unit ball. *Math. Ann.* 190 (1971), 298-305. Correction. *Math. Ann.* 202 (1973), 135-136.
- [Po] POINCARÉ, H. Les fonctions analytiques de deux variables et la représentation conforme. *Rend. Circ. Mat. Palermo* (1907), 185-220.
- [Pr] PRENOWITZ, W. The characterization of plane collineations in terms of homologous families of lines. *Trans. Amer. Math. Soc.* 38 (1935), 564-599.
- [Ra] RADÓ, F. Non-injective collineations on some sets in Desarguesian projective planes and extension of non-commutative valuations. *Aequationes Math.* 4 (1970), 307-321.
- [Re] REIDEMEISTER, K. Topologische Fragen der Differentialgeometrie, V. *Math. Z.* 29 (1929), 427-435.
- [Ru] RUDIN, W. *Function Theory in the Unit Ball of \mathbf{C}^n* . Springer-Verlag, New York, 1980.
- [Ta] TANAKA, N. On pseudo-conformal geometry of hypersurfaces of the space of n complex variables. *J. Math. Soc. Japan* 14 (1962), 397-429.
- [We] WEBSTER, S. On the mapping problem for algebraic real hypersurfaces. *Inventiones Math.* 43 (1977), 53-68.

(Reçu le 5 août 1994)

Bernard Shiffman

Department of Mathematics
 Johns Hopkins University
 Baltimore, MD 21218
 U.S.A.

vide-leer-empty