Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	41 (1995)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	PLURIDIMENSIONAL ABSOLUTE CONTINUITY FOR DIFFERENTIAL FORMS AND THE STOKES FORMULA
Autor:	Jurchescu, Martin / Mitrea, Marius
Kapitel:	5. The global form of the Stokes formula on \$C^1\$ manifolds
DOI:	https://doi.org/10.5169/seals-61826

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

as $\varphi(Q) = 0$ for any Q of the types (1)-(2) described above, and since φ is additive, it follows that $\psi(Q'_i) = \varphi(Q_i)$ for any $i \in J$. In particular,

$$\left|\sum_{i \in J} \Psi(Q'_i)\right| = \left| \varphi(P) \right| \leq \left| \int_{\overset{\circ}{\Omega} \cap \partial P} u \right| + \left| \int_{\overset{\circ}{P} \cap b\Omega} u \right| + \left| \iint_{P \cap \Omega} f \right|.$$

By (4.2), the uniformly local (n-1)-integrability of u, the integrability of u on $b\Omega$ and the integrability of f on Ω , the right hand side of the above equality can be made arbitrarily small, provided $\sum_{i \in J} \mu_{n-1}(Q'_i)$ is sufficiently small. However, since A' has Lebesgue measure zero in \mathbb{R}^{n-1} , this can be readily taken care of and this completes the proof of the theorem. \Box

REMARK 4.3. As an inspection of the proofs shows, Theorem 4.1 and Lemma 4.2 continue to hold in the case when the locally (n - 1)-integrable form u is uniformly (n - 1)-integrable only in a small neighborhood of $\mathcal{S}(u)$.

5. The global form of the Stokes formula on C^1 manifolds

In this section we shall present a coordinate free version of the main result of section 4. Throughout this section, we let M be a fixed, oriented, Hausdorff, differentiable manifold of class C^1 , and real dimension n.

DEFINITION 5.1. A subset Ω of M is called a C^1 domain if for any $a \in \Omega \setminus \mathring{\Omega}$, there exist an open neighborhood U of a in M and a C^1 diffeomorphism $f = (f_1, f_2, ..., f_n)$ of U onto an open neighborhood V of the origin in \mathbb{R}^n , such that

$$U \cap \Omega = \{x \in U; f_n(x) \leq 0\}.$$

Clearly, the border of the domain Ω , $b\Omega := \Omega \setminus \mathring{\Omega}$ is either the empty set or a (n-1)-dimensional C^1 -submanifold of M assumed with the standard induced orientation. Note that a simple application of the implicit function theorem shows that any C^1 domain is also a Lipschitz domain in \mathbb{R}^n .

It is not difficult to see that the class of Lipschitz domains described in Definition 1.1 is not invariant under the action of bi-Lipschitz diffeomorphisms of \mathbf{R}^n . In particular, Theorem 4.1 cannot be reformulated invariantly. To remedy this, for the rest of this section we shall slightly adjust our previous definitions to the C^1 framework by carrying out the following simple modification. That is, whenever applicable, we shall replace "Lipschitz embedding" by " C^1 -embedding", i.e. Lipschitz embeddings which are C^1 functions. Note that, in particular, the condition (1.1) is in this case equivalent with the continuity of the functions

$$\frac{\partial h(s,x)}{\partial x_i}: S \times \omega \to \Omega, \quad i = 1, 2, ..., n-1 .$$

Assuming this modification, all the previously introduced notions become invariant to C^1 diffeomorphisms and, hence, meaningful on C^1 manifolds. More specifically, we make the following.

DEFINITION 5.2. Let Ω be a C^1 domain of M. A (n-1)-form uis said to be absolutely continuous (uniformly (n-1)-locally integrable) on Ω if for any point $P \in \Omega$ there exists a local coordinate map $h: U \to \mathbb{R}^n$ of M with $P \in U$ such that $(h^{-1})^*u$ is absolutely continuous (uniformly (n-1)-locally integrable, respectively) on $h(U \cap \Omega)$.

Let u and f be locally integrable forms on M, having degrees (n-1)and n, respectively. Recall that du = f on a open set Ω of M in the distribution sense, if for any $\varphi \in C_0^1(\Omega)$,

$$\int_M d\phi \wedge u = -\int_M \phi f \, .$$

THEOREM 5.3. Let Ω be a C^1 domain of M, and u a (n-1)-form compactly supported in M. Assume that u is uniformly (n-1)-locally integrable and absolutely continuous on Ω , and that the singular set

$$\mathscr{S}(u) := (\overline{\Omega} \setminus \Omega) \cap \sup u$$

has (n-1)-dimensional Hausdorff measure zero.

If u is integrable on $b\Omega$ and du (taken in the sense of distribution theory) is integrable on Ω , then

$$\int_{b\Omega} u = \iint_{\hat{\Omega}} du .$$

Proof. Using a smooth partition of unity and then working in local coordinates we can assume that $M = \mathbb{R}^n$. In this case, the conclusion is provided by Theorem 4.1.

Note that, here again it suffices to have the "uniform" part of the local (n-1)-integrability condition for u fulfilled only on a small neighborhood of $\mathcal{S}(u)$ (cf. also Remark 4.3).

DEFINITION 5.4. A closed subset A of M is said to have an almost regular boundary if A coincides with the closure of its interior and if there exist a family $(S_i)_{i \in I}$ of C^1 submanifolds of M and a locally finite family $(C_i)_{i \in I}$ of compact subsets of M such that:

(1) $C_i \subset S_i$, for any $i \in I$, and $\mathring{C}_i \cap \mathring{C}_j = \emptyset$, for all $i \neq j$ (the interiors are taken in S_i and in S_j , respectively);

(2) $C_i \cap C_j$ has (n-1)-dimensional Hausdorff measure zero for all $i \neq j$;

 $(3) \quad \partial A = \cup_{i \in I} C_i.$

Note that if A has an almost regular boundary, then

$$\Omega := \overset{\circ}{A} \cup \left(\bigcup_{i \in I} \overset{\circ}{C}_i \right)$$

(the interior of A is taken in M) is a C^1 domain with border $b\Omega = \bigcup_{i \in I} \check{C}_i$. If u is an integrally continuous (n-1)-form on M, it follows that u is integrable on each oriented submanifold \mathring{C}_i (with the standard orientation induced by \mathring{A}). Since ∂C_i has zero measure in S_i , we can define

$$\int_{\partial A} u := \sum_{i \in I} \int_{\mathring{C}_i} u$$

whenever $A \cap \text{supp } u$ is compact. Hence, without further proof, we can state the following.

THEOREM 5.5. Let A be a subset of M with an almost regular boundary. If u is a (n-1)-form which is uniformly (n-1)-locally integrable on M, absolutely continuous on M, and for which $A \cap \text{supp } u$ is compact, then u is integrable on ∂A , du is integrable on $\overset{\circ}{A}$ and

$$\int_{\partial A} u = \iint_{A} du .$$