

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	41 (1995)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	RATIONALITY PROBLEMS FOR K-THEORY AND CHERN-SIMONS INVARIANTS OF HYPERBOLIC 3-MANIFOLDS
Autor:	Neumann, Walter D. / Yang, Jun
Kapitel:	5. Chern-Simons invariants and the regulator map
DOI:	https://doi.org/10.5169/seals-61828

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

5. CHERN-SIMONS INVARIANTS AND THE REGULATOR MAP

As we have seen from the above discussion, hyperbolic 3-manifolds and their volumes all have interesting K -theoretic interpretation. Another important invariant in the theory of hyperbolic 3-manifolds is the Chern-Simons invariant of [4] and [9]. What is its relation with K -theory? As discussed in sect. 1, any hyperbolic 3-manifold M represents a class $\beta(M)$ in $\mathcal{B}(\mathbf{C})$. For M compact, the next theorem follows from Theorem 1.11 of Dupont [5]. A proof for cusped M can be found in [14].

THEOREM 5.1. *The Chern-Simons invariant of $M \bmod \mathbf{Q}(2)$ is equal to the real part of the image of $\beta(M)$ under the regulator map*

$$\rho : \mathcal{B}(\mathbf{C}) \rightarrow \mathbf{C}/\mathbf{Q}(2) .$$

□

Now given a hyperbolic 3-manifold M with invariant trace field k and the associated embedding $\sigma : k \hookrightarrow \mathbf{C}$, the class $\beta(M) \in \mathcal{B}(\mathbf{C})$ associated to M is in the image of $\mathcal{B}(k)_{\mathbf{Q}} \xrightarrow{\sigma} \mathcal{B}(\mathbf{C})$ (Theorem 1.1). If σ is a CM -embedding, then it follows from Corollary 3.2 that the real part of ρ is trivial on $\mathcal{B}(F) \otimes \mathbf{Q}$. Theorem A in the introduction now follows immediately from Theorem 5.1. Similarly, as in the previous section, Corollary 3.2 implies that the irrationality conjecture for Chern Simons invariant of the Introduction would be implied by the Ramakrishnan Conjecture.

We have seen that the imaginary part of ρ is essentially the volume map while the real part of ρ can be called the Chern-Simons map. Reinterpreting the discussion of the previous section, this means that if $k = \bar{k} \subset \mathbf{C}$, then the volume map on $\mathcal{B}(k)$ factors through $\mathcal{B}_-(k)$ and Chern-Simons map on $\mathcal{B}(K)$ factors over $\mathcal{B}_+(k)$. By Theorem B we thus get bounds of $\frac{1}{2}(r_2 + r'_2)$ and $\frac{1}{2}(r_2 - r'_2)$ on the number of rationally independent volumes resp. Chern-Simons invariants for manifolds having invariant trace field contained in our given k . Ramakrishnan's Conjecture says the image of ρ has \mathbf{Q} -rank r_2 . This is equivalent to the conjecture that the \mathbf{Q} -ranks of the images of $vol : \mathcal{B}(k) \rightarrow \mathbf{R}$ and $CS : \mathcal{B}(k) \rightarrow \mathbf{R}/\mathbf{Q}$ are exactly $\frac{1}{2}(r_2 + r'_2)$ and $\frac{1}{2}(r_2 - r'_2)$.