Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	41 (1995)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	CUBIC FORMS AND COMPLEX 3-FOLDS
Autor:	Okonek, Ch. / Van de Ven, A.
Kapitel:	1.1 HOMEOMORPHISM TYPES AND \$C^\infty\$-STRUCTURES
DOI:	https://doi.org/10.5169/seals-61829

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

It is related to the well-known inequality  $c_1^2 \leq 3c_2$  and has been solved to a considerable extent.

Though in the case of 6-folds the corresponding question about the realisability of cubic forms is definitely weaker than the question which 6-folds carry a complex or algebraic structure, it still remains of much interest. In the second half of this paper we say something about algebra and arithmetic of cubic forms and consider the apparently largely untouched question of the realisability of *complex* forms by complex manifolds. Apart from a considerable number of examples some conditions for Kähler manifolds are given. And to show how few 6-folds of the type in question actually carry Kähler structures, we add a theorem about Kähler structures on the set of 6-folds with  $b_2 = 1$ ,  $b_3 \leq$  constant and  $w_2 \neq 0$ .

The first part of this paper surveys the results of Wall and Jupp referred to before, and deals with the homotopy classification. By putting together (for the first time?) all this in a rather systematic way we hope to contribute to the knowledge of complex 3-folds from a topological point of view.

Acknowledgements: We would like to thank the following mathematicians for very helpful remarks and suggestions: F. Grunewald, G. Harder, F. Hirzebruch, and R. Schulze-Pillot. We also want to acknowledge support by the Science project "Geometry of Algebraic Varieties" SCI-0398-C(A); by the Max-Planck-Institut für Mathematik in Bonn, and by the Schweizer Nationalfond (Nr. 21-36111.92).

# 1. TOPOLOGICAL CLASSIFICATION OF CERTAIN 6-MANIFOLDS

The topological classification of 1-connected, closed, oriented, 6-dimensional manifolds has been developped in a sequence of papers by C.T.C. Wall [W], P. Jupp [J], and A. Žubr [Z1], [Z2], [Z3]. Roughly speaking, their main result is that the topological classification of these 6-manifolds is equivalent to the arithmetic classification of certain systems of invariants naturally associated with them.

The aim of this section is to review these results and to reformulate the arithmetic classification problem in a way which makes it accessible to further investigation.

## 1.1 Homeomorphism types and $C^{\infty}$ -structures

Let X be a closed, oriented, 6-dimensional topological manifold; we assume that X is 1-connected with torsion-free homology. The *basic invariants* of X are [J]:

- i)  $H^{2}(X, \mathbb{Z})$ , a finitely generated free abelian group;
- ii)  $b_3(X) = rk_{\mathbb{Z}}H^3(X, \mathbb{Z})$ , a natural number which is even since  $H^3(X, \mathbb{Z})$  admits a non-degenerate symplectic form;
- iii)  $F_X: H^2(X, \mathbb{Z}) \otimes H^2(X, \mathbb{Z}) \otimes H^2(X, \mathbb{Z}) \to \mathbb{Z}$ , a symmetric trilinear form given by the cup-product evaluated on the orientation class;
- iv) p<sub>1</sub>(X) ∈ H<sup>4</sup>(X, Z), the first Pontrjagin class which is always integral because the inclusion of BO in BTOP induces an isomorphism H<sup>4</sup>(BTOP, Z) → H<sup>4</sup>(BO, Z) [J];
- v)  $w_2(X) \in H^2(X, \mathbb{Z}_{/2})$ , the second Stiefel-Whitney class;  $w_2(X)$  is determined by the Steenrod square  $Sq^2: H^4(X, \mathbb{Z}_{/2}) \to H^6(X, \mathbb{Z}_{/2})$ ,  $Sq^2(\xi) = w_2(X) \cdot \xi \ \forall \xi \in H^4(X, \mathbb{Z}_{/2}) \ [W];$
- vi)  $\tau(X) \in H^4(X, \mathbb{Z}_{/2})$ , the triangulation class which is the obstruction to lifting the stable tangent bundle of Y to a *PL* bundle [J].

These invariants satisfy one fundamental relation

(\*) 
$$W^3 \equiv (p_1(X) + 24T) \cdot W \pmod{48}$$

for all integral classes  $W \in H^2(X, \mathbb{Z})$ ,  $T \in H^4(X, \mathbb{Z})$  with  $\overline{W} \equiv w_2(X) \pmod{2}$ ,  $\overline{T} \equiv \tau(x) \pmod{2}$ .

For smooth manifolds (\*) is simply the  $\hat{A}$ -integrality theorem of A. Borel and F. Hirzebruch [B/H], whereas for topological manifolds additional surgery arguments are necessary [J].

In the sequel we shall use Poincaré duality to identify  $H^4(X, \mathbb{Z})$  with  $\operatorname{Hom}_{\mathbb{Z}}(H^2(X, \mathbb{Z}), \mathbb{Z})$ , so that  $p_1(X)$  can be considered as a linear form on  $H^2(X, \mathbb{Z})$ , and we will write  $x \cdot y \cdot z$  instead of  $F_X(x \otimes y \otimes z)$  for elements  $x, y, z \in H^2(X, \mathbb{Z})$ .

DEFINITION 1. A system of invariants is a 6-tuple  $(r, H, w, \tau, F, p)$  consisting of a non-negative integer r, a finitely generated free abelian group H, elements  $w \in H/_{2H}$  and  $\tau \in H^{\vee}/_{2H^{\vee}}$ , a symmetric trilinear form  $F \in S^3 H^{\vee}$ , and a linear form  $p \in H^{\vee}$ . The system  $(H, r, w, \tau, F, p)$  is admissible iff for every  $W \in H$  and  $T \in H^{\vee}$  with  $\overline{W} \equiv w \pmod{2}$  and  $\overline{T} \equiv \tau \pmod{2}$  the following congruence holds:

(\*) 
$$W^3 \equiv (p + 24T) (W) \pmod{48}$$
.

Two systems of invariants  $(H, r, w, \tau, F, p)$  and  $(H', r', w', \tau', F', p')$ are equivalent iff r = r', and there exists an isomorphism  $\alpha: H \rightarrow H'$ such that:

$$\alpha(w) = w', \quad \alpha^*(\tau') = \tau, \quad \alpha^*(F') = F, \quad \alpha^*(p') = p.$$

The main classification result can now be formulated in the following way:

THEOREM 1 (Jupp). The assignment

 $X \mapsto \left(\frac{b_3(X)}{2}, H^2(X, \mathbb{Z}), w_2(X), \tau(X), F_X, p_1(X)\right)$ 

induces a 1-1 correspondence between oriented homeomorphism classes of 1-connected, closed, oriented, 6-dimensional topological manifolds with torsion-free homology, and equivalence classes of admissible systems of invariants.

Furthermore, a topological manifold X as above admits a  $C^{\infty}$ -structure if and only if the triangulation class  $\tau(X)$  vanishes; the  $C^{\infty}$ -structure is then unique.

REMARK 1. The classification theorem is due to C.T.C. Wall in the special case of differentiable spin-manifolds [W]; the final form above was obtained by P. Jupp [J].

A. Žubr generalized Wall's result in another direction; he proved a classification theorem for 1-connected, smooth spin-manifolds with not necessarily torsion-free homology [Z1]; in two further papers [Z2], [Z3] he also obtains P. Jupp's classification, and he asserts in addition, that algebraic isomorphisms of systems of invariants can always be realized by orientation preserving homeomorphisms (diffeomorphisms in the smooth case).

Note that the first invariant  $\frac{b_3(X)}{2}$  of the system is completely independent of the remaining invariants, so that the following splitting theorem holds:

COROLLARY 1. Every 1-connected, closed, oriented, 6-dimensional, topological (differentiable) manifold X with torsion-free homology admits a topological (differentiable) splitting  $X = X_0 \# \frac{b_3(X)}{2} (S^3 \times S^3)$  as a connected sum of a core  $X_0$  with  $b_3(X_0) = 0$ , and  $\frac{b_3(X)}{2}$  copies of  $S^3 \times S^3$ . The oriented homeomorphism (diffeomorphism) type of  $X_0$ is unique.

EXAMPLE 1. The 1-connected, closed, oriented 6-manifolds X with  $H_2(X, \mathbb{Z}) = 0$  are  $S^6$  and the connected sums  $\#_r S^3 \times S^3$  of  $r \ge 1$  copies of  $S^3 \times S^3[Sm]$ .

## 1.2 Homotopy types

In order to describe the homotopy classification of the 6-manifolds above, we need some more preparations.