Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CUBIC FORMS AND COMPLEX 3-FOLDS
Autor: Okonek, Ch. / Van de Ven, A.
Kapitel: 2.2 Homotopy types with a given cohomology ring
DOI: https://doi.org/10.5169/seals-61829

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 25.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

Proof. $W_\circ$ is characteristic for F if and only if $q_{\bar{F}} = \bar{F}^t(W_\circ)$.

In terms of a $\mathbf{Z}$-basis $\{e_1, \ldots, e_b\}$ for H the condition $q_{\bar{F}} \in \operatorname{Im}(\bar{F}^t)$ translates into a simple rank condition over $\mathbf{Z}_{/2}$: the $\mathbf{Z}_{/2}$-rank of the $b \times \binom{b+1}{2}$-matrix A representing $\bar{F}^t$ must be equal to the $\mathbf{Z}_{/2}$-rank of the matrix A extended by the column $(\bar{e}_i \cdot \bar{e}_j \cdot (\bar{e}_i + \bar{e}_j))_{1 \leqslant i \leqslant j \leqslant b}$

EXAMPLE 3. Let $H = \mathbf{Z}e_1 \oplus \mathbf{Z}e_2$ be free of rank 2, $F \in S^3 H^\vee$ given by $e_1^3 = a$, $e_1^2 e_2 = b$, $e_1 e_2^2 = c$, $e_2^3 = d$ with $a, b, c, d \in \mathbf{Z}$. The rank condition becomes

$$rk_2 \begin{bmatrix} \bar{a} & \bar{b} \\ \bar{c} & \bar{d} \\ \bar{b} & \bar{c} \end{bmatrix} = rk_2 \begin{bmatrix} \bar{a} & \bar{b} & \bar{0} \\ \bar{c} & \bar{d} & \bar{0} \\ \bar{b} & \bar{c} & \overline{b+c} \end{bmatrix}$$

2.2 HOMOTOPY TYPES WITH A GIVEN COHOMOLOGY RING

Our next task is to describe the set of oriented homotopy types of 1-connected, closed, oriented, 6-dimensional manifolds with a fixed torsion-free cohomology ring.

From Žubr's classification theorem we know that in algebraic terms this means the following: fix a non-negative integer $r_\circ$, a finitely generated free abelian group $H_\circ$, and a symmetric trilinear form $F_\circ \in S^3 H_\circ^\vee$ which admits characteristic elements.

Let $\mathscr{M}(r_\circ, H_\circ, F_\circ)$ be the set of 1-connected, closed, oriented, 6-dimensional manifolds X with $b_3(X) = 2r_\circ$, such that there exists an isomorphism $\alpha: H_\circ \to H^2(X, \mathbf{Z})$ with $\alpha^* F_X = F_\circ$. Denote by $\operatorname{Aut}(F_\circ)$ the subgroup of $\mathbf{Z}$-automorphisms of $H_\circ$ which leave $F_\circ \in S^3 H_\circ^\vee$ invariant; $\operatorname{Aut}(F_\circ)$ acts on pairs $(w, [l]) \in \bar{H}_\circ \times H_\circ^\vee/_{48H_\circ^\vee}/_{U_{F_\circ}}$ in a natural way:

$$\gamma \cdot (w, [l]) := (\gamma(w), (\gamma^{-1})^* [l]) .$$

Let ${}_{\operatorname{Aut}(F_\circ)}\backslash \bar{H}_\circ \times H_\circ^\vee/_{48H_\circ^\vee}/_{U_{F_\circ}}$ be the set of $\operatorname{Aut}(F_\circ)$-orbits.

A manifold X in $\mathscr{M}(r_\circ, H_\circ, F_\circ)$ and an isomorphism $\alpha: H_\circ \to H^2(X, \mathbf{Z})$ with $\alpha^* F_X = F_\circ$ yields a well-defined $\operatorname{Aut}(F_\circ)$-orbit:

$$(\alpha^{-1}(w_2(X)), \alpha^*[p_1(X) + 24T]) \text{ (modulo } \operatorname{Aut}(F_\circ)) ,$$

where $T \in H^4(X, \mathbf{Z})$ is an arbitrary integral lifting of $\tau(X) \in H^4(X, \mathbf{Z}_{/2})$.

The set of oriented homotopy types $\mathscr{M}(r_\circ, H_\circ, F_\circ)/_\simeq$ of manifolds in $\mathscr{M}(r_\circ, H_\circ, F_\circ)$ can now be described in the following way:

PROPOSITION 3. *The assignment* $X \mapsto (\alpha^{-1}(w_2(X)), \alpha^*[p_1(X) + 24T])$ *(modulo* $\mathrm{Aut}(F_0)$*) defines an injection.*

$$I: \mathcal{M}(r_0, H_0, F_0)/_{\simeq} \to {}_{\mathrm{Aut}(F_0)}\backslash \bar{H}_0 \times H_0^{\vee}/_{48H_0^{\vee}}/_{U_{F_0}}.$$

Proof. Suppose X and X' are manifolds in $\mathcal{M}(r_0, H_0, F_0)$, $\alpha: H_0 \to H^2(X, \mathbf{Z})$ and $\alpha': H_0 \to H^2(X', \mathbf{Z})$ isomorphisms with $\alpha^* F_X = F_0$ and $(\alpha')^* F_{X'} = F_0$. X and X' have the same image under I iff there exists an automorphism $\gamma \in \mathrm{Aut}(F_0)$ with $\gamma\alpha^{-1}(w_2(X)) = (\alpha')^{-1} w_2(X')$ and $(\gamma^{-1})^* \alpha^* [p_1(X) + 24T] = (\alpha')^*[p_1(X') + 24T']$. Consider $\beta := \alpha \circ \gamma \circ \alpha^{-1}: H^2(X, \mathbf{Z}) \to H^2(X', \mathbf{Z})$; β is obviously an isomorphism with $\beta^* F_{X'} = F_X$, $\beta w_2(X) = w_2(X')$, and $\beta^*[p_1(X') + 24T'] = [p_1(X) + 24T]$; but this means that the systems of invariants associated with X and X' are weakly equivalent, and therefore X and X' oriented homotopy equivalent.

A complete description of the set $\mathcal{M}(r_0, H_0, F_0)/_{\simeq}$ i.e. of the image of I is only possible if the automorphism group $\mathrm{Aut}(F_0)$ is known; this can be a serious problem, but we will see that the 'general' automorphism group is finite (and usually small), so that the next proposition gives a reasonable estimate for the number of elements in $\mathcal{M}(r_0, H_0, F_0)/_{\simeq}$.

PROPOSITION 4. *Fix* $r_0 \in \mathbf{N}$, *a finitely generated free abelian group* H_0, *and a symmetric trilinear form* $F_0 \in S^3 H_0^{\vee}$ *which admits characteristic elements. Set* $b := rk_{\mathbf{Z}} H_0$, $s := rk_{\mathbf{Z}/2}(\bar{F}_0^t)$, *and let* $t := rk_{\mathbf{Z}/2}(\cdot_{\bar{F}_0})$ *be the* $\mathbf{Z}/_2$*-rank of the* $\mathbf{Z}/_2$*-linear square map* $\cdot_{\bar{F}_0}: \bar{H}_0 \to \bar{H}_0^{\vee}$ *sending* $\bar{u} \in \bar{H}_0$ *to* $\bar{u}^2 \in \bar{H}_0^{\vee}$. *Then* $\mathcal{M}(r_0, H_0, F_0)/_{\simeq}$ *contains at most* 2^{2b-s-t} *elements.*

Proof. Fix any admissible system of invariants $(r_0, H_0, w_0, \tau_0, F_0, p_0)$ for a manifold in $\mathcal{M}(r_0, H_0, F_0)$. Given (r_0, H_0, F_0), we know from the last lemma that the possible elements w_0 form a coset of $\mathrm{Ker}(\bar{F}_0^t)$ in $\bar{H}_0$, so that there exist precisely 2^{b-s} such elements. It remains to count the classes $[l] \in H_0^{\vee}/_{48H_0^{\vee}}/_{U_{F_0}}$, such that the $\mathrm{Aut}(F_0)$-orbit of $(w_0, [p_0 + 24T_0 + l])$ lies in the image of I.

To understand the latter condition we fix integral liftings $W_0, \in H_0$, $T_0 \in H_0^{\vee}$ of w_0 and τ_0 satisfying the admissibility conditions

i) $W_0^3 \equiv (p_0 + 24T_0)(W_0) \pmod{48}$

ii) $p_0(x) \equiv 4x^3 + 6x^2 W_0 + 3x W_0^2 \pmod{24}$ $\forall x \in H_0$.

Clearly the $\mathrm{Aut}(F_0)$-orbit of $(w_0, [p_0 + 24T_0 + l])$ lies in the image of I if and only if

i′) $W_o^3 \equiv (p_o + 24T_o + l)(W_o) \pmod{48}$,

ii′) $(p_o + l)(x) \equiv 4x^3 + 6x^2 W_o + 3x W_o^2 \pmod{24}$ $\forall x \in H_o$,

which is equivalent to $l(W_o) \equiv 0 \pmod{48}$, and $l \equiv 0 \pmod{24 H_o^\vee}$ because of i) and ii).

Now, by definition of the subgroup $U_{F_o} \subset H_o^\vee/_{48H_o^\vee}$ we have the following commutative diagram with exact rows and columns:

$$\begin{array}{ccccccc}
 & & \mathrm{Ker}(\cdot_{\bar F_o}) & & 0 & & \\
 & & \downarrow & & \downarrow & & \\
0 \to \mathrm{Ker}(24\cdot_{\bar F_o}) & \hookrightarrow & H_o/_{2H_o} & \xrightarrow{24\cdot\bar F_o} & U_{F_o} & \to & 0 \\
 & & \cdot_{\bar F_o}\downarrow & & \downarrow & & \\
0 & \to & H_o^\vee/_{2H_o^\vee} & \xrightarrow{24} & H_o^\vee/_{48H_o^\vee} & \to & H_o^\vee/_{24H_o^\vee} \to 0 \\
 & & \downarrow & & \downarrow & & \| \\
0 & \to & \mathrm{Coker}(\cdot_{\bar F_o}) & \to & H_o^\vee/_{48H_o^\vee}/_{U_{F_o}} & \to & H_o^\vee/_{24H_o^\vee} \to 0 \\
 & & \downarrow & & \downarrow & & \\
 & & 0 & & 0 & &
\end{array}$$

The number of elements $[l] \in H_o^\vee/_{48H_o^\vee}/_{U_{F_o}}$ to be counted coincides therefore with the cardinality of the kernel of the map $ev(w_o)\colon \mathrm{Coker}(\cdot_{\bar F_o}) \to \mathbf{Z}_{/2}$ induced by evaluation in w_o. This number is at most $2^{b-t}(2^{b-t-1}$ if $w_o \neq 0$ and $t \neq b$).

COROLLARY 2. *If the* $\mathbf{Z}_{/2}$*-rank* $s = rk_{\mathbf{Z}/2}(\cdot_{\bar F_o})$ *is maximal, then* $\mathscr{M}(r_o, H_o, F_o)/_{\simeq}$ *contains at most one class.*

Proof. Suppose $\cdot_{\bar F_o}\colon \bar H_o \to \bar H_o^\vee$ is surjective; then $\bar F_o^t\colon \bar H_o \to S^2 \bar H_o^\vee$ must have a trivial kernel, since $\bar h \bar x^2 = 0$ for all $\bar x \in \bar H_o$ implies $\bar h = 0$ if every linear form is a square. But this means $s = t = b$, so that $\mathscr{M}(r_o, H_o, F_o)/_{\simeq}$ has at most one element.

EXAMPLE 4. Let $H_o = \mathbf{Z}e_1 \oplus \mathbf{Z}e_2$, $e_1^3 = a$, $e_1^2 e_2 = b$, $e_1 e_2^2 = c$, $e_2^3 = d$. If $\bar b \equiv \bar c \pmod 2$, and $\bar a \bar d - \bar b \bar c \equiv 1 \pmod 2$, then $\mathscr{M}(r_o, H_o, F_o)/_{\simeq}$ contains precisely one class for every $r_o \geqslant 0$.