

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CUBIC FORMS AND COMPLEX 3-FOLDS
Autor: Okonek, Ch. / Van de Ven, A.
Kapitel: 4.1 Chern numbers of almost complex structures
DOI: <https://doi.org/10.5169/seals-61829>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 24.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

The number of integral classes in these orbits is therefore finite. We have, however, an even stronger finiteness theorem for stable ternary cubics:

PROPOSITION 7. *Let H be a free \mathbf{Z} -module of rank 3. There exist only finitely many classes of symmetric trilinear forms $F \in S^3 H^\vee$ with a fixed discriminant $\Delta \neq 0$.*

Proof. In terms of Arnhold's invariants S and T , Δ is given by $\Delta = S^3 - T^2$. By a theorem of C. Siegel [Si], the diophantine equation $S^3 - T^2 = \Delta$ has only finitely many integral solution (S, T) for any integer $\Delta \neq 0$. For each of these solutions the corresponding point in $S^3 H_C^\vee /_{SL(H_C)}$ lies outside of the discriminant curve, so that the π -fiber over it is a closed $SL(H_C)$ -orbit. The finiteness of the class number then follows from the Borel/Harish-Chandra theorem.

A famous special case of Siegel's theorem is Bachet's equation $S^3 - T^2 = 2$; it has only the two obvious solutions $(3, \pm 5)$.

REMARK 10. To get finiteness results for ternary cubic forms it is not sufficient to fix the J -invariant (instead of the discriminant): The forms $f_m = X^3 + XZ^2 + Z^3 + mY^2Z$, $m \in \mathbf{Z} \setminus \{0\}$, all have the same J -invariant, but they are not equivalent, even over \mathbf{Q} , since they have bad reduction at different primes $p \mid m$.

4. INVARIANTS OF COMPLEX 3-FOLDS

In this section we begin to investigate the topology of 1-connected, compact, complex 3-folds. After a brief discussion of the possible systems of Chern numbers of almost complex 6-manifolds, we study the behaviour of the topological invariants of complex 3-folds under certain standard constructions, like e.g. branched coverings, or blow-ups of points and curves. Then we describe some interesting examples of 1-connected, non-Kählerian 3-folds, including a new construction method which generalizes the Calabi-Eckmann manifolds. These examples will be needed in the next section in order to realize complex types of cubic forms as cup-forms of complex 3-folds.

4.1 CHERN NUMBERS OF ALMOST COMPLEX STRUCTURES

Let X be a closed, oriented, 6-dimensional differentiable manifold. The tangent bundle of X is induced by a classifying map $t_X: X \rightarrow BSO(6)$ which is unique up to homotopy. By an almost complex structure on X we mean the homotopy class $[\tilde{t}_X]$ of a lifting $\tilde{t}_X: X \rightarrow BU(3)$ of t_X to $BU(3)$.

PROPOSITION 8. *Every closed, oriented, 6-dimensional C^∞ -manifold X without 2-torsion in $H^3(X, \mathbf{Z})$ admits an almost complex structure. There is a 1-1 correspondence between almost complex structures on X and integral lifts $W \in H^2(X, \mathbf{Z})$ of $w_2(X)$. The Chern classes c_i of the almost complex manifold (X, W) are given by $c_1 = W$, $c_2 = \frac{1}{2}(W^2 - p_1(X))$.*

Proof (cf. [W]). The obstructions against lifting t_X to $BU(3)$ lie in the cohomology groups $H^{i+1}(X, \pi_i(SO(6)/_{U(3)}))$, $i = 0, 1, \dots, 5$. Since $SO(6)/_{U(3)} = \mathbf{P}^3$ has only one nontrivial homotopy group $\pi_2(SO(6)/_{U(3)}) \cong \mathbf{Z}$ in dimensions $i \leq 5$, there is in fact only one obstruction $o(t_X) \in H^3(X, \mathbf{Z})$, and this obstruction can be identified with the image of $w_2(X)$ under the Bockstein homomorphism $\beta : H^2(X, \mathbf{Z}/2) \rightarrow H^3(X, \mathbf{Z})$. Since $H^3(X, \mathbf{Z})$ has no 2-torsion by assumption, $\beta w_2(X)$ must be equal to zero, so that X has at least one almost complex structure $[\tilde{t}_X] \in [X, BU(3)]$. Standard homotopy arguments show now that the map, which assigns to an almost complex structure $[\tilde{t}_X]$ its first Chern class $\tilde{t}_X^* c_1$, induces a 1-1 correspondence between integral lifts $W \in H^2(X, \mathbf{Z})$ of $w_2(X)$ and homotopy classes of liftings of $[t_X]$ to $BU(3)$.

The second Chern class c_2 of the almost complex manifold (X, W) is determined by $W^2 - 2c_2 = p_1(X)$.

The Chern numbers $c_1^3, c_1 c_2, c_3$ of an almost complex manifold X of real dimension 6 satisfy the following congruences: $c_1^3 \equiv 0 \pmod{2}$, $c_1 c_2 \equiv 0 \pmod{24}$, $c_3 \equiv 0 \pmod{2}$. Conversely, given a triple (a, b, c) of integers $a \equiv 0 \pmod{2}$, $b \equiv 0 \pmod{24}$, and $c \equiv 0 \pmod{2}$, there always exists an almost complex manifold X of dimension 6 with Chern numbers $c_1^3 = a$, $c_1 c_2 = b$, $c_3 = c$.

It is not totally clear, however, that one can find a *connected* manifold X with prescribed Chern numbers [H1].

PROPOSITION 9. *Every triple $(a, b, c) \in \mathbf{Z}^{\oplus 3}$ satisfying $a \equiv 0 \pmod{2}$, $b \equiv 0 \pmod{24}$, $c \equiv 0 \pmod{2}$ is realizable as the Chern numbers of an almost complex 6-manifold.*

Proof. Consider the complete intersection $V(f, g) \subset \mathbf{P}^5$ defined by the polynomials $f(z) = z_0^2 + z_1^2 + 2z_2^2 - z_3^2 - z_4^2 - 2z_5^2$, and $g(z) = z_0^4 + z_1^4 + 2z_2^4 - z_3^4 - z_4^4 - 2z_5^4$ [We]. $V(f, g)$ is a singular 3-fold with 90 ordinary double points, and every small resolution V of these nodes is a (not necessarily projective) Calabi-Yau 3-fold with Euler number 4. Suppose now that a prescribed triple $(a, b, c) \in \mathbf{Z}^{\oplus 3}$ is realized by a possibly disconnected almost complex manifold $X = \coprod_{i \in I} X_i$. If we form the connected sum

X' of the X_i , we obtain a connected almost complex manifold X' with Chern numbers $c_1^3 = a$, $c_1 c_2 = b$, but with $c_3 = c - 2(|I| - 1)$.

If $|I| > 1$ take the connected sum of X' with $|I| - 1$ copies of the complex manifold V . Since V is Calabi-Yau, the Chern numbers c_1^3 and $c_1 c_2$ remain unchanged, whereas the Euler number of $X' \#_{|I|-1} V$ becomes $c_3 = c$.

REMARK 11. The above argument has been suggested by F. Hirzebruch after talk at the MPI, in which one of us had sketched a less geometric proof of the proposition.

There is another question which is related to the result above: Fix a closed, oriented, 6-dimensional differentiable manifold X . Which pairs (a, b) of integers with $a \equiv 0 \pmod{2}$ and $b \equiv 0 \pmod{24}$ occur as Chern numbers c_1^3 and $c_1 c_2$ of almost complex structures on X , and in how many ways?

For manifolds with $b_2(X) = 1$ the Chern numbers determine the almost complex structure. For manifolds with $b_2 > 1$ this is no longer true. It is possible to construct infinitely many distinct almost complex structures with the same Chern numbers on a hypersurface of bidegree $(3, 3)$ in $\mathbf{P}^2 \times \mathbf{P}^2$.

An almost complex structure $[\tilde{t}_X]$ on a differentiable 6-manifold X is said to be integrable if \tilde{t}_X is homotopic to the classifying map of a complex 3-fold. We are not aware of any example of an almost complex 6-manifold which is known not to be integrable. On the other hand, it is also unknown whether or not the Chern numbers $c_1^3, c_1 c_2$ of integrable almost complex manifold are topological invariants. The following remark might therefore be of some interest:

PROPOSITION 10. *If the Chern numbers of complex 3-folds are topological invariants, then there exist almost complex structures which are not integrable.*

Proof. Consider a closed, oriented differentiable 6-manifold X without 2-torsion in $H^3(X, \mathbf{Z})$. Fix any almost complex structure on X with first Chern class $W \in H^2(X, \mathbf{Z})$.

Every element $x \in H^2(X, \mathbf{Z})$ defines a new almost complex structure on X with first Chern class $W + 2x$, and it is easy to see that these two almost complex structures have the same Chern numbers if and only if x satisfies the equations $p_1(X) \cdot x = 0$, and $3W^2 \cdot x + 6W \cdot x^2 + 4x^3 = 0$.

Suppose now (X, W) is integrable, $p_1(X) \neq 0$, and choose $x \in H^2(X, \mathbf{Z})$ such that $p_1(X) \cdot x \neq 0$. Then clearly, either none of the almost complex manifolds $(X, W + 2x)$ is integrable, or the Chern numbers of complex 3-folds are not topologically invariant.

REMARK 12. It is very likely that there exist non-integrable almost complex structures on manifolds X as above, but probably this is hard to prove. It is also not unlikely that the Chern numbers of complex 3-folds are not topological invariants. A possible way to check this would be, to run a computer search for 3-folds given by certain standard constructions.

4.2 STANDARD CONSTRUCTIONS

For later use we investigate the topological invariants of complex 3-folds which can be obtained by certain simple standard constructions like complete intersections, simple cyclic coverings, blow-ups of points and curves, and projective bundles.

PROPOSITION 11 (Libgober/Wood). *Let $X \subset \mathbf{P}^{3+r}$ be a smooth complete intersection of multidegree $\underline{d} = (d_1, \dots, d_r)$. Choose a normalized basis $e \in H^2(X, \mathbf{Z})$, and let $\varepsilon \in H^4(X, \mathbf{Z})$ be defined by $\varepsilon(e) = 1$. Then the invariants of X are:*

$$\begin{aligned} F_X(xe) &= dx^3 \text{ where } d = \prod_{i=1}^r d_i, w_2(X) \equiv (4+r - \sum_{i=1}^r d_i)e, \\ p_1(X) &= d(4+r - \sum_{i=1}^r d_i^2)\varepsilon, \text{ and} \\ b_3(X) &= 4 - \frac{d}{6} [(4+r - \sum_{i=1}^r d_i)^3 - 3(4+r - \sum_{i=1}^r d_i)(4+r - \sum_{i=1}^r d_i^2) \\ &\quad + 2(4+r - \sum_{i=1}^r d_i^3)]. \end{aligned}$$

Proof. [L/W].

PROPOSITION 12. *Let X be a smooth, 1-connected, complex projective 3-fold, and let $\pi: X' \rightarrow X$ be a simple cyclic covering of degree d branched along a non-singular ample divisor $B \in |L^{\otimes d}|$. X' is smooth, projective, 1-connected, and $\pi^*: H^2(X, \mathbf{Z}) \rightarrow H^2(X', \mathbf{Z})$ is an isomorphism. The invariants of X and X' are related by the formulae:*

$$\begin{aligned} (\pi^*)^* F_{X'} &= dF_X, w_2(X') - \pi^* w_2(X) \equiv (d-1)\pi^* c_1(L), \\ p_1(X') - \pi^* p_1(X) &= (1-d)(1+d)\pi^* c_1(L)^2, \text{ and} \\ b_3(X') &= db_3(X) + (d-1)(b_2(B) - 2b_2(X)). \end{aligned}$$

Proof. X' is clearly smooth and projective. By a theorem of M. Cornalba $\pi: X' \rightarrow X$ is a 3-equivalence, i.e. $\pi_*: \pi_*(X') \rightarrow \pi_*(X)$ is bijective for $i \leq 2$, and surjective for $i = 3$ [Co]. X' is therefore 1-connected, and $\pi^*: H^2(X, \mathbf{Z}) \rightarrow H^2(X', \mathbf{Z})$ is an isomorphism. The relation between $F_{X'}$ and F_X is obvious, whereas the formula for $b_3(X')$ follows from $\pi_1(B) = \{1\}$ and standard properties of Euler numbers.