Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CUBIC FORMS AND COMPLEX 3-FOLDS
Autor: Okonek, Ch. / Van de Ven, A.
Kapitel: 4.2 Standard constructions
DOI: https://doi.org/10.5169/seals-61829

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 25.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

REMARK 12. It is very likely that there exist non-integrable almost complex structures on manifolds X as above, but probably this is hard to prove. It is also not unlikely that the Chern numbers of complex 3-folds are not topological invariants. A possible way to check this would be, to run a computer search for 3-folds given by certain standard constructions.

4.2 STANDARD CONSTRUCTIONS

For later use we investigate the topological invariants of complex 3-folds which can be obtained by certain simple standard constructions like complete intersections, simple cyclic coverings, blow-ups of points and curves, and projective bundles.

PROPOSITION 11 (Libgober/Wood). *Let* $X \subset \mathbf{P}^{3+r}$ *be a smooth complete intersection of multidegree* $\underline{d} = (d_1, \ldots, d_r)$. *Choose a normalized basis* $e \in H^2(X, \mathbf{Z})$, *and let* $\varepsilon \in H^4(X, \mathbf{Z})$ *be defined by* $\varepsilon(e) = 1$. *Then the invariants of* X *are:*

$$F_X(xe) = dx^3 \quad \text{where} \quad d = \prod_{i=1}^r d_i,\ w_2(X) \equiv (4 + r - \textstyle\sum_{i=1}^r d_i)e,$$

$$p_1(X) = d(4 + r - \textstyle\sum_{i=1}^r d_i^2)\varepsilon, \quad \text{and}$$

$$b_3(X) = 4 - \tfrac{d}{6}[(4 + r - \textstyle\sum_{i=1}^r d_i)^3 - 3(4 + r - \sum_{i=1}^r d_i)(4 + r - \sum_{i=1}^r d_i^2) + 2(4 + r - \sum_{i=1}^r d_i^3)] .$$

Proof. [L/W].

PROPOSITION 12. *Let* X *be a smooth, 1-connected, complex projective 3-fold, and let* $\pi : X' \to X$ *be a simple cyclic covering of degree* d *branched along a non-singular ample divisor* $B \in |L^{\otimes d}|$. X' *is smooth, projective, 1-connected, and* $\pi^* : H^2(X, \mathbf{Z}) \to H^2(X', \mathbf{Z})$ *is an isomorphism. The invariants of* X *and* X' *are related by the formulae:*

$$(\pi^*)^* F_{X'} = dF_X,\ w_2(X') - \pi^* w_2(X) \equiv (d-1)\pi^* c_1(L),$$

$$p_1(X') - \pi^* p_1(X) = (1-d)(1+d)\pi^* c_1(L)^2, \quad \text{and}$$

$$b_3(X') = db_3(X) + (d-1)(b_2(B) - 2b_2(X)) .$$

Proof. X' is clearly smooth and projective. By a theorem of M. Cornalba $\pi : X' \to X$ is a 3-equivalence, i.e. $\pi_* : \pi_i(X') \to \pi_i(X)$ is bijective for $i \leqslant 2$, and surjective for $i = 3$ [*Co*]. X' is therefore 1-connected, and $\pi^* : H^2(X, \mathbf{Z}) \to H^2(X', \mathbf{Z})$ is an isomorphism. The relation between $F_{X'}$ and F_X is obvious, whereas the formula for $b_3(X')$ follows from $\pi_1(B) = \{1\}$ and standard properties of Euler numbers.

In order to calculate $w_2(X')$ and $p_1(X')$ we compute the Chern classes of $X': c_1(X') - \pi^* c_1(X) = (1-d)\pi^* c_1(L), c_2(X') - \pi^* c_2(X) = (1-d)\pi^*[c_1(X)c_1(L) - dc_1(L)^2]$.

The latter formulae follow from the description of X' as a divisor in the total space of the line bundle L.

EXAMPLE 9. Let X be a d-fold, simple cyclic covering of $\mathbf{P}^3$ branched along a smooth surface $B \subset \mathbf{P}^3$ of degree $dl, l \geqslant 1$. Let $e \in H^2(X, \mathbf{Z})$ correspond to the preimage of a plane in $\mathbf{P}^3$. The invariants of X are then given by:

$$F_X(xe) = dx^3,\ w_2(X) \equiv (4 + (1-d)l)e, p_1(X) = d[4 + (1-d)(1+d)l^2]\varepsilon$$
$$(\varepsilon(e) = 1), b_3(X) = (d-1)(d^2l^2 - 4dl + 6)dl\ .$$

PROPOSITION 13. *Let* $\sigma: \hat{X} \to X$ *be the blow-up of a complex 3-fold* X *in a point, and let* $e \in H^2(\hat{X}, \mathbf{Z})$ *be the class of the exceptional divisor. The invariants of* $\hat{X}$ *and* X *are related by the following formulae:*

$$F_{\hat{X}}(\sigma^* h + xe) = F_X(h) + x^3\ \forall\ h \in H^2(X, \mathbf{Z}), x \in \mathbf{Z}, w_2(\hat{X}) = \sigma^* w_2(X),$$
$$p_1(\hat{X}) = \sigma^* p_1(X) + 4(e^2 - \sigma^* c_1(X) \cdot e), b_3(\hat{X}) = b_3(X)\ .$$

Proof. Standard arguments, see [G/H]. The Chern classes are related by $c_1(\hat{X}) = \sigma^* c_1(X) - 2e, c_2(\hat{X}) = \sigma^* c_2(X)$.

PROPOSITION 14. *Let* $\sigma: \hat{X} \to X$ *be the blow-up of a complex 3-fold* X *along a smooth curve* C *of genus* g, *and let* $e \in H^2(\hat{X}, \mathbf{Z})$ *be the class of the exceptional divisor. The invariants of* $\hat{X}$ *and* X *are related by:*

$$F_{\hat{X}}(\sigma^* h + xe) = F_X(h) - 3h \cdot Cx^2 - \deg N_{C/X} x^3\ \forall\ h \in H^2(X, \mathbf{Z}),$$
$$x \in \mathbf{Z}, w_2(\hat{X}) \equiv \sigma^* w_2(X) + e, p_1(\hat{X}) = \sigma^* p_1(X) + (e^2 - 2\sigma^* C),$$
$$b_3(\hat{X}) = b_3(X) + 2g\ .$$

Proof. [*G/H*]. The Chern classes are given by $c_1(\hat{X}) = \sigma^* c_1(X) - c, c_2(\hat{X}) = \sigma^*(c_2(X) + C) - \sigma^* c_1(X) \cdot e$.

PROPOSITION 15. *Let* E *be a holomorphic vector bundle of rank 2 with Chern classes* $c_i(E), i = 1, 2$ *over a 1-connected, compact complex surface* Y, *and let* $\pi: \mathbf{P}(E) \to Y$ *be the projective bundle of lines in the fibers of* E. *The cup-form of* $\mathbf{P}(E)$ *is given by*

$$F_{\mathbf{P}(E)}(h + x\xi) = x[(3h^2) - (3c_1(E) \cdot h)x + (c_1(E)^2 - c_2(E))x^2]\ ,$$

where $\xi = c_1(\mathscr{O}_{\mathbf{P}(E)}(1))$, $h \in H^2(Y, \mathbf{Z})$, *and* $x \in \mathbf{Z}$. *The other topological invariants of* $\mathbf{P}(E)$ *are:*

$$w_2(\mathbf{P}(E)) \equiv \pi^*(w_2(Y) + c_1(E)), p_1(E))$$
$$= \pi^*[c_1(Y)^2 - 2c_2(Y) + c_1(E)^2 - 4c_2(E)],\ b_3(\mathbf{P}(E)) = 0\ .$$

Proof. The Leray-Hirsch theorem identifies the cohomology ring $H^*(\mathbf{P}(E), \mathbf{Z})$ with the ring $H^*(Y, \mathbf{Z})[\xi]/_{<\xi^2 + c_1(E)\cdot\xi + c_2(E)>}$; this determines the cup-form. In order to calculate the characteristic classes one uses the exact sequence $0 \to \mathscr{O}_{\mathbf{P}(E)} \to \pi^*E \otimes \mathscr{O}_{\mathbf{P}(E)}(1) \to T_{\mathbf{P}(E)} \to \pi^*T_Y \to 0$. $b_3(\mathbf{P}(E)) = 0$ follows from $b_1(Y) = 0$ and the Leray-Hirsch theorem.

4.3 Examples of 1-connected non-Kählerian 3-folds

Recall that the Hessian of a symmetric trilinear form $F \in S^3H^\vee$ on a free $\mathbf{Z}$-module H of finite rank was defined as the composition $H_F: H \xrightarrow{F^t} S^2H^\vee \xrightarrow{\text{disc}} \mathbf{Z}$. In terms of coordinates $\xi_1, \ldots, \xi_b$ on H it is given by the determinant $\det\left(\frac{\partial^2 f}{\partial\xi_i\partial\xi_j}\right)$, where $f \in \mathbf{C}[H_{\mathbf{C}}]_3$ is the homogeneous cubic polynomial associated with F.

Proposition 16. *Let* F *be a symmetric trilinear form whose Hessian vanishes identically. Then* F *is not realizable as cup-form of a Kählerian 3-fold.*

Proof. Let X be a complex 3-fold with a Kähler metric g. The Kähler class $[\omega_g] \in H^2(X, \mathbf{R})$ defines a multiplication map $\cdot[\omega_g]: H^2(X, \mathbf{R}) \to H^4(X, \mathbf{R})$, which is an isomorphism by the Hard Lefschetz Theorem [G/H]. In section 3.1 we have seen that this is not possible if the Hessian of the cup-form vanishes.

Corollary 6. *Cubic forms* $f \in \mathbf{C}[H_{\mathbf{C}}]_3$ *which depend on strictly less than* $b = rk_{\mathbf{Z}}H$ *variables are not realizable as cup-forms of Kählerian 3-folds with* $b_2 = b$.

By considering the Hessian of a cup-form over the reals one obtains further conditions.

Definition 4. *Let* $F \in S^3H^\vee$ *be a symmetric trilinear form on a free* $\mathbf{Z}$*-module of rank* b.

The Hesse cone of F *is the subset* $\mathscr{H}_F \subset H_{\mathbf{R}}$ *defined by* $\mathscr{H}_F := \{h \in H_{\mathbf{R}} \mid (-1)^b \det(F^t(h)) < 0\}$.