Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	41 (1995)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	CUBIC FORMS AND COMPLEX 3-FOLDS
Autor:	Okonek, Ch. / Van de Ven, A.
Kapitel:	4.3 EX AMPLES OF 1-CONNECTED NON-KÄHLERIAN 3-FOLDS
DOI:	https://doi.org/10.5169/seals-61829

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

where $\xi = c_1(\mathscr{O}_{\mathbf{P}(E)}(1)), h \in H^2(Y, \mathbb{Z}), and x \in \mathbb{Z}$. The other topological invariants of $\mathbf{P}(E)$ are:

$$w_2(\mathbf{P}(E)) \equiv \pi^*(w_2(Y) + c_1(E)), p_1(E))$$

= $\pi^*[c_1(Y)^2 - 2c_2(Y) + c_1(E)^2 - 4c_2(E)], b_3(\mathbf{P}(E)) = 0.$

Proof. The Leray-Hirsch theorem identifies the cohomology ring $H^*(\mathbf{P}(E), \mathbf{Z})$ with the ring $H^*(Y, \mathbf{Z})[\xi]/_{\langle\xi^2+c_1(E)\cdot\xi+c_2(E)\rangle}$; this determines the cup-form. In order to calculate the characteristic classes one uses the exact sequence $0 \to \mathscr{O}_{\mathbf{P}(E)} \to \pi^*E \otimes \mathscr{O}_{\mathbf{P}(E)}(1) \to T_{\mathbf{P}(E)} \to \pi^*T_Y \to 0$. $b_3(\mathbf{P}(E)) = 0$ follows from $b_1(Y) = 0$ and the Leray-Hirsch theorem.

4.3 Examples of 1-connected non-Kählerian 3-folds

Recall that the Hessian of a symmetric trilinear form $F \in S^3 H^{\vee}$ on a free **Z**-module *H* of finite rank was defined as the composition $H_F: H \xrightarrow{F^l} S^2 H^{\vee} \xrightarrow{\text{disc}} \mathbf{Z}$. In terms of coordinates $\xi_1, ..., \xi_b$ on *H* it is given by the determinant det $\left(\frac{\partial^2 f}{\partial \xi_i \partial \xi_j}\right)$, where $f \in \mathbb{C}[H_{\mathbb{C}}]_3$ is the homogeneous cubic polynomial associated with *F*.

PROPOSITION 16. Let F be a symmetric trilinear form whose Hessian vanishes identically. Then F is not realizable as cup-form of a Kählerian 3-fold.

Proof. Let X be a complex 3-fold with a Kähler metric g. The Kähler class $[\omega_g] \in H^2(X, \mathbb{R})$ defines a multiplication map $\cdot [\omega_g]: H^2(X, \mathbb{R}) \to H^4(X, \mathbb{R})$, which is an isomorphism by the Hard Lefschetz Theorem [G/H]. In section 3.1 we have seen that this is not possible if the Hessian of the cup-form vanishes.

COROLLARY 6. Cubic forms $f \in \mathbb{C}[H_{\mathbb{C}}]_3$ which depend on strictly less than $b = rk_{\mathbb{Z}}H$ variables are not realizable as cup-forms of Kählerian 3-folds with $b_2 = b$.

By considering the Hessian of a cup-form over the reals one obtains further conditions.

DEFINITION 4. Let $F \in S^3 H^{\vee}$ be a symmetric trilinear form on a free **Z**-module of rank b.

The Hesse cone of F is the subset $\mathcal{H}_F \subset H_R$ defined by $\mathcal{H}_F := \{h \in H_R \mid (-1)^b \det (F^t(h)) < 0\}.$

The index cone \mathscr{J}_F of F is the subset $\mathscr{J}_F := \{h \in \mathscr{H}_F | F^t(h) \in S^2 H_{\mathbb{R}}^{\vee}$ has signature $(1, -1, ..., -1)\}.$

Clearly \mathscr{J}_F is an open subcone of \mathscr{H}_F which coincides with \mathscr{H}_F iff $b \leq 2$.

THEOREM 5. Let $F_X \in S^3 H^2(X, \mathbb{Z})^{\vee}$ be the cup-form of a smooth projective 3-fold with $h^{0,2}(X) = 0$. Then F_X has a non-empty index cone.

Proof. Let $h \in H^2(X, \mathbb{Z})$ be the dual class of a hyperplane section Y in some projective embedding. The inclusion $i: Y \hookrightarrow X$ induces a monomorphism $i^*: H^2(X, \mathbb{Z}) \to H^2(Y, \mathbb{Z})$ by the weak Lefschetz theorem. The symmetric bilinear form $F_X^t(h) \in S^2 H^2(X, \mathbb{Z})^{\vee}$ is simply the pull-back of the cup-form of Y under the inclusion i^* ; it is therefore non-degenerate by the Hard Lefschetz theorem [L]. Applying the Hodge index theorem to Y we see that the real bilinear form $F_X^t(h) \in S^2 H^2(X, \mathbb{R})^{\vee}$ must have one positive and b - 1 negative eigenvalues. In other words: $h \in I_{F_X}$.

REMARK 13. This result has two applications: it provides topological 'upper bounds' for the ample cone of a projective 3-fold with $h^{0,2} = 0$, and if gives further restrictions on symmetric trilinear forms to be realizable as cup-forms of projective 3-folds with $h^{0,2} = 0$ if $b \ge 4$.

These applications will be discussed in section 5.

We will now describe examples of 1-connected, non-Kählerian, complex 3-folds and determine their topological structure.

EXAMPLE 10 (Calabi-Eckmann). E. Calabi and B. Eckmann have defined complex structures X_{τ} , depending on a parameter τ , on the product $S^3 \times S^3[C/E]$. Their manifolds are principal fiber bundles over $\mathbf{P}^1 \times \mathbf{P}^1$ whose fiber and structure group is the elliptic curve $E_{\tau} = \mathbf{C}/_{\mathbf{Z} \oplus \mathbf{Z} \tau}$, Im $(\tau) > 0$.

The Calabi-Eckmann manifolds are homogeneous, non-Kählerian 3-folds of algebraic dimension 2.

EXAMPLE 11 (Maeda). H. Maeda has generalized the Calabi-Eckmann construction. He constructed fiber bundles X'_{τ} over Hirzebruch surfaces $\mathbf{F}_n, n \ge 0$, whose fiber and structure group are an elliptic curve E_{τ} and Aut (E_{τ}) respectively [M]. X'_{τ} is again diffeomorphic to $S^3 \times S^3$, and therefore non-Kählerian. Maeda's manifolds X'_{τ} are homogeneous if and only if n = 0 in which case they are Calabi-Eckmann 3-folds. The Calabi-Eckmann construction can also be generalized in the following way:

Let $S^2 \times S^4$ be the non-trivial S^4 -bundle over S^2 , i.e. $S^2 \times S^4$ is the unique 1-connected, closed, oriented, differentiable 6-manifold with $H_2(S^2 \times S^4, \mathbb{Z}) \cong \mathbb{Z}$ and $b_3 = 0$, whose cup-form and Pontrjagin class vanish, but whose Stiefel-Whitney class w_2 is non-zero.

THEOREM 6. For any integer $b \ge 0$ there exist compact complex 3-folds X_b , and $X_{\tilde{b}}$ if $b \ge 1$, which are homeomorphic to $\#_b S^2 \times S^4 \#_{b+1} S^3 \times S^3$, and $S^2 \tilde{\times} S^4 \#_{b-1} S^2 \times S^4 \#_{b+1} S^3 \times S^3$.

Proof. Let Y be a 1-connected, compact complex surface with $p_g(Y) = 0$ and $b_2(Y) \ge 2$, and let $E = \mathbb{C}/_{\Gamma}$ be the elliptic curve associated to the lattice $\Gamma \subset \mathbb{C}$. We want to construct the required 3-folds as total spaces of principal E-bundles over Y. Let $c: H_2(Y, \mathbb{Z}) \to \Gamma$ be an arbitrary epimorphism. The corresponding cohomology class $c \in H^2(Y, \Gamma)$ defines a topological principal bundle over Y with fiber and structure group $E = \mathbb{C}/_{\Gamma}$ as follows immediately from the identification of the classifying space $BE \simeq K(\Gamma, 2)$.

Let $\mathscr{O}_Y(E)$ be the sheaf of germs of holomorphic maps from Y to E. We have a short exact sequence $0 \to \Gamma \to \mathscr{O}_Y \to \mathscr{O}_Y(E) \to 0$ and a corresponding exact cohomology sequence

$$\rightarrow H^1(Y, \mathscr{O}_Y) \rightarrow H^1(Y, \mathscr{O}_Y(E)) \stackrel{\circ}{\rightarrow} H^2(Y, \Gamma) \rightarrow H^2(Y, \mathscr{O}_Y) \rightarrow$$

By our assumptions δ is an isomorphism, so that every topological principal *E*-bundle admits a holomorphic structure. Let *X* be the total space of such a bundle corresponding to a surjective map $c: H_2(Y, \mathbb{Z}) \to \Gamma$. The homotopy sequence of the fibration $p: X \to Y$ yields the sequence

$$0 \to \pi_2(X) \stackrel{p_*}{\to} \pi_2(Y) \to \pi_1(E) \to \pi_1(X) \stackrel{p_*}{\to} \pi_1(Y) \to 0 .$$

Since Y is 1-connected, $\pi_2(Y)$ can be identified with $H_2(Y, \mathbb{Z})$, and then the boundary map $\pi_2(Y) \to \pi_1(E)$ becomes the characteristic map $c: H_2(Y, \mathbb{Z}) \to \Gamma$ of the bundle. This implies $\pi_1(X) = \{1\}$, whereas $H_2(X, \mathbb{Z})$ is given by: $0 \to H_2(X, \mathbb{Z}) \xrightarrow{p_*} H_2(Y, \mathbb{Z}) \xrightarrow{s} \Gamma \to 0$.

In particular, $H_2(X, \mathbb{Z})$ is free as a submodule of $H_2(Y, \mathbb{Z})$, and by dualizing the last sequence we obtain an identification (via p^*)

$$H^2(X, \mathbf{Z}) = H^2(Y, \mathbf{Z})/_{\Gamma^{\vee}}$$
.

The cup-form F_X of X is therefore trivial. In order to calculate $p_1(X)$ and $w_2(X)$, we use the exact sequence of tangent sheaves: $0 \to T_{X/Y} \to T_X$ → $p^*T_Y \rightarrow 0$. Since $T_{X/Y}$ is a trivial bundle, the characteristic classes of X are simply the pullbacks of the corresponding classes of Y. But the map $p^*: H^4(Y, \mathbb{Z}) \rightarrow H^4(X, \mathbb{Z})$ is zero, since $\langle p^*(\varepsilon) \cup p^*(\alpha), [X] \rangle$ = $\langle \varepsilon \cup \alpha, p_*[X] \rangle = 0$ for all classes $\varepsilon \in H^4(Y, \mathbb{Z})$, and $\alpha \in H^2(Y, \mathbb{Z})$.

Thus $p_1(X) = 0$, and $w_2(X)$ is the residue of $w_2(Y) \in H^2(Y, \mathbb{Z}_{/2})$ modulo $\Gamma^{\vee}/_{2\Gamma^{\vee}}$.

The Euler characteristic of X is zero, so that from $b_2(X) = b_2(Y) - 2$ we find $b_3(X) = 2(b_2(Y) - 1)$. The system of invariants associated to the manifold X is therefore given by

$$(b_2(Y) - 1, H^2(Y, \mathbb{Z})/_{\Gamma^{\vee}}, w_2(Y) \pmod{\Gamma^{\vee}/_{2\Gamma^{\vee}}}, 0, 0, 0),$$

i.e. X is diffeomorphic to

 $\#_{b_2(Y)-2}S^2 \times S^4 \#_{b_2(Y)-1}S^3 \times S^3 \text{ if } w_2(Y) \in \Gamma^{\vee}/_{2\Gamma^{\vee}},$

and to $S^2 \tilde{\times} S^4 \#_{b_2(Y)-3} S^2 \times S^4 \#_{b_2(Y)-1} S^3 \times S^3$ if $b_2(Y) \ge 3$, and $w_2(Y) \notin \Gamma^{\vee}/_{2\Gamma^{\vee}}$.

EXAMPLE 12 (Kato). In the two papers [K1], [K2] M. Kato studies the class of compact, complex 3-folds X containing smooth rational curves with neighborhoods biholomorphic to those of projective lines in \mathbf{P}^3 . On this class of 3-folds, called class L, he defines a semi-group structure + with neutral element \mathbf{P}^3 .

Kato's connecting operation + is defined by removing 'lines' $L_i \,\subset X_i$ from 3-folds X_i , i = 1, 2, and by identifying the complements $X_i \setminus L_i$ along open sets $U_i \setminus L_i$ obtained from suitable neighborhoods $U_i \subset X_i$.

Starting with a certain elliptic fiber space X_1 over the blow-up of $\mathbf{P}^1 \times \mathbf{P}^1$ in a point, he constructs a sequence of 3-folds $X_n := X_1$ $+ X_{n-1}, n \ge 2$. The 3-folds X_n are 1-connected spin-manifolds with $H_2(X_n, \mathbf{Z}) = \mathbf{Z}$. Their cup-forms F_{X_n} , and their Pontrjagin classes $p_1(X_n)$ are in terms of a (normalized) generator $e_n \in H^2(X_n, \mathbf{Z})$ and its dual class $\varepsilon_n \in H^4(X_n, \mathbf{Z})$ given by $F_{X_n}(xe_n) = (n-1)x^3$, and $p_1(X_n)$ $= 4(n-1)\varepsilon_n (\varepsilon_n(e_n) = 1)$. The third Betti-number of X_n is 4n.

In particular, X_1 is diffeomorphic to $S^2 \times S^4 \#_2 S^3 \times S^3$, and X_2 is diffeomorphic to $\mathbf{P}^3 \#_4 S^3 \times S^3$. It is interesting to note that the Chernnumbers c_1^3 , $c_1 c_2$ of the X_n are $c_1^3 = 64(1 - n)$, $c_1 c_2 = 24(1 - n)$, i.e. they satisfy $8c_1 c_2 = 3c_1^3$. For projective manifolds of general type this equality is characteristic for ball quotients [Y]. EXAMPLE 13 (Twistor spaces). Let $p: Z \to M$ be the twistor fibration of a closed, oriented Riemannian 4-manifold (M, g). Z carries a natural almost complex structure which is integrable if and only if g is self-dual [A/H/S].

Examples of 1-connected 4-manifolds which admit self-dual structures are S^4 , $\#_n \mathbf{P}^2$, and K3-surfaces.

The total spaces of their twistor fibrations are 1-connected complex 3-folds which may be Moishezon for S^4 and $\#_n \mathbf{P}^2$ [C], but which are usually non-Kähler [Hi]. We leave it to the reader to calculate the topological invariants of these 3-folds. There is an interesting relation between Twistor spaces of connected sums and Kato's connection operation + for class L manifolds [K2], [D/F].

EXAMPLE 14 (Oguiso). In a recent preprint [O1] K. Oguiso constructs examples of 1-connected, Moishezon Calabi-Yau 3-folds with very interesting cup-forms. He proves that for every integer $d \ge 1$ there exists a smooth complete intersection X'_d of type (2, 4) in \mathbf{P}^5 which contains a non-singular rational curve C_d of degree d with normal bundle $N_{C_d/X_d} = \mathscr{O}_{C_d}(-1)^{\oplus 2}$.

The 3-fold X'_d can now be flopped along C_d , i.e. C_d can be blown up to $\mathbf{P}(N_{C_d/X_d}) \cong \mathbf{P}^1 \times \mathbf{P}^1$, and then 'blown down in the other direction'. The resulting 3-fold X_d is a 1-connected Moishezon manifold with trivial canonical bundle and cup-form F_{X_d} given by $F_{X_d}(xe_d) = (d^3 - 8)x^3$. Here $e_d \in H^2(X_d, \mathbb{Z})$ is the normalized generator corresponding to the strict transform of the negative of a hyperplane section of X'_d . The Pontrjagin class of X_d is $p_1(X_d) = (112 + 4d)\varepsilon_d$ where $\varepsilon_d \in H^4(X_d, \mathbb{Z})$ denotes the generator with $\varepsilon_d(e_d) = 1$. Since the Euler-number does not change under a flop we have $b_3(X_d) = 180$ for every d.

5. Complex 3-folds with small b_2

In this section we investigate the following natural problem: Which cubic forms can be realized as cup-forms of compact complex 3-folds? For small b_2 something can be said: Any core of a 1-connected, closed, oriented differentiable 6-manifold with $H_2(X, \mathbb{Z}) \cong \mathbb{Z}$ is homotopy equivalent to the core of a 1-connected complex 3-fold. In the case $b_2 = 2$, at least every discriminant Δ is realizable by a complex manifold. If $b_2 = 3$ we can realize all types of complex cubics with one exception, the union of a smooth conic and a tangent line. In addition to these realization results we prove a finiteness