

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CUBIC FORMS AND COMPLEX 3-FOLDS
Autor: Okonek, Ch. / Van de Ven, A.
Kapitel: 5.1 3-folds with $b_2 = 1$
DOI: <https://doi.org/10.5169/seals-61829>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 01.05.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

theorem for 3-folds with $b_2 = 1$, $w_2 \neq 0$, and we give examples which show that the condition $I_{F_X} \neq \emptyset$ for the index cone of a projective 3-fold with $h^{0,2} = 0$ is non-trivial in general.

5.1 3-FOLDS WITH $b_2 = 1$

Recall from section 1.1 that every closed, oriented, 1-connected differentiable 6-manifold X with torsion-free homology has a connected sum decomposition $X \cong X_\circ \#_r S^3 \times S^3$ where $r = \left(\frac{b_3(X)}{2}\right)$, which is unique up to orientation preserving diffeomorphisms; the manifold X_\circ with $b_3(X_\circ) = 0$ is the core of X .

THEOREM 6. *Let X_\circ be a 1-connected, closed, oriented differentiable 6-manifold with $H_2(X_\circ, \mathbf{Z}) \cong \mathbf{Z}$ and $b_3(X_\circ) = 0$. There exists a compact complex 3-fold X whose core is orientation preservingly homotopy equivalent to X_\circ .*

Proof. The oriented homotopy type of X_\circ is determined by the invariants d , w_2 , and $p_1 \pmod{48}$; more precisely: for $d \equiv 1 \pmod{2}$ there is a single homotopy type whereas for $d \equiv 0 \pmod{2}$ there are three; one of these 3 types has $w_2 \neq 0$, the other two are spin, they are distinguished by $p_1 \equiv 4d \pmod{48}$, $p_1 \equiv 4d + 24 \pmod{48}$ respectively. In order to realize these homotopy types as cores of complex 3-folds we first look at simple cyclic coverings of \mathbf{P}^3 . Given a positive integer d , let $\pi: X \rightarrow \mathbf{P}^3$ be a simple cyclic covering of \mathbf{P}^3 branched along a smooth surface B of degree dl . Then X has the correct ‘degree’ d and the characteristic classes $w_2 \equiv (d-1)l \pmod{2}$, and $p_1 = 4d + (1-d)(1+d)dl^2$, see 4.2. For odd d there is nothing to prove. For even d we can realize $w_2 = 0$ or $w_2 \neq 0$ by choosing $l \equiv 0 \pmod{2}$ or $l \equiv 1 \pmod{2}$. Taking $l \equiv 0 \pmod{4}$ gives $w_2 = 0$, $p_1 \equiv 4d \pmod{48}$, taking $l \equiv 2 \pmod{4}$ yields $w_2 = 0$, and $p_1 \equiv 4d + 24 \pmod{48}$. It remains to treat the special case $d = 0$, where the 3 homotopy types are given by $w_2 \neq 0$, by $w_2 = 0$, $p_1 \equiv 0 \pmod{16}$, and by $w_2 = 0$, $p_1 \equiv 8 \pmod{16}$. The first two homotopy types are realizable as cores of elliptic fiber bundles over the projective plane blown up in two points.

The third homotopy type is realized by the core of Oguiso’s Calabi-Yau 3-fold X_2 with vanishing cup-form and $p_1(X_2) = 120\varepsilon_2$.

The result just proven suggests a natural question: given a manifold X_\circ as above, which (even) integers $b_3 \geq 0$ occur as the third Betti numbers of complex 3-folds X whose core is homotopy equivalent to X_\circ ?

There will certainly be some gaps for algebraic 3-folds. In order to show this, we prove the following finiteness theorem for families of Kähler structures:

THEOREM 7. *Fix a positive constant c . There exist only finitely many families of 1-connected, smooth projective 3-folds X with $H_2(X, \mathbf{Z}) \cong \mathbf{Z}$, $w_2(X) \neq 0$, and with $b_3(X) \leq c$.*

Proof. Let X be a smooth projective 3-fold with $H_1(X, \mathbf{Z}) = \{0\}$, $H_2(X, \mathbf{Z}) \cong \mathbf{Z}$, and with $w_2(X) \neq 0$. Clearly $\text{Pic}(X) \cong H^2(X, \mathbf{Z})$, and we can choose a basis $e \in H^2(X, \mathbf{Z})$ corresponding to the ample generator of $\text{Pic}(X)$.

Let $c_1(X) = c_1 e$, $c_2(X) = c_2 \varepsilon$ where $e^2 = d\varepsilon$, $\varepsilon(e) = 1$. If c_1 is positive, then X is Fano, and there are only finitely many possibilities [Mu]. The case $c_1 = 0$ is excluded, so that we are left with $c_1 < 0$, i.e. the canonical bundle of X is ample.

The Riemann-Roch formula $\chi(X, \mathcal{O}_X) = 1 - h^3(X, \mathcal{O}_X) = \frac{1}{24} c_1 c_2$ shows that the set of possible Chern numbers $c_1 c_2$ is bounded from below: $24(1 - c) \leq c_1 c_2$. Using Yau's inequality $8c_1(X)c_2(X) \leq 3c_1(X)^3$ we find that $d | c_1 |^3 \leq 64(c - 1)$, i.e. the degree d and the order of divisibility $|c_1|$ of $c_1(X)$ is bounded. Now Kollar's finiteness theorem [Ko2] yields the assertion.

EXAMPLE 15. Let X be a 1-connected, smooth projective 3-fold with $H_2(X, \mathbf{Z}) \cong \mathbf{Z}$ and $w_2(X) \neq 0$. If $b_3(X) \leq 2$, then $h^3(X, \mathcal{O}_X) \leq 1$ and X must be Fano of index 1 or 3. For $b_3(X) = 4$ we have that X is either Fano, or $h^3(X, \mathcal{O}_X) = 2$ and X is of general type with $d | c_1 |^3 \leq 64$.

Note that the assumption $w_2 \neq 0$ was only used to exclude Calabi-Yau 3-folds.

5.2 3-FOLDS WITH $b_2 = 2$

Let X be a 1-connected, closed, oriented, 6-dimensional differentiable manifold with $H_2(X, \mathbf{Z}) \cong \mathbf{Z}^2$.

We choose a basis e_1, e_2 for $H^2(X, \mathbf{Z})$ and set $a_0 = e_1^3$, $a_1 = e_1^2 e_2$, $a_2 = e_1 e_2^2$, $a_3 = e_2^3$; the cubic polynomial f associated to the cup-form of X is then given by $f = a_0 X^3 + 3a_1 X^2 Y + 3a_2 X Y^2 + a_3 Y^3$. The discriminant of f is by definition $\Delta(f) = a_0^2 a_3^2 - 3a_1^2 a_2^2 - 6a_0 a_1 a_2 a_3 + 4a_0 a_2^3 + 4a_1^3 a_3$; up to a factor it is simply the discriminant of the Hessian $H_f = 6^2[(a_0 a_2 - a_1^2)X^2 + (a_0 a_3 - a_1 a_2)XY + (a_1 a_3 - a_2^2)Y^2]$ of f : $\Delta(f) = (a_0 a_3 - a_1 a_2)^2 - 4(a_0 a_2 - a_1^2)(a_1 a_3 - a_2^2)$.