

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CUBIC FORMS AND COMPLEX 3-FOLDS
Autor: Okonek, Ch. / Van de Ven, A.
Kapitel: 5.3 3-folds with $b_2 \geq 3$
DOI: <https://doi.org/10.5169/seals-61829>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

REMARK 14. The Hessian of a binary form $F \in S^3 H^\vee$ is identically zero iff F is degenerate; it is negative semi-definite if F is non-degenerate and $\Delta(F) \leq 0$; it is indefinite iff $\Delta(F) > 0$ [Ca]. Only in the indefinite case $\Delta(F) > 0$ can the closure $\overline{\mathcal{H}}_F := \{h \in H_{\mathbf{R}} \mid \det F'(h) \leq 0\}$ of the Hesse cone be a proper subset of $H_{\mathbf{R}}$.

EXAMPLE 16. Let $P = \mathbf{P}_{\mathbf{P}^2}(E)$ be the projectivization of a rank-2 vector bundle E with Chern classes $c_i = c_i(E)$. The cup-form of P yields the cubic polynomial $f = (c_1^2 - c_2)X^2 + 3(-c_1)X^2Y + 3XY^2$ whose Hessian is $H_f = (-c_2)X^2 + c_1XY - Y^2$. Rewriting H_f as $H_f = -\frac{1}{4}[(2Y - c_1X)^2 + X^2(4c_2 - c_1^2)] = \frac{-1}{4}[(2Y - c_1X)^2 - \Delta(f)X^2]$ we find 3 possibilities for the Hesse cone:

- i) $\Delta(f) < 0$: $\mathcal{H}_f = H^2(P, \mathbf{R}) \setminus \{0\}$
- ii) $\Delta(f) = 0$: $\mathcal{H}_f = H^2(P, \mathbf{R}) \setminus L_{c_1}$ for a real line L_{c_1} depending on c_1 ($L_{c_1} = \mathbf{R}(2, c_1)$ in the coordinates X, Y)
- iii) $\Delta(f) > 0$: \mathcal{H}_f is an open cone whose angle is determined by $\Delta(f) ((Z + \sqrt{\Delta(f)}X)(Z - \sqrt{\Delta(f)}X) > 0$ in coordinates $X, Z := 2Y - c_1X$).

5.3 3-FOLDS WITH $b_2 \geq 3$

Let X be a 1-connected, compact complex 3-fold with $H_2(X, \mathbf{Z}) \cong \mathbf{Z}^{\oplus 3}$. The cup-form of X gives rise to a curve C_X of degree 3 in the projective plane $\mathbf{P}(H^2(X, \mathbf{C}))$:

$$C_X := \{ \langle h \rangle \in \mathbf{P}(H^2(X, \mathbf{C})) \mid h^3 = 0 \}.$$

A first natural question is which types of plane cubic curves occur in this way?

Recall that there are 10 types of plane cubics, namely: 1) non-singular cubics, 2) irreducible cubics with a node, 3) irreducible cubics with a cusp, 4) reducible cubics consisting of a smooth conic and a transversal line, 5) smooth conics with a tangent line, 6) three lines forming a triangle, 7) three distinct lines through a common point, 8) a double line with a third skew line, 9) a triple line, 10) the trivial 'cubic' with equation 0.

LEMMA 4. *If the 3-fold X has a non-trivial Hodge number $h^{2,0}(X) \neq 0$, then C_X is of type 4), 6) 9) or 10).*

Proof. Choose basis vectors $e^{k,l} \in H^{k,l}(X)$, so that every $h \in H^2(X, \mathbf{C})$ can be uniquely written as $h = xe^{2,0} + ye^{1,1} + ze^{0,2}$.

Then clearly $h^3 = y[y^2(e^{1,1})^3 + 6xz(e^{2,0} \cdot e^{1,1} \cdot e^{0,2})]$.

We now realize the cubics of types 7)-10). These cubics are degenerate, i.e. they are cones, and therefore their Hessians vanish identically. From section 4.3 we know that they can not be realized by Kählerian 3-folds.

PROPOSITION 20. *The plane cubics of types 7)-10) can all be realized by 1-connected, non-Kählerian 3-folds.*

Proof. ‘Cubics’ of type 10) can be realized by elliptic fibre bundles over surfaces Y with $b_2(Y) = 5$. In order to realize cubics of type 9) or 7) one blows up one or two points in an elliptic fibre bundle over a surface with $b_2 = 4$ or 3 respectively. The realization of a type 8) cubic is a little trickier: One starts with an elliptic fibre bundle over a surface Y with $b_2(Y) = 3$, and blows up one of its fibers. The resulting 3-fold X' has $b_2(X') = 2$ and $F_{X'} \equiv 0$. Now choose a line l in the exceptional divisor E of X' , and let X be the blow-up of X' along l . The cup-form of X yields the cubic polynomial $x^2[y(-3l \cdot E) - x(\deg N_{C/X'})]$ with a non-zero coefficient $-3l \cdot E = 3$.

There are four types of complex cubics which we have been able to realize by projective 3-folds.

PROPOSITION 21. *Cubics of type 1), 3), 4) and 6) are realizable by 1-connected projective 3-folds.*

Proof. Type 1) occurs for blow-ups of complete intersections in two distinct points. The product $\mathbf{P}^1 \times \mathbf{P}^1 \times \mathbf{P}^1$ realizes a triangle, whereas most projective bundles over a surface with $b_2 = 2$ lead to the union of a smooth conic and a transversal line.

Irreducible cubics with a cusp can be obtained by blowing-up a line and a point in \mathbf{P}^3 . The resulting 3-fold yields the cubic polynomial $X^3 - 3XY^2 - 2Y^3 + Z^3 = (X + Y)^2(X - 2Y) + Z^3$.

The remaining two types of cubics are cubics with a node (type 2)), and smooth conics with a tangent line (type 5)). We do not know if these types are realizable by projective 3-folds. A non-Kählerian 3-fold whose cup-form yields a nodal cubic can be constructed: one just takes the blow-up of two suitable curves in Oguiso’s Calabi-Yau 3-fold with $b_2 = 1$ and vanishing cup-form.

Finally we like to show that the non-emptiness condition on the index cone of a projective 3-fold with $h^{0,2} = 0$ gives non-trivial restrictions for the possible cup-forms if $b_2 \geq 4$. Further investigations of this condition will appear elsewhere [Sch].

EXAMPLE 17. Let H be a free \mathbf{Z} -module of rank 4 with basis $(e_i)_{i=1,\dots,4}$. Consider a trilinear form $F \in S^3 H^\vee$ and its adjoint map $F^t: H \rightarrow S^2 H^\vee$. The image $F^t(h)$ of an element $h \in H$ is in terms of the chosen basis $(e_i)_{i=1,\dots,4}$ represented by the symmetric 4×4 -matrix $[[he_i e_j]]_{i,j=1,\dots,4}$. Suppose this matrix is a diagonal sum $[[he_i e_j]]_{i,j=1,2} \oplus [[he_k e_l]]_{k,l=3,4}$ such that the determinants of both 2×2 -matrices are negative for every $h \in H \setminus \{0\}$.

In this case $F^t(h)$ were of signature $(1, -1, 1, -1)$ for every $h \in H \setminus \{0\}$, and we would have $I_F = \mathcal{H}_F = \emptyset$.

All these conditions can be met, e.g. by setting $e_1^2 e_2 = e_2^3 = e_3^2 e_4 = e_4^3 = 1$, $e_1 e_2^2 = e_3 e_4^2 = 2$, and $e_i e_j e_k = 0$ otherwise. In this particular case the image of $h = \sum_{i=1}^4 h_i e_i$ under F^t is represented by the matrix

$$\left[\begin{array}{cc|cc} h_2 & h_1 + 2h_2 & & \\ h_1 + 2h_2 & 2h_1 + h_2 & & \\ \hline & & h_4 & h_3 + 2h_4 \\ & 0 & h_3 + 2h_4 & 2h_3 + h_4 \end{array} \right],$$

which has a positive determinant unless $h = 0$.

REFERENCES

- [A] ARNDT, F. Zur Theorie der binären kubischen Formen. *Crelle's Journal* 53, 309-321 (1857).
- [Ar1] ARONHOLD, S. Zur Theorie der homogenen Funktionen dritten Grades von drei Variabeln. *Crelle's Journal* 39, 140-159 (1850).
- [Ar2] ——— Theorie der homogenen Funktionen dritten Grades von drei Veränderlichen. *Crelle's Journal* 55, 93-191 (1858).
- [At] ATIYAH, M. Some examples of complex manifolds. *Bonner Math. Schriften* 5, 1-28 (1958).
- [A/H/S] ATIYAH, M., N. HITCHIN and I. SINGER. Self-duality in four-dimensional Riemannian geometry. *Proc. R. Soc. Lond. A.* 362, 425-461 (1978).