

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CUBIC FORMS AND COMPLEX 3-FOLDS
Autor: Okonek, Ch. / Van de Ven, A.
Bibliographie
DOI: <https://doi.org/10.5169/seals-61829>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Finally we like to show that the non-emptiness condition on the index cone of a projective 3-fold with $h^{0,2} = 0$ gives non-trivial restrictions for the possible cup-forms if $b_2 \geq 4$. Further investigations of this condition will appear elsewhere [Sch].

EXAMPLE 17. Let H be a free \mathbb{Z} -module of rank 4 with basis $(e_i)_{i=1,\dots,4}$. Consider a trilinear form $F \in S^3 H^\vee$ and its adjoint map $F^t: H \rightarrow S^2 H^\vee$. The image $F^t(h)$ of an element $h \in H$ is in terms of the chosen basis $(e_i)_{i=1,\dots,4}$ represented by the symmetric 4×4 -matrix $[[he_i e_j]]_{i,j=1,\dots,4}$. Suppose this matrix is a diagonal sum $[[he_i e_j]]_{i,j=1,2} \oplus [[he_k e_l]]_{k,l=3,4}$ such that the determinants of both 2×2 -matrices are negative for every $h \in H \setminus \{0\}$.

In this case $F^t(h)$ were of signature $(1, -1, 1, -1)$ for every $h \in H \setminus \{0\}$, and we would have $I_F = \mathcal{H}_F = \emptyset$.

All these conditions can be met, e.g. by setting $e_1^2 e_2 = e_2^3 = e_3^2 e_4 = e_4^3 = 1$, $e_1 e_2^2 = e_3 e_4^2 = 2$, and $e_i e_j e_k = 0$ otherwise. In this particular case the image of $h = \sum_{i=1}^4 h_i e_i$ under F^t is represented by the matrix

$$\left[\begin{array}{cc|cc} h_2 & h_1 + 2h_2 & & 0 \\ h_1 + 2h_2 & 2h_1 + h_2 & & \\ \hline & & h_4 & h_3 + 2h_4 \\ 0 & & h_3 + 2h_4 & 2h_3 + h_4 \end{array} \right],$$

which has a positive determinant unless $h = 0$.

REFERENCES

- [A] ARNDT, F. Zur Theorie der binären kubischen Formen. *Crelle's Journal* 53, 309-321 (1857).
- [Ar1] ARONHOLD, S. Zur Theorie der homogenen Funktionen dritten Grades von drei Variablen. *Crelle's Journal* 39, 140-159 (1850).
- [Ar2] —— Theorie der homogenen Funktionen dritten Grades von drei Veränderlichen. *Crelle's Journal* 55, 93-191 (1858).
- [At] ATIYAH, M. Some examples of complex manifolds. *Bonner Math. Schriften* 5, 1-28 (1958).
- [A/H/S] ATIYAH, M., N. HITCHIN and I. SINGER. Self-duality in four-dimensional Riemannian geometry. *Proc. R. Soc. Lond. A.* 362, 425-461 (1978).

- [B] BEAUVILLE, A. Variétés Kählériennes dont la première classe de Chern est nulle. *J. Diff. Geom.* 18, 755-782 (1983).
- [Bo] BOREL, A. *Introduction aux groupes arithmétiques*. Publ. de L'Institut Math. de L'Univ. de Strasbourg. Hermann, Paris 1969.
- [B/H-C] BOREL, A., HARISH-CHANDRA. Arithmetic subgroups of algebraic groups. *Annals of Math.* (2) 75, 485-535 (1962).
- [B/H] BOREL, A. and F. HIRZEBRUCH. Characteristic classes and homogeneous spaces, III. *Amer. J. Math.* 82, 491-504 (1960).
- [B/V] BRIESKORN, E. and A. VAN DE VEN. Some complex structures on products of homotopy spheres. *Topology* 7, 389-393 (1968).
- [C/E] CALABI, E. and B. ECKMANN. A class of compact, complex manifolds which are not algebraic. *Ann. of Math.* 58, 494-500 (1959).
- [C] CAMPANA, F. On Twistor Spaces of the class C. *J. Diff. Geom.* 33, 541-549 (1991).
- [C/P] CAMPANA, F. and T. PETERNELL. Rigidity theorems for primitive Fano 3-folds. Preprint, Bayreuth 1991.
- [Ca] CASSELS, J. *An Introduction to the Geometry of Numbers*. Grundlehren Bd. 99, Springer, Berlin 1959.
- [Cay] CAYLEY, A. Tables of the binary cubic forms for the negative discriminants, $\equiv 0 \pmod{4}$, from -4 to -400; and $\equiv 1 \pmod{4}$, from -3 to -99; and for five irregular negative determinants. *The Quart. J.* 11, 246-261 (1871).
- [Co] CORNALBA, M. Una osservazione sulla topologia dei rivestimenti ciclici di varietà algebriche. *Bulletino U.M.I.* (5) 18-A, 323-328 (1981).
- [D/G/M] DELIGNE, P., P. GRIFFITHS and J. MORGAN. Real homotopy theory of Kähler manifolds. *Invent. math.* 29, 245-274 (1975).
- [D] DICKSON, L. *History of the theory of numbers. Vol. III. Quadratic and higher forms*. Reprinted by: Chelsea Publishing Company, New York, N.Y. 1971.
- [D/F] DONALDSON, S. and R. FRIEDMANN. Connected sums of self-dual manifolds and deformations of singular spaces. *Nonlinearity* 2, 197-239 (1989).
- [E/L1] EIN, L. and R. LAZARSFELD. Syzygies and Koszul cohomology of smooth projective varieties of arbitrary dimension. *Invent. math.* 111, 373-392 (1993).
- [E/L2] EIN, L. and R. LAZARSFELD. Global generation of pluricanonical and adjoint linear series on smooth projective threefolds. Preprint, 1992.
- [F] FRIEDMANN, R. On threefolds with trivial canonical bundle. *Proc. Symp. Pure Math.* 53, 103-134 (1991).
- [F/M] FRIEDMANN, R. and J. MORGAN. *Smooth four-manifolds and complex surfaces*. Ergeb. Math. 3. Folge, Springer, Heidelberg 1994.
- [G/N] GORDAN, P. and M. NOETHER. Über die algebraischen Formen, deren Hesse'sche Determinante identisch verschwindet. *Math. Ann.* 10, 547-568 (1876).
- [G/H] GRIFFITHS, P. and J. HARRIS. *Principles of Algebraic Geometry*. J. Wiley and Sons, New York 1978.
- [G/S] GRUNEWALD, F. and D. SEGAL. Some general algorithms. I: Arithmetic groups. *Ann. of Math.* 112, 531-583 (1980).
- [H] HILBERT, D. Über die vollen Invariantensysteme. *Math. Ann.* 42, 313-373 (1893).

- [H1] HIRZEBRUCH, F. Komplexe Mannigfaltigkeiten. In: *Proc. Int. Congress of Math. 1958*, 119-136. Cambridge University Press, 1960.
- [H2] —— Some examples of threefolds with trivial canonical bundle. Notes by J. Werner. Preprint, MPI Bonn 1985.
- [Hi] HITCHIN, N. Kählerian Twistor Spaces. *Proc. London Math. Soc.* 43, 133-150 (1981).
- [J1] JORDAN, C. Sur l'équivalence des formes algébriques. *C. R. Acad. Sc. XC*, 1422-1423 (1880).
- [J2] —— Mémoire sur l'équivalence des formes. *J. Ec. Polytechn. XLVIII*, 111-150 (1880).
- [J] JUPP, P. Classification of certain 6-manifolds. *Proc. Camb. Phil. Soc.* 73, 293-300 (1973).
- [K1] KATO, M. Examples of simply connected compact complex 3-folds. *Tokyo J. Math.* 5, 341-364 (1982).
- [K2] —— On compact complex 3-folds with lines. *Japan J. Math.* 11, 1-58 (1985).
- [K/Y] KATO, M. and A. YAMADA. Examples of simply connected complex 3-folds II. *Tokyo J. Math.* 9, 1-28 (1986).
- [K/M/M] KAWAMATA, Y., K. MATSUDA and K. MATSUKI. Introduction to the minimal model problem. *Adv. Stud. Pure Math.* 10, 283-360 (1987).
- [Ko1] KOLLÁR, J. Flips, flops, minimal models, etc. *Survey in Diff. Geom.* I, 113-199 (1991).
- [Ko2] KOLLÁR, J. Effective base point freeness. Preprint, Utah 1992.
- [Kr] KRAFT, H. *Geometrische Methoden in der Invariantentheorie*. Aspekte der Math., Vieweg, Braunschweig 1984.
- [L] LAMOTKE, K. The topology of complex projective varieties after S. Lefschetz. *Topology* 20, 15-51 (1981).
- [L/S] LANTERI, A. and D. STRUPPA. Projective manifolds with the same homology as \mathbf{P}_k . *Monatsh. Math.* 101, 53-58 (1986).
- [L/W] LIBGOBER, A. and J. WOOD. Differentiable structures on complete intersections - I. *Topology* 21, 469-482 (1982).
- [M] MAEDA, H. Some complex structures on the product of spheres. *J. Fac. Sci. Univ. Tokyo* 21, 161-165 (1974).
- [Mi] MIYAKOYA, Y. The Chern classes and Kodaira dimension of a minimal variety. *Adv. Stud. Pure Math.* 10, 449-476 (1987).
- [Mo] MORI, S. Threefolds whose canonical bundle is not numerically effective. *Ann. Math.* 116, 133-176 (1982).
- [M/M1] MORI, S. and S. MUKAI. On Fano 3-folds with $b_2 \geq 2$. *Manus. math.* 36, 147-162 (1981).
- [M/M2] MORI, S. and S. MUKAI. Classification of Fano 3-folds with $b_2 \geq 2$. *Adv. Studies Pure Math.* I, 101-129 (1981).
- [M/F] MUMFORD, D. and J. FOGARTY. *Geometric invariant theory*. 2nd ed. Ergeb. Math. 34, Springer, Heidelberg 1982.
- [Mu] MURRE, J. *Classification of Fano threefolds according to Fano and Iskovskikh*. Springer Lecture Notes in Mathematics 947, 35-92.
- [N] NEWSTEAD, P. *Introduction to moduli problems and orbit spaces*. Tata Inst. Lecture Notes, Springer, Heidelberg 1978.
- [O1] OGISO, K. On polarized Calabi-Yau 3-folds. *J. Fac. Sci. Univ. Tokyo* 38, 395-429 (1991).

- [O2] —— On algebraic fiber space structures on a Calabi-Yau 3-fold. Preprint, Bonn 1992.
- [O3] —— A note on the graded ring of a polarized Calabi-Yau 3-fold. Preprint, Bonn 1993.
- [O4] —— Two remarks on Moishezon Calabi-Yau 3-folds. Preprint, Bonn 1993.
- [O/V] OKONEK, Ch. and A. VAN DE VEN. *Stable Bundles, Instantons and C^∞ -structures on Algebraic Surfaces*. Encyclopaedia of Math. Sciences, Vol. 69, 198-249, Springer, Berlin-Heidelberg 1990.
- [P] POINCARÉ, H. Formes cubiques ternaires et quaternaires. *J. Ec. Polytechn.* 51, 45- 91 (1882).
- [S] SESHADRI, C. Geometric Reductivity over Arbitrary Base. *Advances in Math.* 26, 225-274 (1977).
- [Sch] SCHMITT, A. Zur Topologie dreidimensionaler komplexer Mannigfaltigkeiten. Ph. D. thesis, Zürich 1995.
- [Si] SIEGEL, C. The integer solutions of the equation $y^2 = ax^n + bx^{n-1} + k$. *J. London Math. Soc.* 1, 66-68 (1926).
- [Sm] SMALE, S. On the structure of manifolds. *Amer. J. Math.* 84, 387-399 (1962).
- [St] STURMFELS, B. *Algorithms in Invariant Theory*. Springer, New York 1993.
- [W] WALL, C. T. C. Classification problems in differential topology, V. On certain 6-Manifolds. *Invent. math.* 1, 355-374 (1966).
- [We] WERNER, J. Neue Beispiele dreidimensionaler Varietäten mit $c_1 = 0$. *Math. Gottingensis* 20, 1988.
- [Wi] WILSON, P. Calabi-Yau manifolds with large Picard number. *Invent. math.* 98, 139-155 (1989).
- [Y] YAU, S. On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère équation, I. *Comm. Pure & Appl. Math.* 31, 339-411 (1978).
- [Z1] ŽUBR, A. Classification of simply-connected six-dimensional Spin-manifolds. *Izv. Akad. Nauk SSSR, Ser. Mat.*, 39, 793-812 (1975).
- [Z2] —— Classification of simply connected six-dimensional manifolds. *Dokl. Akad. Nauk SSSR*, 255, 828-831 (1980).
- [Z3] —— Classification of simply-connected topological 6-manifolds. (*Rohlin-volume*) *LNM* 1346, 325-339 (1988).

(Reçu le 20 janvier 1995)

Ch. Okonek

Institut für Mathematik
 Universität Zürich
 Winterthurer Strasse 190
 8001 Zürich
 Switzerland

A. Van de Ven

Department of Mathematics
 Leiden University
 Niels Bohrweg 1
 Postbus 9512
 2300 RA Leiden

vide-leer-empty