

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 41 (1995)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CONCERNING A REAL-VALUED CONTINUOUS FUNCTION ON THE INTERVAL WITH GRAPH OF HAUSDORFF DIMENSION 2
Autor: Wingren, Peter
Kapitel: 0. Introduction
DOI: <https://doi.org/10.5169/seals-61818>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

CONCERNING A REAL-VALUED CONTINUOUS FUNCTION ON THE INTERVAL WITH GRAPH OF HAUSDORFF DIMENSION 2

by Peter WINGREN

ABSTRACT. A real-valued continuous nowhere-differentiable function on $[0, 1]$ is constructed. Its graph F is proved to have the following property. If B is a Borel subset of F and if the projection of B on $[0, 1]$ has positive Lebesgue measure, then the Hausdorff dimension of B is two.

0. INTRODUCTION

In 1903 Takagi [TAK, p. 176] gave an extremely simple construction of a nowhere differentiable real-valued continuous function on $[0, 1]$. Takagi's construction is

$$(1) \quad T(x) = \sum_{p=0}^{\infty} 2^{-p} \operatorname{dist}(2^p x, \mathbf{Z})$$

where each term is a scaled version of the sawtooth function

$$(2) \quad \operatorname{dist}(x, \mathbf{Z}) := \inf \{ |x - y| : y \in \mathbf{Z} \}.$$

Later, in 1930, van der Waerden [WAE] gave a similar example, which de Rham [RHA], in 1957, improved to an example identical with Takagi's.

It follows from a proof of Mauldin and Williams [M-W, pp. 795-797] that the graph of the Takagi function has a σ -finite linear Hausdorff measure and hence is of Hausdorff dimension 1.

In 1937 Besicovitch and Ursell [B-U, p. 29] constructed for an arbitrary α , $1 < \alpha < 2$, a real-valued nowhere-differentiable function in $C[0, 1]$ with graph of Hausdorff dimension α . They too used the sawtooth function $\operatorname{dist}(x, \mathbf{Z})$ as a building block in their construction.

In this paper we construct a real valued continuous function $f(x)$, $x \in [0, 1]$, whose graph has an optimal property with respect to Hausdorff dimension and measure.

We prove that for an arbitrary α , $1 < \alpha < 2$, $f(x)$ has the property

$\mathcal{P}(\alpha)$: Every Borel subset $B \subset \text{graph}(f)$, with projection on the x -axis of positive Lebesgue measure $m(\text{Proj}(B)) > 0$, has infinite α -dimensional Hausdorff measure

$$(3) \quad H^\alpha(B) = +\infty.$$

It is easy to see that

$$\mathcal{P}(\alpha) \forall \alpha < 2 \Leftrightarrow \mathcal{P}$$

where

\mathcal{P} : Every Borel set $B \subset \text{graph}(f)$ with $m(\text{Proj}(B)) > 0$ has Hausdorff dimension equal to two.

Rather than establish a general theorem valid for a class of functions we shall construct a single function with the desired property. The rationale is to provide a simple construction accompanied by a short, clear and instructive proof.

Our function is

$$(4) \quad f(x) = \sum_{p=0}^{\infty} 2^{-p} \text{dist}(2^{2^p} x, \mathbf{Z}).$$

Even though \mathcal{P} is established for only a single function f , the proof contains general methods extracted as Lemma 1 and Lemma 2. It appears that Lemma 1 is well known in more general cases than ours; compare [P-U, p. 159, the beginning of the proof of their Lemma 1]. However the proof is included here for completeness and because in the present case it is particularly simple.

The author is grateful to Professor V.P. Havin [HAV] for suggesting the investigation of fractal graphs with respect to $\mathcal{P}(\alpha)$, $\alpha = 1$.

PROBLEM. We believe that the following problem is unsolved.

Part 1: Construct a real valued function in $C[0, 1]$ with graph of Hausdorff dimension 1 and with property $\mathcal{P}(\alpha)$ for $\alpha = 1$.

Part 2: Determine the optimal smoothness in terms of the second difference of such a function.

Notation. The diameter of U is denoted by $|U|$ and the L^1 -norm of $g \in L^1(\mathbf{R})$ by $\|g\|$. If f is a real valued function in $C[0, 1]$, we write $\tilde{f}(x)$ for $(x, f(x))$. The notation $H^\alpha(F)$ stands for α -dimensional Hausdorff measure of a set $F \subset \mathbf{R}^2$ and $M^\alpha(F)$ is the α -dimensional net measure of F

constructed by closed dyadic cubes. The graph of a real valued function $f \in C[0, 1]$ is denoted by $\text{graph}(f)$. By a dyadic cube we mean a cube which is the Cartesian product of dyadic intervals. If Q is an arbitrary dyadic closed cube, then the band of type $\{(x, y) : (x, z) \in Q \text{ for some } z \in \mathbf{R}\}$ is called a dyadic band. In our construction the dyadic bands of width 2^{-2^p} play a special role. They are called bands of generation $p, p = 0, 1, 2, \dots$.

Acknowledgement. We would like to thank the referee for helpful suggestions.

1. A LEMMA ABOUT MASS DISTRIBUTION

By a mass distribution on a subset A of \mathbf{R}^2 we mean a measure μ on A such that $0 < \mu(A) < \infty$.

LEMMA 1. *Let f be a real valued measurable function defined on $[0, 1]$. Then there is a mass distribution μ on $F := \text{graph}(f)$ such that*

1) *for any two subintervals I and I' of $[0, 1]$, with $m(I) = m(I')$,*

$$\mu(I \times \mathbf{R}) = \mu(I' \times \mathbf{R})$$

and

2) *if for two Borel sets B_1 and B_2 in $[0, 1] \times \mathbf{R}$ there exists $(x_0, y_0) \in \mathbf{R}^2$ such that*

$$B_1 \cap F + (x_0, y_0) = B_2 \cap F$$

then

$$\mu(B_1) = \mu(B_2).$$

Proof. Let B be an arbitrary Borel set in \mathbf{R}^2 . Define

$$(5) \quad \mu(B) = m(\tilde{f}^{-1}(B)).$$

Then it is obvious that μ is a mass distribution on $\text{graph}(f)$ and 1) and 2) follow from the translation invariance of the Lebesgue measure.

2. A LEMMA ABOUT MASS DISTRIBUTION AND SUCCESSIVE TRANSLATIONS

LEMMA 2. *Let $g(y) \geq 0$ and $g(y) \in L^1(\mathbf{R})$. If I is a finite interval and d is a positive real number then*

$$(6) \quad \int_I \sum_{n=-\infty}^{\infty} g(y - nd) dy < \left(1 + \text{int} \frac{m(I)}{d}\right) \cdot \|g\|.$$