

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 41 (1995)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** CHAOTIC GROUP ACTIONS  
**Autor:** Cairns, G. / Davis, G. / Elton, D. / Kolganova, A. / Perversi, P.  
**Kapitel:** 3. Constructions of Chaotic Group Actions  
**DOI:** <https://doi.org/10.5169/seals-61820>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 19.08.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

suppose that groups  $G_i, i \in I$  act faithfully with all orbits finite on the Hausdorff spaces  $M_i, i \in I$  respectively. Now to each space  $M_i$ , add an additional isolated element  $x_i$  and denote  $\tilde{M}_i$  the union  $M_i \cup \{x_i\}$ . Then we define an action of  $G_i$  on  $\tilde{M}_i$  by using the action of  $G_i$  on  $M_i$  and making  $x_i$  a fixed point. Clearly  $G_i$  acts faithfully with all orbits finite on  $\tilde{M}_i$ . Now let  $M$  denote the subset of the infinite product  $\prod_{i \in I} \tilde{M}_i$  composed of all elements  $(y_i)_{i \in I}$  for which only finitely many of the  $y_i$  are different from  $x_i$ . We equip  $M$  with the topology induced by the product topology. Then clearly the infinite direct product  $\prod_{i \in I} G_i$  acts faithfully with all orbits finite on  $M$ .

Finally Part (f) is similar to Part (e); suppose that a group  $G$  acts chaotically on a space  $M$  and that  $H$  is a finite group. Then there is a natural action of the wreath product  $G \text{Wr} H$  on the space  $M \times H$ , where  $H$  is given the discrete topology (see [H]). It is easy to see that this action is chaotic.  $\square$

The following groups are known to be residually finite: Fuchsian groups [LS], the mapping class groups of compact Riemann surfaces [G], arithmetic groups [Se] and the group of  $p$ -adic integers [We]. It would be interesting to find natural chaotic actions of these groups.

### 3. CONSTRUCTIONS OF CHAOTIC GROUP ACTIONS

First recall that there are many examples of chaotic  $\mathbf{Z}$ -actions; that is, chaotic homeomorphisms. Perhaps the most basic example is that of the Anosov diffeomorphisms of tori and infranilmanifolds (see [Sm], [Mann]); these maps are chaotic since their periodic points are dense [BR] and by Anosov's closing lemma (see for instance [Sh]), they are transitive on their nonwandering set. (The Anosov diffeomorphisms of tori are just the linear hyperbolic maps; that is, linear maps with no eigenvalues on the unit circle.) Similarly, the pseudo-Anosov maps of compact surfaces are also chaotic (see Exposé 9 in [FLP] and the diagrams in [Mañ], pages 111-116).

Let us now give some general results.

**THEOREM 2.** *Consider a Hausdorff space  $M$  and the group  $\text{Hom}(M)$  of homeomorphisms of  $M$ . Then one has:*

(a) *If there are group inclusions*

$$G \leq H \leq K \leq \text{Hom}(M)$$

*then the action of  $H$  on  $M$  is chaotic if the actions of  $G$  and  $K$  on  $M$  are chaotic.*

- (b) If  $G \leq H \leq \text{Hom}(M)$  and  $G$  has finite index in  $H$  and if the action of  $G$  on  $M$  is chaotic, then the action of  $H$  on  $M$  is chaotic.
- (c) If  $M$  is locally compact and if  $\text{Hom}(M)$  is given the compact-open topology, then the action of  $G$  on  $M$  is chaotic if and only if the action on  $M$  of the closure  $\bar{G}$  of  $G$  in  $\text{Hom}(M)$  is chaotic.

*Proof.* In Part (a), notice that if a point  $x \in M$  has finite orbit under  $K$ , then  $x$  obviously has finite orbit under  $H$ . So if the action of  $K$  has finite orbits dense, then the action of  $H$  has finite orbits dense. On the other hand, if the action of  $G$  is topologically transitive, then clearly the action of  $H$  is also topologically transitive. So Part (a) holds. Part (b) is similar to Part (a).

In Part (c), again if the action of  $\bar{G}$  has finite orbits dense, then the action of  $G$  has finite orbits dense. Now suppose that the action of  $\bar{G}$  is topologically transitive. Let  $U$  and  $V$  be two non-empty open subsets of  $M$ . Then there exists  $g \in \bar{G}$  such that  $g(U) \cap V$  is non-empty. Let  $x$  be an element of  $U \cap g^{-1}(V)$  and let  $\Theta$  be the open subset of  $\bar{G}$  composed of elements that send  $x$  into  $V$ . Then  $g \in \Theta$  and since  $G$  is dense in  $\bar{G}$ , there exists  $h \in G \cap \Theta$ . So  $h(U) \cap V$  is non-empty and hence the action of  $G$  is topologically transitive.

Conversely, if  $M$  is locally compact, then the natural map  $\text{Hom}(M) \times M \rightarrow M$  is continuous. So, if a point  $x \in M$  has finite orbit under  $G$ , then since  $G$  is dense in  $\bar{G}$ , one has that  $G(x)$  is dense in  $\bar{G}(x)$ . Hence  $\bar{G}(x)$  is finite. So if the action of  $G$  has finite orbits dense, then the action of  $\bar{G}$  has finite orbits dense. Finally, if the action of  $G$  is topologically transitive, then obviously so too is the action of  $\bar{G}$ .  $\square$

#### 4. MANIFOLDS THAT ADMIT CHAOTIC GROUP ACTIONS

Chaotic homeomorphisms of the 2-dimensional disc can be constructed as follows. Starting with any Anosov diffeomorphism of the torus  $\mathbf{T}^2$ , one can quotient by the map  $\sigma: x \mapsto -x$ , to obtain a chaotic homeomorphism on the sphere  $\mathbf{S}^2$ . (This map was used in [Wa], p. 140 to show that expansiveness is not preserved under semi-conjugation.) Then, by blowing up the origin to a circle, one obtains a chaotic homeomorphism on the closed disc. Unfortunately this latter homeomorphism is not the identity on the boundary. This can be rectified by making a slight modification of the above construction. Instead of starting with an Anosov diffeomorphism of  $\mathbf{T}^2$ , one starts with linked twist map [D1] of the torus  $\mathbf{T}^2$ . A linked twist map is an