

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CHARACTERISTIC CLASSES, ELLIPTIC OPERATORS AND COMPACT GROUP ACTIONS
Autor: HEITSCH, James L.
Kapitel: 0. Introduction
DOI: <https://doi.org/10.5169/seals-87879>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

CHARACTERISTIC CLASSES, ELLIPTIC OPERATORS AND COMPACT GROUP ACTIONS

by James L. HEITSCH¹⁾

0. INTRODUCTION

Our aim in this paper is to introduce characteristic classes of vector bundles, to relate them to the indices of elliptic operators and the Lefschetz theorems for such operators, and finally to show how to use these to prove the non-existence of non-trivial actions preserving a Spin structure for compact connected Lie groups. Specifically, we will show how to prove the following.

THEOREM 5.2 ([HL 2]). *Let F be an oriented foliation of a compact oriented manifold M and assume that F admits a Spin structure. If a compact connected Lie group acts non-trivially on M as a group of isometries taking each leaf of F to itself and preserving the Spin structure on F , then the \hat{A} genus of F is zero.*

In [HL 1] and [HL 2], we assumed that F admitted a transverse invariant measure. In this paper, we show how to remove that rather restrictive hypothesis by employing the Haefliger forms of F . In particular, the traces we use here have values in those forms rather than in the complex numbers. A transverse invariant measure defines a map from the Haefliger zero forms to the reals, and applying it to the traces we use in this paper produces the traces used in [HL 1] and [HL 2]. Note in particular that all the results of [HL 1] and [HL 2] are still valid even if F does not admit a transverse invariant measure. One need merely ignore the transverse invariant measure and interpret the traces used as taking values in the Haefliger zero forms instead of the complex

¹⁾ Partially supported by NSF Grant DMS 9400676.

numbers. For another application of Haefliger forms and their cohomology, see [He].

An immediate corollary is the following theorem of Atiyah and Hirzebruch.

THEOREM 5.3 ([AH]). *Let M be a compact connected, oriented manifold which admits a Spin structure. If a compact connected Lie group acts non-trivially on M , then the \widehat{A} genus of M is zero.*

Theorem 5.2 is an application of the Lefschetz fixed point theorem for complexes elliptic along the leaves of a foliated manifold. We explain the classical Lefschetz theorem for elliptic complexes and give an outline of how to prove it. The original proofs of this theorem relied on the fact that the underlying manifold was compact. We outline a proof which does not rely on that fact, and so can be generalized to complexes defined along the leaves of a compact foliated manifold. Note that such leaves are in general not compact, but the fact that they come from a foliation of a compact manifold means that they have uniformly bounded geometry. It is this property which allows us to prove the foliation version of the Lefschetz theorem. We then show how the Lefschetz theorem leads to Theorem 5.2. Finally, we give a brief explanation of a very general rigidity theorem conjectured by Witten and proven by Bott and Taubes.

This paper is based on lectures given at the conference Actions Différentiables de Groupes Compacts, Espaces d'Orbites et Classes Caractéristiques, held at the Université des Sciences et Techniques du Languedoc in Montpellier in January, 1994. The author wishes to thank the organizers, especially Daniel Lehmann and Pierre Molino, for extending the invitation to him to speak at the conference and for making his stay in Montpellier so pleasant.

1. CHARACTERISTIC CLASSES AND MULTIPLICATIVE SEQUENCES

All objects considered in this paper will be smooth. Let E be an n dimensional complex vector bundle over the real manifold M . Denote the space of smooth sections of E by $C^\infty(E)$. A connection on E is a linear map $\nabla : C^\infty(E) \rightarrow C^\infty(T^*M \otimes E)$ satisfying

$$\nabla(f \cdot \sigma) = df \otimes \sigma + f \cdot \nabla\sigma$$

for any $\sigma \in C^\infty(E)$ and $f \in C^\infty(M)$, the smooth functions on M . T^*M denotes the cotangent bundle of M .