

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 42 (1996)  
**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** CHARACTERISTIC CLASSES, ELLIPTIC OPERATORS AND  
COMPACT GROUP ACTIONS  
**Autor:** HEITSCH, James L.  
**Kapitel:** 1. CHARACTERISTIC CLASSES AND MULTIPLICATIVE  
SEQUENCES  
**DOI:** <https://doi.org/10.5169/seals-87879>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. [Siehe Rechtliche Hinweise.](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. [Voir Informations légales.](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. [See Legal notice.](#)

**Download PDF:** 18.05.2025

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

numbers. For another application of Haefliger forms and their cohomology, see [He].

An immediate corollary is the following theorem of Atiyah and Hirzebruch.

**THEOREM 5.3 ([AH]).** *Let  $M$  be a compact connected, oriented manifold which admits a  $Spin$  structure. If a compact connected Lie group acts non-trivially on  $M$ , then the  $\hat{A}$  genus of  $M$  is zero.*

Theorem 5.2 is an application of the Lefschetz fixed point theorem for complexes elliptic along the leaves of a foliated manifold. We explain the classical Lefschetz theorem for elliptic complexes and give an outline of how to prove it. The original proofs of this theorem relied on the fact that the underlying manifold was compact. We outline a proof which does not rely on that fact, and so can be generalized to complexes defined along the leaves of a compact foliated manifold. Note that such leaves are in general not compact, but the fact that they come from a foliation of a compact manifold means that they have uniformly bounded geometry. It is this property which allows us to prove the foliation version of the Lefschetz theorem. We then show how the Lefschetz theorem leads to Theorem 5.2. Finally, we give a brief explanation of a very general rigidity theorem conjectured by Witten and proven by Bott and Taubes.

This paper is based on lectures given at the conference Actions Différentiables de Groupes Compacts, Espaces d'Orbites et Classes Caractéristiques, held at the Université des Sciences et Techniques du Languedoc in Montpellier in January, 1994. The author wishes to thank the organizers, especially Daniel Lehmann and Pierre Molino, for extending the invitation to him to speak at the conference and for making his stay in Montpellier so pleasant.

## 1. CHARACTERISTIC CLASSES AND MULTIPLICATIVE SEQUENCES

All objects considered in this paper will be smooth. Let  $E$  be an  $n$  dimensional complex vector bundle over the real manifold  $M$ . Denote the space of smooth sections of  $E$  by  $C^\infty(E)$ . A connection on  $E$  is a linear map  $\nabla : C^\infty(E) \rightarrow C^\infty(T^*M \otimes E)$  satisfying

$$\nabla(f \cdot \sigma) = df \otimes \sigma + f \cdot \nabla \sigma$$

for any  $\sigma \in C^\infty(E)$  and  $f \in C^\infty(M)$ , the smooth functions on  $M$ .  $T^*M$  denotes the cotangent bundle of  $M$ .

If  $\sigma_1, \dots, \sigma_n$  is a local basis of  $C^\infty(E)$  on an open set  $U$ , the local connection form  $\theta_U$  (which is an  $n \times n$  matrix of 1 forms) is defined by

$$\nabla \sigma_i = \sum_{j=1}^n \theta_{U,j}^i \otimes \sigma_j.$$

The local curvature form  $\Omega_U$  is the  $n \times n$  matrix of 2 forms

$$\Omega_U = d\theta_U - \theta_U \wedge \theta_U.$$

It is not difficult to show that if  $\tau_1, \dots, \tau_n$  is another local basis on the open set  $V$  with  $\tau_i = \sum_{j=1}^n g_{ij} \sigma_j$  on  $U \cap V$ , with  $g_{ij} \in C^\infty(U \cap V)$ , then on  $U \cap V$

$$(1.1) \quad \Omega_U = g \Omega_V g^{-1}$$

where  $g = [g_{ij}]$ .

Now consider the local differential form on  $U$ ,  $\det(I - \frac{1}{2\pi i} \Omega_U)$ . Because of (1.1), this is actually a well defined *global* form on  $M$ . This form depends only on  $\nabla$  and it is closed, so it defines a cohomology class  $c(E)$ , the total Chern class of  $E$ , which actually takes values in the real de Rham cohomology of  $M$ . This class depends only on  $E$  and we may write

$$c(E) = 1 + c_1(E) + \dots + c_n(E)$$

where  $c_k(E) \in H^{2k}(M, \mathbf{R})$  is the  $k$ th Chern class of  $E$ .

If  $E$  is an  $n$  dimensional real vector bundle over  $M$ , it is easy to show that  $c_{2k+1}(E \otimes_{\mathbf{R}} \mathbf{C}) = 0$ , and the  $k$ th Pontrjagin class of  $E$  is defined to be

$$p_k(E) = (-1)^k c_{2k}(E \otimes_{\mathbf{R}} \mathbf{C}).$$

For more on this see [KN] and [M].

Let  $Q(z) = \sum_{i=0}^{\infty} b_i z^i$  be a formal power series in  $z$ . Associated to  $Q$  is the *multiplicative sequence*  $K = (K_0, K_1, K_2, \dots)$  where each  $K_j$  is a polynomial in  $j$  indeterminants,  $K_j(\sigma_1, \dots, \sigma_j)$  given as follows. Denote by  $Q_j$  the degree  $j$  part of  $Q(z_1) \dots Q(z_j)$ , where each  $z_i$  has degree 1.  $Q_j$  is a symmetric polynomial in the  $z_i$  so it can be written as a polynomial in the elementary symmetric polynomials  $\sigma_1, \dots, \sigma_j$  in  $z_1, \dots, z_j$ , i.e.

$$Q_j = K_j(\sigma_1, \dots, \sigma_j).$$

For example, if  $Q(z) = 1 + z$ , then  $Q_j = z_1 \dots z_j = \sigma_j$  and  $K_j(\sigma_1, \dots, \sigma_j) = \sigma_j$ .

If  $Q(z)$  is an even power series,  $Q(z) = \sum_{i=0}^{\infty} b_{2i} z^{2i}$ , then the degree  $2j$  part of  $Q(z_1) \dots Q(z_j)$  can be written as a polynomial in the elementary symmetric polynomials  $\gamma_1, \dots, \gamma_j$  in  $z_1^2, \dots, z_j^2$ . We set  $K_j(\gamma_1, \dots, \gamma_j)$  to be this polynomial.

DEFINITION 1.2. (a) Let  $E$  be an  $n$  dimensional complex vector bundle over  $M$  and  $Q(z)$  a formal power series with associated multiplicative sequence  $K = (K_0, K_1, \dots)$ . The  $K$  genus of  $E$ ,  $K(E)$  is the de Rham cohomology class

$$K(E) = \sum_{j=0}^{\infty} K_j(c_1(E), \dots, c_j(E)).$$

(as  $K_j(c_1(E), \dots, c_j(E)) \in H^{2j}(M, \mathbf{R})$ , this is actually a finite sum).

(b) Let  $E$  be an  $n$  dimensional real vector bundle over  $M$  and  $Q(z)$  an even formal power series with associated multiplicative sequence  $K = (K_0, K_1, \dots)$ . Then the  $K$  genus of  $E$  is  $K(E) = \sum_{j=0}^{\infty} K_j(p_1(E), \dots, p_j(E))$ .

$K$  is called a multiplicative sequence because  $K(E_1 \oplus E_2) = K(E_1) \cdot K(E_2)$ .

#### IMPORTANT EXAMPLES

1.  $Q(z) = 1 + z$ . Then

$$K(z) = 1 + c_1(E) + c_2(E) + \dots = c(E).$$

2.  $Q(z) = z/\tanh(z)$ , which is even and gives the  $L$  genus of Hirzebruch.

Recall that the signature  $\text{Sign}(M)$  of a compact oriented  $4k$  dimensional manifold  $M$  is the signature of the quadratic form on  $H^{2k}(M, \mathbf{R})$  given by  $\alpha, \beta \mapsto \int_M \alpha \cdot \beta$ .

THEOREM 1.3 (Hirzebruch [H]).

$$\text{Sign}(M) = \int_M L_k(p_1(TM), \dots, p_k(TM)),$$

where  $TM$  is the tangent bundle of  $M$ .

Thus  $\text{Sign}(M)$  is completely determined by the Pontrjagin classes of  $M$ .

3.  $Q(z) = z/(1 - e^{-z})$ , gives the Todd genus  $Td$ .

Let  $M$  be a compact, complex  $n$  dimensional manifold and  $V$  a holomorphic vector bundle over  $M$ . Recall the Dolbeault complex of  $V$

$$0 \rightarrow A^{0,0}(V) \xrightarrow{\bar{\partial}_0} A^{0,1}(V) \xrightarrow{\bar{\partial}_1} \dots \xrightarrow{\bar{\partial}_{n-1}} A^{0,n}(V) \rightarrow 0$$

where  $A^{0,q}(V)$  is the space of differential forms on  $M$  of type  $0, q$  with coefficients in  $V$ .  $H^q(M, V) = \text{kernel } \bar{\partial}_q / \text{image } \bar{\partial}_{q-1}$  and it is finite dimensional (and isomorphic to  $H^q(M, \mathcal{O}(V))$  where  $\mathcal{O}(V)$  is the sheaf of germs of holomorphic sections of  $V$ ).

A fundamental invariant of  $V$  is its Euler class

$$\chi(M, V) = \sum_{q=0}^n (-1)^q \dim H^q(M, V).$$

The Riemann-Roch problem is to calculate this integer from topological information about  $M$  and  $V$ . The solution is given as follows. Suppose dimension  $V = k$ .  $e^{z_1} + \cdots + e^{z_k}$  is symmetric in the  $z_i$  so may be written as a power series in  $\sigma_1, \dots, \sigma_k$ , i.e.  $\sum_{i=1}^k e^{z_i} = k + \text{ch}_1(\sigma_1) + \text{ch}_2(\sigma_1, \sigma_2) + \cdots$  where

$$\text{ch}_j(\sigma_1(z_1, \dots, z_j), \dots, \sigma_j(z_1, \dots, z_j)) = \sum_{i=1}^k z_i^j / j!.$$

Set  $\text{ch}(V) = k + \text{ch}_1(c_1(V)) + \text{ch}_2(c_1(V), c_2(V)) + \cdots$ .

THEOREM 1.4 (The Riemann-Roch Theorem, [AS]).

$$\chi(M, V) = \int_M Td(TM) \cdot \text{ch}(V).$$

Thus  $\chi(M, V)$  is completely determined by the Chern classes of  $M$  and  $V$ .

4.  $Q(z) = (z/2)/\sinh(z/2) = z/(e^{z/2} - e^{-z/2})$  is an even function and gives the  $\hat{\mathcal{A}}$  genus.

Recall that  $\text{Spin}(n)$  is the simply connected double cover of  $SO(n)$ . A Spin structure on an oriented Riemannian manifold  $M$  of dimension  $n$  is a principal  $\text{Spin}(n)$  bundle  $P$  over  $M$  and an isomorphism of oriented bundles

$$P \times_{\text{Spin}(n)} \mathbf{R}^n \simeq TM.$$

$\text{Spin}(n)$  has a complex representation space  $\Delta$  of dimension  $2^n$ . See [ABS], [LM]. If  $n = 2k$ ,  $\Delta$  may be written as  $\Delta = \Delta^+ \oplus \Delta^-$  where  $\Delta^\pm$  are irreducible representations of dimension  $2^{n-1}$ . Set  $E^\pm = P \times_{\text{Spin}(n)} \Delta^\pm$ . The metric connection on  $M$  defines one on  $E = E^+ \oplus E^-$ , denoted  $\nabla$ . The Dirac operator  $D^+ : C^\infty(E^+) \rightarrow C^\infty(E^-)$  is defined as follows. Let  $c : C^\infty(T^*M \otimes E) \rightarrow C^\infty(E)$  be Clifford multiplication (we identify  $T^*M$  with  $TM$  using the metric on  $M$ ). Then  $D = c \cdot \nabla : C^\infty(E) \rightarrow C^\infty(E)$  and

$D$  maps  $C^\infty(E^+)$  to  $C^\infty(E^-)$  and vice-versa, since  $c$  does. Thus we may write

$$(1.5) \quad D = \begin{bmatrix} 0 & D^- \\ D^+ & 0 \end{bmatrix}$$

where  $D^\pm : C^\infty(E^\pm) \rightarrow C^\infty(E^\mp)$ . (See [AS] or [LM]). Now  $\ker D^+$  and  $\operatorname{coker} D^+ (\simeq \ker D^-)$  are finite dimensional and the Spinor index of  $M$ ,

$$\operatorname{Spin}(M) = \dim \ker D^+ - \dim \operatorname{coker} D^+.$$

THEOREM 1.6 ([AS]). *If  $M$  is a Spin manifold of  $\dim 2k$  then*

$$\operatorname{Spin}(M) = \int_M \hat{\mathcal{A}}(TM).$$

In particular, if  $2k \equiv 2(4)$ , then  $\operatorname{Spin}(M) = 0$  as  $\hat{\mathcal{A}}(TM)$  involves only the Pontrjagin classes of  $M$  and these occur only in dimensions  $\equiv 0(4)$ .

More generally, we may construct the twisted spinor complex. For this, let  $F$  be a complex bundle over  $M$  with hermitian metric and connection. Combining the connection on  $E$  with that on  $F$  we obtain a connection on  $E \otimes F$ . Composing this with Clifford multiplication

$$c : C^\infty(T^*M \otimes E \otimes F) \rightarrow C^\infty(E \otimes F)$$

we obtain the twisted Dirac operator  $D_F$  on  $E \otimes F$ . As before  $D_F$  interchanges  $E^+ \otimes F$  and  $E^- \otimes F$  and we get the *twisted Spin complex*

$$0 \rightarrow C^\infty(E^+ \otimes F) \xrightarrow{D_F^+} C^\infty(E^- \otimes F) \rightarrow 0.$$

The kernel and cokernel of  $D_F^+$  are finite dimensional and the twisted spinor index is

$$\operatorname{Spin}(M, F) = \dim \ker D_F^+ - \dim \operatorname{coker} D_F^+.$$

THEOREM 1.7 ([AS]).

$$\operatorname{Spin}(M, F) = \int_M \hat{\mathcal{A}}(TM) \cdot \operatorname{ch}(F).$$