

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: CHARACTERISTIC CLASSES, ELLIPTIC OPERATORS AND
COMPACT GROUP ACTIONS
Autor: HEITSCH, James L.
Kapitel: 6. The Rigidity Theorem of Witten
DOI: <https://doi.org/10.5169/seals-87879>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 30.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$w_\alpha^L(z)$ on $N_\alpha \cap L$ which represents the cohomology class $\widehat{A}(N_\alpha \cap L)B(N_\alpha \cap L, z)$. Then $w_\alpha^L(z)$ is the form a_α^L given in the foliation Lefschetz theorem for z acting on the leafwise Spin complex, and it defines a smooth form $w_\alpha(z)$ on N_α . Thus for $z \in S^1$, z not a root of unity, we have

$$L(z) = \int_N w(z) = \sum_\alpha \int_{N_\alpha} w_\alpha(z).$$

Now notice that the right side of this equation defines a function $A(F, z)$ on the complex plane with values in the Haefliger forms of F . Also note that $A(F, z)$ has poles only at roots of unity and no pole at $z = \infty$, since $w_\alpha(z)$ has poles only at roots of unity and no pole at $z = \infty$. Because of the factor of $(z^d)^{1/2}$, $A(F, 0) = 0$. For $z \in S^1$, z not a root of unity, $A(F, z) = L(z)$. But $L(z)$ is defined for all $z \in S^1$ and by Theorem 5.5 it is continuous on S^1 . Thus $A(F, z)$ has no poles at all. Since it is analytic and bounded, it is constant and hence is identically zero. Therefore $L(z) = 0$ for all $z \in S^1$, but $L(1) = \widehat{A}(F)$ so we are done.

The compactness of G is essential, as in [HL2], we give an example of an infinite discrete group acting by leaf preserving isometries on a compact oriented foliated manifold M, F and G preserves a Spin structure on F . The foliation F admits an invariant transverse measure which defines a map from the Haefliger zero forms of F to \mathbf{C} . The image of $\widehat{A}(F)$ under this map is non-zero, so $\widehat{A}(F) \neq 0$.

6. THE RIGIDITY THEOREM OF WITTEN

In 1986, Witten [W] predicted rigidity theorems for the indices of certain elliptic operators on manifolds with S^1 actions. The genesis for Witten's conjecture was his study of the Dirac operator on the free loop space $\mathcal{L}M$ (an infinite dimensional manifold) of a Spin manifold M . $\mathcal{L}M$ admits a natural S^1 action whose fixed point set is diffeomorphic to M . The sequences of bundles $R(q)$ and $R'(q)$ described below were derived from the normal bundle of M in $\mathcal{L}M$ and from the formal analogue on $\mathcal{L}M$ of the fixed point formula for the Dirac operator in the finite dimensional case.

Let $D : C^\infty(E_1) \rightarrow C^\infty(E_2)$ be an elliptic operator on a compact manifold M and suppose M admits an S^1 action preserving D . Then as noted above, $\text{Index}(D)$ is a virtual S^1 module and has a decomposition into a finite sum of irreducible complex one dimensional representations

$$\text{Index}(D) = \sum a_m L^m$$

where $z \in S^1$ acts on L^m by multiplication by z^m . D is called *rigid* if all the a_m for $m \neq 0$ are zero, i.e. if the representation L^m , $m \neq 0$ occurs in kernel D with multiplicity a then it occurs in cokernel D with the same multiplicity a .

Denote by $S^k(T)$ and $\lambda^k(T)$ the k th symmetric and exterior powers of $T = TM$ and set

$$S_a(T) = \sum_{k=0}^{\infty} a^k S^k(T)$$

$$\lambda_a(T) = \sum_{k=0}^{\infty} a^k \lambda^k(T).$$

Let R_n and R'_n be the sequences of bundles defined by the formal power series

$$R(q) = \sum_{n=0}^{\infty} q^n R_n = \bigotimes_{\ell=1}^{\infty} \lambda_{q^\ell}(T) \bigotimes_{m=1}^{\infty} S_{q^m}(T)$$

$$R'(q) = \sum_{n=0}^{\infty} q^{n/2} R'_n = \bigotimes_{\ell=\frac{1}{2}, \frac{3}{2}, \dots}^{\infty} \lambda_{q^\ell}(T) \bigotimes_{m=1}^{\infty} S_{q^m}(T)$$

Now suppose M is a $2n$ dimensional compact Riemannian Spin manifold and denote by D^+ the Dirac operator of M . For each n we may form the operators

$$D^+ \otimes (E^+ \oplus E^-) \otimes R_n \quad \text{and} \quad D^+ \otimes R'_n.$$

THEOREM 6.1. *These operators are rigid under any S^1 action on M by isometries, i.e. the induced action on the index of any of these operators is the trivial action.*

This is the theorem conjectured by Witten and first proven by Taubes [T]. A beautiful proof of it appears in [BT].

Roughly speaking Bott and Taubes' proof goes as follows. First they show that the Signature operator $d_S = D^+ \otimes (E^+ \oplus E^-)$ is rigid by an argument similar to that presented above. Combining this result with the power series $R(q)$ and interpreting $\text{ch}(\text{Index}(d_S \otimes R(q)))$ as a meromorphic function on the complex torus $T_{q^2} = \mathbf{C}^*/q^2$, they show that it has poles only at roots of unity and no poles on a certain circle $S^1 \subset T_{q^2}$. The Spin hypothesis then implies

that it has no poles at all and hence is constant. Thus the character of S^1 given by its action on $\text{Index}(d_S \otimes R(q))$ is constant, and so the action must be trivial as claimed. They then give separate arguments to extend this result to $D^+ \otimes R'_n$.

These results all extend in a straight forward way to S^1 actions preserving a foliation (see [HL 2]).

REFERENCES

- [AB] ATIYAH, M.F. and R. BOTT. The Lefschetz fixed point formula for elliptic complexes I and II. *Annals of Math.* 86 (1967), 374–407; 88 (1968), 451–491.
- [ABP] ATIYAH, M.F., R. BOTT and V.K. PATODI. On the Heat Equation and the Index Theorem. *Invent. Math.* 19 (1973), 279–330; errata *ibid* 28 (1975), 277–280.
- [ABS] ATIYAH, M.F., R. BOTT and A. SHAPIRO. Clifford Modules. *Topology* 3, Suppl. 1 (1964), 3–38.
- [AH] ATIYAH, M.F. and F. HIRZEBRUCH. Spin Manifolds and Group Actions. *Essays on Topology and Related Topics* (Mémoires dédiés à Georges de Rham), Springer Verlag, New York (1970), 18–27.
- [AS] ATIYAH, M.F. and I.M. SINGER. The index of elliptic operators : III, *Annals of Math.* 87 (1968), 546–604.
- [Be] BENAMEUR, M-T. Un théorème de Lefschetz longitudinal en K-théorie équivariante. *C.R.A.S. (Série I)* 317 (1993), N° 5, 491–494.
- [B] BISMUT, J.M. The Atiyah-Singer index theorem for families of Dirac operators : two heat equation proofs. *Invent. Math.* 83 (1986), 91–151.
- [BT] BOTT, R. and C. TAUBES. On the Rigidity Theorems of Witten. *Journal of AMS* 2 (1989), 137–186.
- [C] CONNES, A. Sur la Théorie Non-Commutative de l'Intégration, Algèbres d'Opérateurs. Springer Verlag, Berlin. *Lecture Notes in Math.* 725 (1979), 19–143.
- [G] GETZLER, E. A short proof of the local Atiyah-Singer Index Theorem. *Topology* 25 (1986), 111–117.
- [Gi] GILKEY, P. Invariance Theory, the Heat Equation, and the Atiyah-Singer Index Theorem. Publish or Perish Press, Wilmington, Del. *Math. Lec. Series 11* (1984).
- [Ha] HAEFLIGER, A. Some remarks on foliations with minimal leaves. *J. Diff. Geo.* 15 (1980), 269–284.
- [He] HEITSCH, J.L. Bismut superconnections and the Chern character for Dirac operators on foliated manifolds. *K-Theory* 9 (1995), 507–528.
- [HL 1] HEITSCH, J.L. and C. LAZAROV. A Lefschetz Theorem for Foliated Manifolds. *Topology* 29 (1990), 127–162.
- [HL 2] HEITSCH, J.L. and C. LAZAROV. Rigidity theorems for foliations by surfaces and spin manifolds. *Mich. J. Math.* 38 (1991), 285–297.