Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

**Band:** 42 (1996)

**Heft:** 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SYSTEMS OF CURVES ON A CLOSED ORIENTABLE SURFACE

Autor: EDMONDS, Allan L.

Kapitel: 4. SUFFICIENCY FOR INDEPENDENT HOMOLOGY CLASSES

**DOI:** https://doi.org/10.5169/seals-87881

## Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

**Download PDF:** 05.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

then we can conclude that A is connected and we are done. If some R has more than two boundary components, then it contains two positive curves or two negative curves, and we can proceed as above to reduce the number of components of A.

It remains to consider the case where each component  $R_k$  of  $\widehat{F}$  has exactly two boundary components of the form  $A_i^+$  and  $A_j^-$ , where  $A_i$  and  $A_j$  are distinct components of A. In this case we conclude that we can arrange the components of A in a sequence  $A_1, A_2, \ldots, A_n$ , so that  $A_1$  is homologous to  $A_2, A_2$  is homologous to  $A_3, \ldots, A_n$  is homologous to  $A_1$ . In this case, then, the number n of components is exactly the divisibility of  $\alpha$ .

# 4. SUFFICIENCY FOR INDEPENDENT HOMOLOGY CLASSES

In this section we complete the proof of Theorem 2, dealing with the case of a set of homology classes consisting of independent elements.

LEMMA 4.1. Let F be a closed orientable surface and let  $\alpha_1, \ldots, \alpha_n \in H_1(F)$  be independent homology classes that span a summand of  $H_1(F)$  on which the intersection pairing of F vanishes. Then there exists  $\gamma \in H_1(F)$  such that  $\gamma \cdot \alpha_n = 1$  and  $\gamma \cdot \alpha_i = 0$  for i < n.

Proof. This is a consequence of Poincaré Duality.

PROPOSITION 4.2. Let F be a closed orientable surface and let  $\alpha_1, \ldots, \alpha_n \in H_1(F)$  be independent homology classes that span a summand of  $H_1(F)$  on which the intersection pairing of F vanishes. Then there exist pairwise disjoint simple closed curves  $A_1, \ldots, A_n$  in F representing the homology classes  $\alpha_1, \ldots, \alpha_n$ .

*Proof.* The proof will proceed by induction on n. The case n=1 is given by Proposition 3.3.

Now inductively consider the case of n>1 homology classes. By Proposition 3.3 we can find a simple closed curve  $A_n$  in F representing  $\alpha_n$ . We claim that there is a simple closed curve  $B_n$  in F representing a homology class  $\beta_n$  such that  $B_n$  meets  $A_n$  in exactly one point and such that  $[B_n] \cdot \alpha_i = 0$  for i < n. By Lemma 4.1 there is a homology class  $\gamma_n \in H_1(F)$  such that  $\alpha_i \cdot \gamma_n = \delta_{i,n}$ . We begin by representing  $\gamma_n$  by a simple closed curve B transverse to  $A_n$ . By tubing together neighboring pairs of intersection of B with  $A_n$  of opposite sign we can transform B into a disjoint union B'

of simple closed curves meeting  $A_n$  in exactly one point. Now we can band together the components of B', using bands in the complement of  $A_n$  to create a closed curve B'' representing  $\gamma_n$  and meeting  $A_n$  in exactly one point. But B'' may now have self-intersections. We may then eliminate the self-intersections by sliding segments of B'' over  $A_n$ . This creates a simple closed curve  $B_n$  meeting  $A_n$  in exactly one point, and representing a homology class of the form  $\beta_n = \gamma_n + k\alpha_n$ , which proves the claim.

Now the union of the two curves  $A_n$  and  $B_n$  has a small neighborhood N of the form of a once punctured torus. Let  $F_n$  denote the result of removing N and replacing it with a disk D. Then  $F_n - D = F - N \subset F$  and inclusion identifies  $H_1(F_n)$  with the orthogonal complement of  $\alpha_n$  and  $\beta_n$  in  $H_1(F)$ . Thus the homology classes  $\alpha_1, \ldots, \alpha_{n-1}$  determine well-defined classes in  $H_1(F_n)$ , which we continue to call by the same names. By induction there are pairwise disjoint simple closed curves  $A_1, \ldots, A_{n-1}$  in  $F_n$  representing the homology classes  $\alpha_1, \ldots, \alpha_{n-1}$ . Then these curves also live in F, determining the same homology classes, and are disjoint from the curve  $A_n$ . This completes the proof.

Here is a sketch of a standard but somewhat more learned proof of Proposition 4.2, suggested by M. Kervaire: Any basis for a self-annihilating summand of a skew-symmetric inner product space over **Z** can be extended to be part of a symplectic basis. Any two symplectic bases are related by an isometry of the inner product space. Half of a fixed standard symplectic basis is clearly represented by standard pairwise disjoint simple closed curves in a standard model of the surface. And any isometry is induced by a homeomorphism of the surface, so that the standard curves are taken to the desired curves. To see that any isometry is induced by a homeomorphism one can argue that the symplectic group is generated by certain elementary automorphisms and that these elementary automorphisms are induced by Dehn twist homeomorphisms around standard curves on the surface.

## 5. DISJOINT SIMPLE CLOSED CURVES ON A PLANAR SURFACE

Subsequent proofs of realizability of non-independent homology classes will proceed by cutting the surface along curves representing a basis for homology until it becomes a punctured 2-sphere and then representing the remaining homology classes by disjoint curves on this planar surface. We