

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: FACTOR EQUIVALENCE RESULTS FOR INTEGERS AND UNITS
Autor: De Smit, Bart
Kapitel: 3. Rings of integers
DOI: <https://doi.org/10.5169/seals-87884>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

(2.5) PROPOSITION. *If N and M are factor equivalent then for any $A[G]$ -linear embedding $j: M \rightarrow N$ the function $H \mapsto [N^H : j(M^H)]_A$ is factorizable.*

Proof. We have $j = \varphi i$, where i is an embedding as in (2.4) and φ is a $K[G]$ -linear automorphism of $N \otimes_A K$. Using [15, Ch. III, § 1, Prop. 2] and the notation of (2.3) we see that

$$[N^H : j(M^H)]_A = d_\varphi(H) \cdot [N^H : i(M^H)]_A .$$

This is a product of two factorizable functions by (2.3) and by our choice of i . \square

The fact that “factor equivalence” is an equivalence relation is an easy consequence of (2.5). If \mathfrak{p} is a prime of K not dividing $\#G$ then condition (1) of (2.4) implies that the \mathfrak{p} -part of $[N^H : i(M^H)]_A$ is factorizable. One can prove this with [16, § 15.2] and [16, § 14.4, Lemma 21].

(2.6) REMARK. The definitions of factorizability given by Fröhlich [8; 9] and Burns [2] for abelian groups G are in agreement with our definitions. They also define the notion called \mathbf{Q} -factorizability in the abelian case, which is a stronger condition than factorizability. However, the function that one wants to be factorizable in the definition of factor equivalence automatically satisfies this stronger condition if it is factorizable. Thus, \mathbf{Q} -factor equivalence is the same as factor equivalence.

In [4, § 3] a factorizable function f with values in $I(\mathbf{Q})$ must also satisfy an additional condition: there should be a map g from the group of complex characters $R_{\mathbf{C}}(G)$ to $I(E)$, where E is some normal number field containing all character values of G , such that g is $\text{Gal}(E/\mathbf{Q})$ -equivariant, and such that $g(1_H^G)$ is the E -ideal generated by $f(H)$. It is not hard to see that this condition is satisfied by all functions that are factorizable in our sense.

3. RINGS OF INTEGERS

Let A be a Dedekind domain with quotient field K of characteristic zero and let L a Galois extension of K with Galois group G . The integral closure B of A in L is again a Dedekind domain. Assume that for all primes of L the residue class field extension is separable.

(3.1) THEOREM. *The $A[G]$ -lattices B and $A[G]$ are factor equivalent.*

Proof. Define a $B[G]$ -module structure on $B \otimes_A B$ by letting B act on the left factor and G on the right. We will show first that $B \otimes_A B$ and $B[G]$ are factor equivalent as $B[G]$ -lattices. Define the canonical $B[G]$ -linear map $\varphi: B \otimes_A B \rightarrow B[G]$ by

$$x \otimes y \mapsto \sum_{\sigma \in G} x\sigma(y) \cdot \sigma^{-1} .$$

Let H be a subgroup of G . If $\sigma_1, \dots, \sigma_n$ are the K -embeddings of L^H in L , and if there is an A -basis $\omega_1, \dots, \omega_n$ of B^H , then the restriction $(B \otimes_A B)^H \rightarrow B[G]^H$ of φ is a B -linear map with matrix $(\sigma_i(\omega_j))_{ij}$ on the bases $\{1 \otimes \omega_j\}$ and $\{b_i\}$, where b_i is the formal sum of those $\sigma \in G$ for which σ^{-1} restricts to σ_i . The square of the determinant of this matrix generates the discriminant $\Delta(B^H/A)$ as an A -ideal. By localization it follows that even if B is not free over A , we have

$$[B[G]^H : \varphi(B \otimes_A B)^H]_B^2 = \Delta(B^H/A) \cdot B .$$

By Hasse's conductor discriminant product formula [15, Ch. VI, §3] the ideal $\Delta(B^H/A)$ is a factorizable function of H , so $B \otimes_A B$ and $B[G]$ are factor equivalent $B[G]$ -lattices.

In order to descend to $A[G]$ -lattices, note that there exists an $A[G]$ -linear injection $i: A[G] \rightarrow B$ by the normal basis theorem, and consider the induced $B[G]$ -linear map $i_*: B[G] \rightarrow B \otimes_A B$ that sends $b\sigma$ to $b \otimes i(\sigma)$ for $b \in B$ and $\sigma \in G$. We have

$$[(B \otimes_A B)^H : i_*(B[G])^H]_B = [B^H : i(A[G])^H]_A \cdot B ,$$

and by (2.5) we know that the left hand side is a factorizable function of H . But then the A -index $[B^H : i(A[G])^H]_A$ is also factorizable. \square

4. S -UNITS

Let L/K be a Galois extension of number fields with Galois group G , and let S be a finite G -stable set of primes of L containing the infinite primes. The ring of S -integers of L consists of all elements of L that are integral outside S . Its class number is written as $h_S(L)$ and its unit group, the group of S -units of L , is denoted by $U_S(L)$. The group of roots of unity in L is denoted by μ_L and its order is written as $w(L)$.