

**Zeitschrift:** L'Enseignement Mathématique  
**Herausgeber:** Commission Internationale de l'Enseignement Mathématique  
**Band:** 42 (1996)  
**Heft:** 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

**Artikel:** DOUBLE VALUED REFLECTION IN THE COMPLEX PLANE  
**Autor:** Webster, S. M.  
**Kapitel:** 5. INVOLUTIONS ON A TORUS  
**DOI:** <https://doi.org/10.5169/seals-87870>

### **Nutzungsbedingungen**

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

### **Terms of use**

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

**Download PDF:** 14.03.2026

**ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>**

must descend to the Green's function of  $E^0$ . For the annulus  $\tilde{G}$  may be constructed, for example, by the method of electrostatic images, using the reflections (4.3) in the boundary circles of  $A_1^\mu$  (see [2], [7]).

The lemniscate (4.16) may serve as a useful model for domains with corners.

## 5. INVOLUTIONS ON A TORUS

We return to the situation at the beginning of section 3, but with a non-simply connected Riemann surface  $\Gamma$ . Let  $\pi: \tilde{\Gamma} \rightarrow \Gamma$  be the universal covering space, and  $\Lambda \subseteq \text{Aut}(\tilde{\Gamma})$  be the group of covering transformations. We consider liftings

$$(5.1) \quad \tilde{\tau}_i, \tilde{\rho}: \tilde{\Gamma} \rightarrow \tilde{\Gamma}$$

of  $\tau_i, \rho$ . For each  $\gamma \in \Lambda$  there is a  $\gamma_1 \in \Lambda$  with

$$(5.2) \quad \tilde{\rho} \circ \gamma = \gamma_1 \circ \tilde{\rho},$$

and similarly for  $\tau_i$ . Also

$$(5.3) \quad \tilde{\tau}_i^2, \tilde{\rho}^2 \in \Lambda.$$

In this section we take  $\tilde{\Gamma} = \mathbf{C}$ , and  $\Lambda$  a group of translations, which we shall also identify with an additive subgroup of  $(\mathbf{C}, +)$  of rank one or two over  $\mathbf{Z}$ . We shall determine what restrictions on  $\Lambda$  are forced if  $\Gamma$  is the complexification of a real curve admitting double valued reflection. We are, of course, interested in the corresponding objects on  $\Gamma = \mathbf{C}/\Lambda$ .

We drop the tilde notation and let  $t \in \mathbf{C}$ . In view of (5.3), we consider

$$(5.4) \quad \tau_i(t) = \varepsilon_i t + c_i, \varepsilon_i^2 = 1, (\varepsilon_i + 1)c_i \in \Lambda, i = 1, 2;$$

$$(5.5) \quad \sigma(t) = \tau_1 \tau_2(t) = \varepsilon_1 \varepsilon_2 t + c_1 + \varepsilon_1 c_2;$$

$$(5.6) \quad \rho(t) = a\bar{t} + b, a\bar{a} = 1, b + a\bar{b} \in \Lambda.$$

In case  $\tau_2 = \rho \tau_1 \rho$ , we have

$$(5.7) \quad \varepsilon_1 = \varepsilon_2, c_2 = a(\varepsilon_1 \bar{b} + \bar{c}_1) + b.$$

The constants  $c_i, b$  are only determined mod  $\Lambda$ . For each  $\tau_i$ , either  $\varepsilon_i = -1$  and  $c_i \in \mathbf{C}$  can be arbitrary, or  $\varepsilon_i = +1$  and  $2c_i \in \Lambda$ .

We set

$$(5.8) \quad a = e^{2\alpha i}, 0 \leq \alpha < \pi, \rho_\alpha(t) = a\bar{t}, \\ l_\alpha = \{\lambda e^{i\alpha} \mid \lambda \in \mathbf{R}\} = \{t \mid \text{Re}(ie^{-i\alpha} t) = 0\}.$$

$\rho_\alpha$  is the reflection the line  $l_\alpha$ . If we apply the condition (3.3) we get

$$(5.9) \quad \rho_\alpha(\Lambda) = \Lambda .$$

Thus,  $\Lambda$  must be symmetric about  $l_\alpha$ . Clearly,  $b + a\bar{b} \in l_\alpha$ , so (5.6) gives

$$(5.10) \quad b + a\bar{b} = \omega_0 \in \Lambda \cap l_\alpha ,$$

and  $b$  lies on the line perpendicular to  $l_\alpha$  and passing through  $\frac{1}{2}\omega_0$ . This line has the equation

$$(5.11) \quad 2\operatorname{Re}(e^{-i\alpha}(t - \omega_0/2)) = e^{-i\alpha}(t + a\bar{t} - \omega_0) = 0 .$$

If  $\Lambda$  satisfies (5.9) for some angle  $\alpha$ , we choose  $\omega_0 \in \Lambda \cap l_\alpha$ , for example  $\omega_0 = 0$ . We then choose  $b$  satisfying (5.10), and construct  $\rho$ . If we replace  $b$  by  $b + \omega_*$ ,  $\omega_* \in \Lambda \cap l_\alpha$ , then  $\omega_0$  gets replaced by  $\omega_0 + 2\omega_*$ . Hence, there are at most two inequivalent choices for  $\omega_0$  on  $l_\alpha$ .

A point  $t_0 \in \mathbf{C}$  represents a fixed-point of  $\rho$  if and only if it lies on a line of the form

$$(5.12) \quad t - a\bar{t} - b = \omega'_0 \in \Lambda .$$

Since  $t - a\bar{t}$  is orthogonal to  $e^{i\alpha}$ ,  $\omega'_0$  must lie on the line perpendicular to  $l_\alpha$  and passing through  $-\frac{1}{2}\omega_0$ ,

$$(5.13) \quad t + a\bar{t} + \omega_0 = 0 .$$

If there is an  $\omega'_0 \in \Lambda$  on this line, then the fixed-point set  $FP(\rho)$  of  $\rho$  is non-empty, and is given by (5.12) for all such  $\omega'_0$ . (5.12) is the line parallel to  $l_\alpha$  and passing through  $\frac{1}{2}(b + \omega'_0)$ ; hence, there are at most two inequivalent choices of  $\omega'_0$ .

First consider the very simple case

$$(5.14) \quad \Lambda = \{2\pi ki \mid k \in \mathbf{Z}\} .$$

From (5.9) we can only have  $\alpha = 0$ , or  $\alpha = \pi/2$ . In the first case,  $l_\alpha$  is the real axis,  $a = 1$ , and  $\omega_0 = 0$ ,  $b = ib_2$  is purely imaginary. We may take  $\omega'_0 = 2ki$ ,  $k = 1, 2$ ; thus

$$(5.15) \quad \rho(t) = \bar{t} + ib_2, \quad FP(\rho) = \{Im t = b_2/2\} \cup \{Im t = b_2/2 + \pi\} .$$

In the second case  $l_\alpha$  is the imaginary axis,  $a = -1$ , and we may take either  $\omega_0 = 0$ , or  $\omega_0 = 2\pi i$ . Then, either  $b = b_1 \in \mathbf{R}$ , or  $b = b_1 + i\pi$ . In the first case we have  $\omega'_0 = 0$ , while in the second case there is no  $\omega'_0$ . Thus,

$$(5.16) \quad \rho(t) = -\bar{t} + b_1, \quad FP(\rho) = \{Re t = b_1/2\} .$$

Of course,  $\Gamma \equiv \mathbf{C}^*$ , and the covering projection  $\pi: \mathbf{C} \rightarrow \Gamma$  is just  $\zeta = \pi(t) = e^t$ . The first choice of  $\rho$  gives reflection in the two rays  $\arg \zeta = \frac{1}{2}b_2, \frac{1}{2}b_2 + \pi$ . The second gives reflection in the circle  $|\zeta| = e^{b_1}$ . We must still make a choice of  $\tau_1$  as in (5.4), and find a "minimal" function  $F$  which is  $\tau_1$ -invariant. Relative to  $\zeta$  we have  $\tau_1(\zeta) = \mu\zeta^{\varepsilon_1}$ ,  $\mu = e^{c_1}$ . For  $\varepsilon_1 = -1$ , we take  $F = f + f \circ \tau_1 = \zeta + \mu\zeta^{-1}$ . For  $\varepsilon_1 = +1$ ,  $c_1 = \pi i$ , we take  $F = f \cdot f \circ \tau_1 = -\zeta^2$ . We have already used these in the case of conics.

Next we consider a rank two lattice (4.8), and after a coordinate change if necessary, choose a normalized basis  $\omega_1 = 1, \omega_2 = \omega$ ,

$$(5.17) \quad \operatorname{Im} \omega > 0, \quad -\frac{1}{2} < \operatorname{Re} \omega \leq \frac{1}{2}, \quad |\omega| \geq 1,$$

$$|\omega| = 1 \Rightarrow \operatorname{Re} \omega \geq 0.$$

We consider those  $\Lambda$  which satisfy the reality condition (5.9) [3], [5]. Since  $a = \rho_\alpha(1) \in \Lambda$ , we have  $a = n_1 + n_2\omega$ , and

$$(5.18) \quad \begin{aligned} 1 = a\bar{a} &= n_1^2 + n_2^2|\omega|^2 + 2n_1n_2\operatorname{Re} \omega \\ &\geq n_1^2 + n_2^2 - 2|n_1n_2\operatorname{Re} \omega| \\ &\geq |n_1|^2 + |n_2|^2 - |n_1n_2| \geq |n_1n_2|. \end{aligned}$$

There are two cases. If  $n_1n_2 = 0$ , then either  $a = \pm 1$ , or  $|\omega| = 1$  and  $a = \pm \omega$ . Otherwise,  $|n_1| = |n_2| = 1$ , and we have the equalities in (5.18). Equality in all three places implies  $|\omega| = 1, \operatorname{Re} \omega = \frac{1}{2}, n_1n_2 \leq 0$ , and  $|n_1| = |n_2|$ . Hence,  $n_2 = -n_1 = \pm 1$ , and  $a = \pm(\omega - 1)$ .  $|\omega| = |\omega - 1| = 1$  implies that  $\omega = (1 + \sqrt{3}i)/2$ . If  $a = \pm 1$ , then both  $\omega, \bar{\omega}$ , and hence  $2\operatorname{Re} \omega$  are in  $\Lambda$ . It follows that either  $\operatorname{Re} \omega = 0$ , or  $\operatorname{Re} \omega = \frac{1}{2}$ .

In summary we have the following classical result.

LEMMA 5.1. *Suppose that  $\mathbf{C}/\Lambda$  admits the reflection (5.8). Then the possibilities for  $\Lambda$  and  $a$  are*

1.  $\operatorname{Re} \omega = 0, |\omega| > 1, a = \pm 1$ ;
2.  $\operatorname{Re} \omega = \frac{1}{2}, |\omega| > 1, a = \pm 1$ ;
3.  $|\omega| = 1, 0 < \operatorname{Re} \omega < \frac{1}{2}, a = \pm \omega$ ;
4.  $\omega = i, a = \pm 1, \pm i$ ;
5.  $\omega = (1 + \sqrt{3}i)/2, a = \pm 1, \pm \omega, \pm(\omega - 1)$ .

In particular, it follows that  $J(\omega)$ , the elliptic modular function [5], is real at  $\omega$ . In each case one has to determine the possible reflections  $\rho$ , determine their fixed-point sets, and add a suitable  $\tau_1$ .

We consider the rectangular case (1) of the lemma, for application in the next section. Let

$$(5.19) \quad \omega_1 = 1, \omega_2 = \omega = i\omega'', \omega'' > 1$$

be a normalized basis. For  $a = 1$ ,  $l_\alpha$  is the real axis,  $\omega_0 = 0$ , or  $\omega_0 = 1$ ,  $b = ib_2$ , or  $b = \frac{1}{2} + ib_2$ ,  $0 \leq b_2 < \omega''$ . In the first case  $\omega'_0 = 0$ , or  $\omega'_0 = \omega$ , while there is no  $\omega'_0$  in the second case. Thus, we have

$$(5.20) \quad \rho(t) = \bar{t} + ib_2, FP(\rho) = \{Im t = b_2/2\} \cup \{Im t = (b_2 + \omega'')/2\}.$$

For  $a = -1$ ,  $l_\alpha$  is the imaginary axis,  $\omega_0 = 0$  or  $\omega_0 = \omega$ ,  $b = b_1$ , or  $b = b_1 + i\omega''/2$ ,  $0 \leq b_1 < 1$ .  $\omega'_0 = 0, 1$  in the first case, and there is no  $\omega'_0$  in the second case. We have

$$(5.21) \quad \rho(t) = -\bar{t} + b_1, FP(\rho) = \{Re t = b_1/2\} \cup \{Re t = (b_1 + 1)/2\}.$$

If  $\varepsilon_1 = -1$ , then

$$(5.22) \quad FP(\tau_1) = \{c_1/2, (c_1 + \omega_1)/2, (c_1 + \omega_2)/2, (c_1 + \omega_1 + \omega_2)/2\}.$$

If we have  $\varepsilon_1 = +1$ ,  $2c_1 \in \Lambda$ ,  $c_1 \notin \Lambda$ , then  $\tau_1$  has no fixed points.  $\tau_1$  is then the deck transformation of an unbranched covering of another torus.

## 6. EMBEDDING OF TORI

We turn to the problem of concretely realizing the data of the previous section in the main case. Given a complex torus  $\Gamma = \mathbf{C}/\Lambda$ , with a pair of holomorphic involutions induced by

$$(6.1) \quad \tau_i(t) = -t + c_i, i = 1, 2,$$

we look for a pair of two-fold branched coverings

$$(6.2) \quad \pi_i: \Gamma \rightarrow \mathbf{P}_1, \pi_i \circ \tau_i = \pi_i, i = 1, 2.$$

The problem is immediately solved by taking

$$(6.3) \quad z_i = \pi_i(t) \equiv \mathcal{P}(t - c_i/2), i = 1, 2,$$