

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: DOUBLE VALUED REFLECTION IN THE COMPLEX PLANE
Autor: Webster, S. M.
Kapitel: 7. A RECTANGULAR LATTICE
DOI: <https://doi.org/10.5169/seals-87870>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

To satisfy (3.11) we *redefine*

$$(6.18) \quad \pi_i(t) = a \mathcal{P}(t - c_i/2),$$

and set

$$(6.19) \quad G_0(z_1, z_2) = a^4 G(z_1/a, z_2/a),$$

so that

$$(6.20) \quad \overline{G_0(z_1, z_2)} = G_0(\bar{z}_2, \bar{z}_1).$$

In summary we have

PROPOSITION 6.1. *Let $\Lambda = \mathbf{C}/\Lambda$ have the holomorphic involutions (6.1) intertwined by the anti-holomorphic involution (5.6). Then (Γ, ρ, τ_i) is realized by the map (6.5), (6.18) onto the quartic curve $G_0(z_1, z_2) = 0$ given by (6.13), (6.14), (6.19). If the fixed-point set of ρ is non-empty, then this is the complexification of the real curve $G_0(z, \bar{z}) = 0$.*

7. A RECTANGULAR LATTICE

We consider the special case of Λ, ρ, τ_i as given in (5.19), (5.6), (6.1), with

$$(7.1) \quad a = +1, b = 0, \bar{c}_2 = c_1 = c'_1 + ic''_1, c = ic''_1.$$

From (6.16), (6.15) it follows that g_2, g_3, β are real, and β' is purely imaginary. Thus, the coefficients $\beta_1, \beta_2, \beta_3$ of $G(z_1, z_2)$ are real. With $t = t' + it''$, we have

$$(7.2) \quad FP(\rho) = \{t'' = 0\} \cup \{t'' = \omega''/2\},$$

$$(7.3) \quad \tau_1\{t'' = 0\} = \{t'' = c''_1\}, \tau_1\{t'' = \omega''/2\} = \{t'' = c''_1 + \omega''/2\}.$$

Let us assume that $0 < c''_1 < \omega''/2$. Then the torus Λ is divided into four annuli

$$A_1 = \{0 < t'' < c''_1\}, A_2 = \{c''_1 < t'' < \omega''/2\},$$

$$A_3 = \{\omega''/2 < t'' < c''_1 + \omega''/2\}, A_4 = \{c''_1 + \omega''/2 < t'' < \omega''\}.$$

The fixed points of τ_1 are, by (5.22),

$$(7.4) \quad c_1/2, (c_1 + 1)/2 \in A_1,$$

$$(7.5) \quad (c_1 + i\omega'')/2, (c_1 + 1 + i\omega'')/2 \in A_3.$$

For the map $z = \pi_1(t)$ (6.3), we get

$$(7.6) \quad D_j = \pi_1(A_j), C_j = \partial D_j, 1 \leq j \leq 4 .$$

Then the z -plane is the disjoint union of $D_1, D_2 = D_4, D_3, C_1$, and C_3 . π_1 maps each of A_2 and A_4 biholomorphically onto D_2 , which is topologically an annulus with boundary $C_2 = C_1 - C_3$. π_1 also gives twofold branched coverings of A_i onto $D_i, i = 1, 3$, branched at (7.4), (7.5). In particular, D_1 is unbounded, containing $\pi_1(c_1/2) = \infty$ in its interior, and C_1 and C_3 are symmetric with respect to the real axis. $\pi_1(t)$ is real on the two horizontal lines through (7.4) and (7.5). It is also real on the two vertical lines $\{t' = c'_1/2\}, \{t' = (c'_1 + 1)/2\}$, which intersect ∂A_1 in the points

$$(7.7) \quad a_1 = \frac{1}{2} c'_1, a_2 = \frac{1}{2} c'_1 + ic''_1 ,$$

and

$$(7.8) \quad b_1 + 1 = \frac{1}{2} (c'_1 + 1), b_2 = \frac{1}{2} (c'_1 + 1) + ic''_1 .$$

$\pi_1(a_1) = \pi_1(a_2) \in C_1 \cap \mathbf{R}$ and $\pi_1(b_1 + 1) = \pi_1(b_2) \in C_1 \cap \mathbf{R}$ are the “vertices” of C_1 .

The “annular” domain D_2 has the same conformal type as A_2 , which is determined by $\frac{1}{2} \omega'' - c''_1$. This depends on both Λ and τ_1 .

Finally we consider the Riemann map, $\zeta = f(z)$, of D_1 onto the right half plane H , which takes $\pi_1(a_1)$ to zero and $\pi_1(b_1 + 1)$ to ∞ . We extend $f \circ \pi_1(t)$ to the entire t -plane by reflection in the lines $\{t'' = 0\}$ and $\{t'' = c''_1\}$. This gives a doubly periodic meromorphic function $\hat{\phi}$ with period module

$$(7.9) \quad \hat{\Lambda} = \{n_1 \cdot 1 + n_2 \cdot 2c''_1 i \mid n_1, n_2 \in \mathbf{Z}\} .$$

$\hat{\phi}$ has the representation in terms of the sigma function \hat{S} relative to $\hat{\Lambda}$,

$$(7.10) \quad \hat{\phi} = \frac{\hat{S}(t - a_1) \hat{S}(t - a_2)}{\hat{S}(t - b_1) \hat{S}(t - b_2)} .$$

The invariance $\hat{\phi} \circ \tau_1 = \hat{\phi}$ follows as in section 4, since $\tau_1(a_1) = a_2, \tau_1(b_1) = b_2$. Since $z = \mathcal{P}(t - c_1/2)$, we have

THEOREM 7.1. *Let Λ be the lattice with periods (5.19), and let $D_1 \subset \mathbf{P}_1$ be the simply connected domain above. The Riemann map, $\zeta = f(z)$, of D_1 onto the right half ζ -plane H is given by*

$$(7.11) \quad \zeta = \hat{\phi}(\mathcal{P}^{-1}(z) + c_1/2) ,$$

where $\hat{\phi}$ is the sigma quotient (7.10) relative to the lattice (7.9), and $\mathcal{P}^{-1}(z)$ is the elliptic integral of the first kind, in Weierstrass normal form, relative to Λ .

REMARK. We have seen that double valued reflection places a severe restriction on a real algebraic curve in the complex plane. In fact our results should provide the basis for a complete and explicit classification. We have also seen how double valued reflection may be used to explicitly determine Riemann maps. Apparently, all known such examples can be so explained. The result in the above theorem seems to be new. It would be interesting to work out more examples in the genus one case.

REFERENCES

- [1] CARATHEODORY, C. *Conformal Representation*. Cambridge University Press (1952).
- [2] COURANT, R. and D. HILBERT. *Methods of Mathematical Physics*, vol. I. Interscience, New York (1953).
- [3] DU VAL, P. *Elliptic Functions and Elliptic Curves*. Cambridge University Press (1973).
- [4] FARKAS, H. and I. KRA. *Riemann Surfaces*. Springer-Verlag (1980).
- [5] FRICKE, R. *Elliptische Functionen*. B.G. Teubner, Leipzig und Berlin (1916).
- [6] HURWITZ, A. and R. COURANT. *Functionentheorie*. Springer-Verlag (1964).
- [7] KELLOGG, O.D. *Foundations of Potential Theory*. Dover Publications, New York (1954).
- [8] MOSER, J.K. and S.M. WEBSTER. Normal forms for real surfaces in \mathbb{C}^2 near complex tangents and hyperbolic surface transformations. *Acta Math.* 150 (1983), 255-296.
- [9] NEHARI, Z. *Conformal Mapping*. Dover Publications, New York (1975).
- [10] SCHWARZ, H.A. *Gesammelte Mathematische Abhandlungen*, vol. II. Berlin (1890).

(Reçu le 15 mai 1995)

S.M. Webster

University of Chicago
 Department of Mathematics
 5734 University Ave.
 Chicago, IL 60637
 U.S.A.