

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	42 (1996)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	CENTRALISERS IN THE BRAID GROUP AND SINGULAR BRAID MONOID
Autor:	Fenn, Roger / Zhu, Jun
Kapitel:	4. Centralisers of braid subgroups
DOI:	https://doi.org/10.5169/seals-87872

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

iterated $2r$ times. Noting that I_p persists in w^* it is easy to argue that $w(A * \sigma_j^{2r}) = w(A)$ is impossible; the contradiction. \square




FIGURE 4
The action of $* \sigma_j^{2r}$ on a (j, k) -arc in case $r = 2$

We now turn to the proof of Theorem 2.2. It has already been observed that $(e) \Rightarrow (d) \Rightarrow (a)$, and it is obvious that $(a) \Rightarrow (c) \Rightarrow (b)$. So it remains to establish that $(b) \Rightarrow (e)$. Thus we assume that, for some $r \neq 0$, $\sigma_j^r \beta = \beta \sigma_k^r$. Since the algebraic crossing number of any two strings of a braid is a well-defined braid invariant, this equation is possible only if $\{j, j+1\} * \beta = \{k, k+1\}$. Now, noting that $\beta^{-1} \sigma_j^r \beta = \sigma_k^r$ and that σ_k^r has a (k, k) -band, we conclude that there is a proper ribbon for $\beta^{-1} \sigma_j^r \beta$ from $[k, k+1] \times 0$ to $[k, k+1] \times 1$. Define $A = \beta * [k, k+1] = [k, k+1] * \beta^{-1}$. Then we may assume (possibly after an isotopy) that the planes $\mathbf{C} \times 1/3$ and $\mathbf{C} \times 2/3$ cut the ribbon in the arcs $A \times 1/3$ and $A \times 2/3$. Moreover, the middle third of the ribbon, and Proposition 1.1, imply that $A * \sigma_j^r = A$. By Lemma 3.2, $A = [j, j+1]$ and the theorem is proved. \square

4. CENTRALISERS OF BRAID SUBGROUPS

We have established the following.

4.1 THEOREM. *The centraliser in B_n of the generator σ_j is the subgroup of all braids which have (j, j) -bands. This subgroup is isomorphic to $B_{n-1}^j \times \mathbf{Z}$ where B_{n-1}^j is the subgroup of B_{n-1} consisting of all $(n-1)$ -braids whose permutations stabilise j .* \square

The goal of this section is to describe the centraliser of B_r in B_n , $r \leq n$, which we will call $C(r, n)$. Here B_r is the r -string braid group with its usual inclusion in B_n , namely as the subgroup generated by $\sigma_1 \dots \sigma_{r-1}$.

4.2 THEOREM. *The centraliser $C(r, n)$ of B_r in B_n consists of all n -braids in which the first r strings lie on a ribbon, disjoint from the other strings, and which intersects $\mathbf{C} \times 0$ and $\mathbf{C} \times 1$ in exactly the straight line intervals from $[1, r] \times 0$ and $[1, r] \times 1$ (up to isotopy).*

Proof. A braid β is in $C(r, n)$ if and only if it commutes with each σ_j , $1 \leq j \leq r - 1$. Thus $[j, j + 1] * \beta = [j, j + 1]$, $1 \leq j \leq r - 1$ and so $[1, r] * \beta = [1, r]$, up to isotopy fixing $\{1, \dots, n\}$. \square

It follows that $C(r, n)$ consists of all n -braids constructible as follows. Let $k = n - r + 1$ and consider the subgroup B_k^1 of k -braids whose associated permutation fixes 1. Then replace the first string of a braid in B_k^1 by r parallel strings lying on a ribbon along that string. The ribbon may be twisted by some integral multiple of 2π (or π in the case $r = 2$); such braids are precisely the central elements of B_r .

4.3 THEOREM. *The centraliser $C(r, n)$ is isomorphic to the direct product $B_{n-r+1}^1 \times \mathbf{Z}$.* \square

A PRESENTATION OF $C(r, n)$. In order to establish a set of generators and defining relations for $C(r, n)$ we recall results of Chow [Ch] regarding B_k^1 . This subgroup of B_k is generated by $\sigma_2, \dots, \sigma_{k-1}$, together with elements a_2, \dots, a_k defined by

$$a_i := \sigma_1^{-1} \sigma_2^{-1} \cdots \sigma_{i-2}^{-1} \sigma_{i-1}^2 \sigma_{i-2} \cdots \sigma_2 \sigma_1.$$

These generators satisfy the usual braid relations:

$$\sigma_i \sigma_j = \sigma_j \sigma_i, \quad |i - j| > 1$$

$$\sigma_i \sigma_{i+1} \sigma_i = \sigma_{i+1} \sigma_i \sigma_{i+1}$$

as well as the following, for $i = 2, \dots, k - 1$:

$$\begin{aligned} \sigma_i a_j \sigma_i^{-1} &= a_j, \quad j \neq i, i + 1 \\ \sigma_i a_i \sigma_i^{-1} &= a_{i+1} \\ \sigma_i a_{i+1} \sigma_i^{-1} &= a_{i+1}^{-1} a_i a_{i+1}. \end{aligned}$$

In fact these are *defining* relations for B_k^1 . Chow also noted that the subgroup of B_k^1 generated by the a_i is a normal subgroup (as is clear from the above relations), in fact a *free* group on the generators a_i , and that B_k^1 could be regarded as the semidirect product of that free subgroup with the

subgroup generated by $\sigma_2 \dots \sigma_{k-1}$, the latter group clearly isomorphic with the braid group on $k - 1$ strings.

Applying this to our situation, for each $i = 1, \dots, n - r$, let A_{r+i} be the n -braid resulting from replacing the first string of the k -braid a_i , defined above, by r parallel strings which lie on an untwisted band. Specifically,

$$\begin{aligned} A_{r+i} = & (\sigma_r^{-1} \sigma_{r+1}^{-1} \cdots \sigma_{r+i-2}^{-1} \sigma_{r+i-1}) (\sigma_{r-1}^{-1} \sigma_r^{-1} \cdots \sigma_{r+i-3}^{-1} \sigma_{r+i-2}) \\ & \cdots (\sigma_1^{-1} \sigma_2^{-1} \cdots \sigma_{i-1}^{-1} \sigma_i) \times (\sigma_i \sigma_{i-1} \cdots \sigma_1) (\sigma_{i+1} \sigma_i \cdots \sigma_2) \\ & \cdots (\sigma_{r+i-1} \sigma_{r+i-2} \cdots \sigma_r) \end{aligned}$$

Also let C denote the well-known generator of the centre of the r -string braid group, namely $C = \sigma_1$ if $r = 2$ and in case $r > 2$:

$$C = (\sigma_1 \sigma_2 \cdots \sigma_{r-1})^r.$$




FIGURE 5
Special generators of $C(r, n)$

4.4 THEOREM. *The centraliser $C(r, n)$ of B_r in B_n has the generators:*

$$\sigma_{r+1}, \sigma_{r+2}, \dots, \sigma_{n-1}, A_{r+1}, \dots, A_n, C$$

and defining relations:

$$\begin{aligned} \sigma_i \sigma_j &= \sigma_j \sigma_i, \quad |i - j| > 1 \\ \sigma_i \sigma_{i+1} \sigma_i &= \sigma_{i+1} \sigma_i \sigma_{i+1} \\ \sigma_i A_j \sigma_i^{-1} &= A_j, \quad j \neq i, i+1 \\ \sigma_i A_i \sigma_i^{-1} &= A_{i+1} \\ \sigma_i A_{i+1} \sigma_i^{-1} &= A_{i+1}^{-1} A_i A_{i+1} \\ C \sigma_i &= \sigma_i C \\ C A_i &= A_i C. \end{aligned}$$

(Subscripts ranging over all values for which the symbols are in the list of generators.) \square