

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: UNIFORM DISTRIBUTION ON DIVISORS AND BEHREND SEQUENCES
Autor: Tenenbaum, Gérald
Kapitel: 3. Functions of excessive growth: the case $f(d) = d^\alpha$
DOI: <https://doi.org/10.5169/seals-87875>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

We choose $T = \varphi_1(x)^{y/4} = (\log x / \varphi(x))^{y/4}$. The upper bound above becomes

$$(56) \quad \ll \varphi(x)^{y/2} (\log x)^{y/2-1} (\log \varphi_1(x))^2.$$

Indeed the last term is easily seen to be negligible by the lower bound (40) imposed on $\varphi(x)$, and because $R(x) = \exp\{(\log x)^{4/7}\}$ is an admissible choice for R . Thus we may define $E_2(x, y)$ as being equal to a suitable constant multiple of the right-hand side of (56), and apply Theorem 7 to obtain that

$$(57) \quad \Delta(n; f) \ll \xi(n) \tau(n) y^{\Omega(n)/2} \sqrt{E_2(n, y)} \quad \text{ppl},$$

provided $0 \leq y \leq y_0 < 4$ and $y = y(n)$ is such that $E_2(n, y)$ is slowly increasing as a function of n . We choose

$$y = 2 \left/ \left(1 + \frac{\log \varphi(n)}{\log_2 n} \right) \right. = 1 \left/ \left(1 - \frac{\log \varphi_1(n)}{2 \log_2 n} \right) \right.,$$

which minimises $(\log n)^{-\log y} E_2(n, y)$ up to a power of $\log_2 \varphi_1(n)$. This value of y is always in the range $[1, 2]$. Inserting into (56) yields

$$E_2(n, y) \asymp (\log \varphi_1(n))^2,$$

which implies that this function is slowly decreasing. The required estimate (42) hence follows from (57). This completes the proof of Theorem 10.

3. FUNCTIONS OF EXCESSIVE GROWTH: THE CASE $f(d) = d^\alpha$

Here, we address the problem of bounding the discrepancy *ppl* for functions which increase too fast for the techniques of the previous section to be applicable. More precisely, let us recall the quantity

$$(58) \quad H_v(x, y) := \sum_{k=1}^{\infty} \left(\frac{y}{4} \right)^{\Omega(k)} \frac{1}{k^{1+\sigma}} \left| \sum_{m=1}^{\infty} \left(\frac{y}{4} \right)^{\Omega(m)} \frac{e(\nu f(km))}{m^{1+\sigma}} \right|^2,$$

with $\sigma := 1/\log x$, which appears implicitly in the upper bound (26) of Theorem 7 for the discrepancy $\Delta(n; f)$. This was primarily defined for $y < 8$, but we restrict it here to values of $y \leq 4$. The functions of moderate growth are essentially those for which the inner m -sum can be estimated by partial summation, using the available results on the mean value of $m \mapsto (y/4)^{\Omega(m)}$. When the rate of growth of f prohibits such a treatment, we may consider $H_v(x, y)$ as a 'type II sum', according to the poetic

terminology of Vinogradov. For all intents and purposes, this means making the trivial estimate $|(y/4)^{\Omega(k)}| \leq 1$, expanding the square and, after permuting summations, estimating the inner k -sum by an *ad hoc* exponential sum method.

This programme may be carried out, in principle, for any smooth function f of, say, at most polynomial growth, and indeed one could even aim at a general theorem established along these lines and providing, under suitable sufficient conditions, explicit upper bounds for the discrepancy. Due to the considerable amount of calculations that this would involve, we have preferred to treat only examples which reflect all the difficulties of the general case, but avoid tedious technicalities that would hide the main stream of the argument. In this context, we believe that the functions

$$d \mapsto d^\alpha \quad (\alpha \in \mathbf{R}^+ \setminus \mathbf{Z}^+), \quad d \mapsto \theta d \quad (\theta \in \mathbf{R} \setminus \mathbf{Q})$$

are of special interest. We treat the first of these immediately and the second one in the next section. The following theorem provides an effective version of the corresponding qualitative result obtained by Hall & Tenenbaum in [13].

THEOREM 11. *Let $\alpha > 0$ be a given real number, not an integer. Then the function $g_\alpha(d) := d^\alpha$ is erd. More precisely, we have*

$$(59) \quad \Delta(n; g_\alpha) < \tau(n) (\log n)^{-\delta} \quad \text{ppl}$$

for all $\delta < \delta_0 := \frac{1}{2} \log \frac{12}{11}$.

We note that $\delta_0 > \frac{1}{23}$. We have not attempted here to find the best exponent available from latest developments in exponential sums theory and have confined ourselves to using a result of Karatsuba [16] on Vinogradov-type bounds which is expressed in an easily applicable form. We remark that van der Corput-type estimates would in general yield weaker bounds for the discrepancy and would actually only save a power of $\log_2 n$ in (59).

It is also worthwhile to note at this stage that the method of proof of Theorem 11 will readily yield that the function

$$f(d) = \exp\{(\log d)^\alpha\}$$

is erd for $0 < \alpha < 1$, and indeed it will provide a bound comparable with (59) for the discrepancy. In particular, this shows that the limitation $\alpha < 3/5$ which arises from mere application of Theorems 4 or 10 is purely technical. The range $1 < \alpha < 3/2$ may also be handled by the

same technique, but with a weaker effective result — see Theorem 2 of Karatsuba [16].

By Theorem 3, we immediately derive from the above result the following corollary.

COROLLARY 5. *Let α, δ be as in the statement of Theorem 11. Then the sequence*

$$(60) \quad \{n \geq 2 : \langle n^\alpha \rangle \leq (\log n)^{-\delta}\}$$

is a Behrend sequence.

Of course we can rewrite the condition in (60) introducing $j := [n^\alpha]$. This yields the following reformulation in terms of block sequences.

COROLLARY 6. *Let $\beta > 0$, $1/\beta \notin \mathbf{Z}$, $\delta < \delta_0 = \frac{1}{2} \log \frac{12}{11}$. Then the sequence*

$$(61) \quad \mathcal{B}(\delta) := \bigcup_{j=2}^{\infty} \left[j^\beta, j^\beta \left(1 + \frac{1}{j(\log j)^\delta} \right) \right] \cap \mathbf{Z}^+$$

is a Behrend sequence.

As far as block sequences are concerned, this is only significant when $\beta > 1$: otherwise the ‘blocks’ have lengths smaller than 1 and looking at $\mathcal{B}(\delta)$ as a block sequence is meaningless. As we remarked in the previous section, the above result is unreachable, in the present state of knowledge, by the technique applied in [24]. The natural conjecture in accord with the results of [15] and [24] would be that $\mathcal{B}(\delta)$ is Behrend for all $\delta < \log 2$, this exponent then being optimal. This is also out of reach of the present technique, which implies a systematic loss due, among other causes, to the trivial estimate for $(y/4)^{\Omega(k)}$ in (58).

We now embark on the proof of Theorem 11. We give ourselves two parameters x_1, x_2 satisfying

$$e^{\sqrt{\log x}} \leq x_1 \leq x, \quad x_2 := x^{\log_2 x},$$

and introduce the following further notation

$$J = J(x) := \frac{\log(x_2/x_1)}{\log 2}, \quad K_j := 2^j x_1 \quad (0 \leq j \leq J),$$

$$B_j(m, n; v, f) := \sum_{K_j < k \leq K_{j+1}} e(vf(kn) - vf(km)) \quad (0 \leq j \leq J).$$

LEMMA 1. Let $\alpha > 0$. Then we have uniformly for $x \geq 3, v \geq 1, 0 \leq y \leq 4$, and a real valued arithmetical function f

$$(62) \quad H_v(x, y) \ll (\log x)^{y/2} (\log x_1)^{y/4} + \sum_{\substack{x_1 < m < n \leq x_2 \\ n^\alpha - m^\alpha \geq 1}} \frac{\left(\frac{1}{4}y\right)^{\Omega(mn)}}{mn} \sum_{0 \leq j \leq J} \frac{\Re B_j(m, n; v, f)}{K_j}.$$

Proof. To lighten the presentation, we temporarily set $z := y/4$. We first split the k -sum in (58) according to whether $k \leq x_1, x_1 < k < x_2$ or $k > x_2$, so as to write correspondingly

$$H_v(x, y) = H_v^{(1)}(x, y) + H_v^{(2)}(x, y) + H_v^{(3)}(x, y).$$

Using the bounds

$$(63) \quad \sum_{k \leq x_1} z^{\Omega(k)} k^{-1} \ll (\log x_1)^z,$$

and

$$(64) \quad \sum_{k > x_2} z^{\Omega(k)} k^{-1-\sigma} \ll x_2^{-\sigma/2} \sum_{k \geq 1} z^{\Omega(k)} k^{-1-\sigma/2} \ll (\log x)^{z-1/2},$$

we readily obtain

$$(65) \quad H_v^{(1)}(x, y) + H_v^{(3)}(x, y) \ll (\log x_1)^z (\log x)^{2z} + (\log x)^{3z-1/2} \ll (\log x)^{y/2} (\log x_1)^{y/4}.$$

Next, we split the inner m -sum in $H_v^{(2)}(x, y)$ at x_1 and x_2 and use the inequality $(a + b + c)^2 \leq 3(a^2 + b^2 + c^2)$ to obtain

$$H_v^{(2)}(x, y) \leq 3H_v^{(21)}(x, y) + 3H_v^{(22)}(x, y) + 3H_v^{(23)}(x, y)$$

with

$$(66) \quad H_v^{(21)}(x, y) \leq \sum_{x_1 < k \leq x_2} z^{\Omega(k)} k^{-1} \left(\sum_{m \leq x_1} z^{\Omega(m)} m^{-1} \right)^2 \ll (\log x)^z (\log x_1)^{2z} \ll (\log x)^{y/2} (\log x_1)^{y/4},$$

by (63), and

$$(67) \quad H_v^{(23)}(x, y) \leq \sum_{x_1 < k \leq x_2} z^{\Omega(k)} k^{-1} \left(\sum_{m > x_2} z^{\Omega(m)} m^{-1-\sigma} \right)^2 \ll (\log x)^{3z-1} = (\log x)^{3y/4-1} \ll (\log x)^{y/2},$$

by (64).

It remains to estimate $H_v^{(22)}(x, y)$. We have

$$(68) \quad H_v^{(22)}(x, y) \leq \sum_{0 \leq j \leq J} K_j^{-1} \sum_{K_j < k \leq K_{j+1}} \left| \sum_{x_1 < m \leq x_2} \frac{z^{\Omega(m)}}{m^{1+\sigma}} e(vf(km)) \right|^2,$$

where we have made the trivial estimate $z^{\Omega(k)} \leq 1$. Expanding the square, we find that it does not exceed

$$(69) \quad 2\Re e \sum_{\substack{m < n \leq x_2 \\ n^\alpha - m^\alpha \geq 1}} \frac{z^{\Omega(mn)}}{(mn)^{1+\sigma}} e(vf(kn) - vf(km)) + 2 \sum_{\substack{x_1 < m \leq n \leq x_2 \\ n^\alpha - m^\alpha < 1}} \frac{1}{mn}.$$

We claim that the second sum on the right is $\ll x_1^{-\min(1, \alpha)}$. This plainly holds if $\alpha \geq 1$ since the summation conditions then imply that $m = n$. When $0 < \alpha < 1$, we note that $n^\alpha - m^\alpha \geq \alpha(n - m)n^{\alpha-1}$, so for fixed n the m -sum is $\ll n^{1-\alpha}/n = n^{-\alpha}$ and the conclusion is still valid. Inserting this estimate into (69) and (68) and using the fact that $J \ll x_1^{\min(1, \alpha)}$, we obtain

$$H_v^{(22)}(x, y) \ll \Re e \sum_{0 \leq j \leq J} \frac{1}{K_j} \sum_{K_j < k \leq K_{j+1}} \sum_{\substack{x_1 < m < n \leq x_2 \\ n^\alpha - m^\alpha \geq 1}} \frac{z^{\Omega(mn)}}{(mn)^{1+\sigma}} e(vf(kn) - vf(km)) + 1.$$

We permute summations on k and m, n and see that the new, inner k -sum equals $B_j(m, n; v, f)$. Together with (65), (66) and (67), this completes the proof of our lemma.

We now apply Karatsuba's estimate to bound the exponential sum

$$(70) \quad B(K; v) := \sum_{K < k \leq 2K} e(vk^\alpha)$$

for relevant values of v, K .

LEMMA 2. *Let $\alpha \in \mathbf{R}^+ \setminus \mathbf{Z}^+$. There exists a constant $c = c(\alpha) > 0$ such that the estimate*

$$(71) \quad B(K; v) \ll K^{1-c} + K \exp \left\{ -c \frac{(\log K)^3}{(1 + \log v)^2} \right\}$$

holds uniformly for $K \geq 1, v \geq 1$.

Proof. If $v \leq K^{1-\alpha}$, and so $0 < \alpha < 1$, we apply a classical estimate of van der Corput (see e.g. Titchmarsh [26], lemma 4.7) to get

$$B(K; v) = \int_K^{2K} e(vt^\alpha) dt + O(1) \ll K^{1-\alpha}.$$

(The same estimate also follows from theorem 2.1 of Graham & Kolesnik [7].)

Thus (71) holds in this case.

If $v > K^{1-\alpha}$, set $n := [3\alpha + 3(\log v)/\log K]$, so that $n \geq 3$ and

$$K^{n/3} \leq vK^\alpha < K^{(n+1)/3}.$$

Put $g(t) := vt^\alpha$. We have for all non-negative integers s

$$\frac{g^{(s)}(t)}{s!} = v \binom{\alpha}{s} t^{\alpha-s}.$$

Writing $\binom{\alpha}{s} = (-1)^s \prod_{1 \leq j \leq s} \{1 - (\alpha + 1)/j\}$, we see that we have for suitable positive constants $c_1 = c_1(\alpha)$, $c_2 = c_2(\alpha)$,

$$(72) \quad c_1 s^{-\alpha-1} \leq \left| \binom{\alpha}{s} \right| \leq c_2 s^{-\alpha-1} \quad (s \geq 0).$$

Hence for large K and $K \leq t \leq 2K$ we have

$$\left| \frac{g^{(n+1)}(t)}{(n+1)!} \right| \leq \frac{2^\alpha c_2}{(n+1)^{\alpha+1}} v K^{\alpha-n-1} \leq 2^\alpha c_2 K^{-2(n+1)/3} \leq K^{-(n+1)/2}.$$

Similarly, a straightforward computation enables us to deduce from (72) that for all s in the range $3n/4 \leq s \leq n$ (so $s \geq 3$) and large K we have

$$K^{-3s/4} \leq c_1 s^{-\alpha-1} 2^{-s} K^{-2/3} \leq \left| \frac{g^{(s)}(t)}{s!} \right| \leq 2^\alpha K^{-5s/9+1/3} \leq K^{-s/3}.$$

By Theorem 1 of Karatsuba [16], it follows that, for suitable positive absolute constants c_3 and c_4 and $K > K_0(\alpha)$ we have

$$|B(K; v)| \leq c_3 K^{1-c_4/n^2}.$$

This implies the required bound.

We are now in a position to complete the proof of Theorem 11. We want to apply Theorem 7 and use Lemmas 1 and 2 to obtain an upper bound for the quantity $E_2(n, y)$. We select, in Lemma 1, $f = g_\alpha$, as defined in the statement of the theorem, and

$$x_1 = \exp\{c(\log x)^{2/3} \log_2 x\},$$

with $c = c(\alpha) > 0$ at our disposal. Then, with the notation of Lemma 2, we have $B_j(m, n; v, g_\alpha) = B(K_j; v)$ where $v := v(n^\alpha - m^\alpha)$, so $v \leq x_2^{1+\alpha}$ provided $v \leq x$. By Lemma 2 there is a positive constant $c_5 = c_5(\alpha)$ such that, for all $m, n \leq x$ with $n^\alpha - m^\alpha \geq 1$,

$$B_j(m, n; v, g_\alpha) / K_j \ll \exp \left\{ -c_5 \frac{(\log K_j)^3}{(\log x_2)^2} \right\} \ll 1 / (\log x)^4$$

provided $c(\alpha)$ is large enough. By (62), we infer that we have uniformly for $1 \leq v \leq x$, $0 \leq y \leq 4$,

$$(73) \quad H_v(x, y) \ll (\log x)^{2y/3} (\log_2 x)^{y/4}.$$

Inserting the above estimate into (24) with, say, $T := \log x$, we see that we may choose

$$(74) \quad E_2(x, y) \asymp (\log x)^{11y/12 - 1} (\log_2 x)^{2 + y/4},$$

which is hence slowly increasing. Therefore we get by Theorem 7 that

$$(75) \quad \Delta(n, g_\alpha) < \tau(n) (\log n)^{11y/24 - 1/2 - (1/2)\log y + o(1)} \text{ ppl}.$$

The required estimate (59) now follows on taking optimally $y = \frac{12}{11}$.

4. FUNCTIONS OF EXCESSIVE GROWTH: THE CASE $f(d) = \theta d$

We now investigate, in a quantitative form, the uniform distribution on divisors of the function

$$h_\theta(d) := \theta d$$

when θ is a given irrational real number. This study is similar in principle to that of the previous section, but more complicated inasmuch as the effective bounds for $\Delta(n; h_\theta)$ will depend on the arithmetic nature of θ . On the other hand we shall not need, as might be expected, any involved tool for the estimation of the relevant exponential sums.

More explicitly, let us define $Q(x) := x / (\log x)^{10}$, and

$$(76) \quad q(x; \theta) := \inf \{ q : 1 \leq q \leq Q(x), \|q\theta\| \leq 1/Q(x) \}$$

where $\|u\|$ denotes the distance of u to the set of integers. Our results depend on a free parameter y , $0 < y \leq 4$, and may be expressed conveniently in terms of any increasing lower bound for $q(x; \theta)$, say $q^*(x; y, \theta)$, with the property that $q^*(x; y, \theta) / (\log x)^{y/4}$ is decreasing. A possible choice is

$$(77) \quad q^*(x; y, \theta) := 4(\log x)^{y/4} / \int_1^x \frac{y(\log t)^{y/4 - 1}}{t \inf_{u \geq t} q(u; \theta)} dt.$$

Unless θ has abnormally good rational approximations, we have

$$(78) \quad q^*(x; y, \theta) \asymp (\log x)^{y/4}.$$