

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: UNIFORM DISTRIBUTION ON DIVISORS AND BEHREND SEQUENCES
Autor: Tenenbaum, Gérald
Kapitel: §5. Additive functions
DOI: <https://doi.org/10.5169/seals-87875>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

§5. ADDITIVE FUNCTIONS

As we noted in the first section, the study of uniform distribution on divisors of additive functions is made much easier by the fact that the Weyl sums are multiplicative. Indeed this is the only case when we are able to achieve estimates for the discrepancy which go beyond the statistical bound $\sqrt{\tau(n)}$.

We shall prove the following theorem, which provides a simple, but nevertheless effective criterion. We write for integer v

$$(102) \quad L_v(x; f) := \sum_{p \leq x} \frac{\|vf(p)\|^2}{p}, \quad C_v(x; f) = \sum_{p \leq x} \frac{1 - |\cos v\pi f(p)|}{p},$$

and note that we have for all x

$$(103) \quad 4L_v(x; f) \leq C_v(x; f) \leq \frac{1}{2} \pi^2 L_v(x; f).$$

THEOREM 13. *Let f be an additive function. Then f is erd if, and only if,*

$$(104) \quad \sum_p \frac{\|vf(p)\|^2}{p} = \infty \quad (v \neq 0).$$

Moreover, if this is the case then we have

$$(105) \quad \sum_{n \leq x} \frac{\Delta(n; f)}{n} \left(\frac{y}{2}\right)^{\Omega(n)} \ll (\log x)^y \left\{ \frac{1}{T} + \sum_{1 \leq v \leq T} \frac{e^{-yC_v(x; f)}}{v} \right\}$$

uniformly for $x \geq 2$, $T \geq 1$, $0 \leq y \leq y_0 < 4$.

The qualitative result corresponding to this statement (i.e. the criterion (104)) was first established by Kátai [18].

By (104) and the lower bound of (103), the expression between curly brackets tends to zero for suitable $T = T(x, y)$. Hence (105) provides, as described in Theorem 6, a family of effective ppl estimates for $\Delta(n; f)$ which may of course be further optimised with respect to y . This process must lead to a non trivial result since $y = 1$ is admissible.

We give below two applications of Theorem 13, respectively devoted to the functions $f(d) = \theta\Omega(d)$ for irrational θ , and $f(d) = \log d$. Recovering a result first obtained independently by Hall and Kátai, the latter case furnishes the best known ppl upper bound for the discrepancy for any function we are aware of, although this falls short of the current conjecture stated in section 2. The former case provides an example of a function which

is erd but has the same rate of growth than, and indeed is asymptotically equal to, a function which is not, namely $\theta \log_2 n$ — see Corollary 3. Actually, if we define $\Omega(n; E)$ as the number of those prime factors of n which belong to some set of primes E , then Theorem 13 implies that $\theta\Omega(n; E)$ is erd for all $\theta \in \mathbf{R} \setminus \mathbf{Q}$ if, and only if, $\sum_{p \in E} 1/p = \infty$. We thus exhibit erd additive functions with an arbitrarily slow growth.

The effective bounds for $\Delta(n; \theta\Omega)$ naturally depend on the rational approximations to θ . We set $Q_1(x) := \sqrt{\log_2 x / \log_3 x}$, and define

$$(106) \quad q_1(x; \theta) := \inf_{t \geq x} \max \{q \leq Q_1(t) : \|\theta q\| \leq 1/Q_1(t)\} .$$

Then $q_1(x; \theta)$ increases to ∞ for all irrational θ and furthermore $q_1(x; \theta) = (\log_2 x)^{1/2 + o(1)}$ for almost all θ .

COROLLARY 10. *Let $\theta \in \mathbf{R} \setminus \mathbf{Q}$. Then the function $\theta\Omega(n)$ is erd and we have*

$$(107) \quad \Delta(n; \theta\Omega) < \tau(n) q_1(n; \theta)^{-1 + o(1)} \quad \text{ppl} .$$

COROLLARY 11 (Hall [8]; Kátai [17]). *Let $\alpha > (\log(4/\pi))/\log 2 \approx 0.34850$. We have*

$$(108) \quad \Delta(n; \log) < \tau(n)^\alpha \quad \text{ppl} .$$

We now embark on the proof of Theorem 13. That the condition is necessary is a straightforward consequence of the definition of uniform distribution on divisors in the form (1). Indeed, suppose that (104) fails to hold for $v \neq 0$. Writing $F(z; n) := \sum_{d|n, \langle f(d) \rangle \leq z} 1$, we have

$$\begin{aligned} g_v(n) &:= \sum_{d|n} e(vf(n)) = \int_0^1 e(vz) dF(z; n) = \int_0^1 e(vz) d(F(z; n) - z\tau(n)) \\ &= 2\pi i v \int_0^1 e(vz) (F(z; n) - z\tau(n)) dz , \end{aligned}$$

hence

$$(109) \quad 2\pi |v| |\Delta(n; f)| \geq |g_v(n)| .$$

Now for all $\varepsilon \in]0, 1[$, we have

$$\begin{aligned} (110) \quad \sum_{n \leq x} \frac{\mu(n)^2 |g_v(n)|}{n\tau(n)} &\geq \sum_{P^+(n) \leq x^\varepsilon} \frac{\mu(n)^2 |g_v(n)|}{n\tau(n)} - \sum_{\substack{n > x \\ P^+(n) \leq x^\varepsilon}} \frac{1}{n} \\ &= \prod_{p \leq x^\varepsilon} \left(1 + \frac{|\cos \pi v f(p)|}{p} \right) - O(e^{-1/2\varepsilon} \log x) , \end{aligned}$$

where the O -estimate follows from (94) by partial summation. The product is

$$\gg \exp \left\{ \sum_{p \leq x^\varepsilon} \frac{|\cos \pi v f(p)|}{p} \right\} \gg \exp \left\{ \sum_{p \leq x^\varepsilon} \frac{1 - \frac{1}{2} \pi^2 \|v f(p)\|^2}{p} \right\} \gg \varepsilon \log x,$$

by our assumption that the series (104) converges. Inserting this into (110) and choosing ε small enough but fixed, we obtain that the left-hand side is $\gg \log x$, whence by (109)

$$\sum_{n \leq x} \frac{\Delta(n; f)}{n \tau(n)} \gg \log x.$$

This cannot hold if $\Delta(n; f) = o(\tau(n))$ pp (or even ppl), and we obtain the required necessity assertion.

The sufficiency part of the theorem readily follows from the upper bound (21) which we recall for convenience: we have uniformly for $x \geq 2$, $T \geq 1$, $0 \leq y \leq y_0 < 4$,

$$(111) \quad \sum_{n \leq x} \frac{\Delta(n; f)}{n} \left(\frac{y}{2}\right)^{\Omega(n)} \ll \frac{(\log x)^y}{T} + \sum_{1 \leq v \leq T} \frac{1}{v} \sum_{n \leq x} \frac{|g_v(n)|}{n} \left(\frac{y}{2}\right)^{\Omega(n)}.$$

The inner n -sum on the right does not exceed

$$\sum_{P^+(n) \leq x} \frac{|g_v(n)|}{n} \left(\frac{y}{2}\right)^{\Omega(n)} \ll \exp \left\{ \sum_{p \leq x} \frac{y |\cos \pi v f(p)|}{p} \right\} \ll (\log x)^y e^{-y C_v(x; f)}.$$

Inserting this into (111) yields (105). Taking $y = 1$ implies $\Delta(n; f) = o(2^{\Omega(n)})$ ppl, from which we deduce in turn that $\Delta(n; f) = o(\tau(n))$ ppl by a familiar argument. Corollary 1 hence implies that f is erd, and this finishes the proof of Theorem 13.

Proof of Corollary 10. We apply the upper bound (105) of Theorem 13 with $y = 1$. We have

$$C_v(x; \theta \Omega) = (1 - |\cos \pi v \theta|) \log_2 x + O(1)$$

and need a lower bound for this. For given x , there exist integers a, q with $(a, q) = 1$, $q \leq Q_1(x) = \sqrt{\log_2 x / \log_3 x}$, and $|\theta - a/q| \leq 1/q Q_1(x)$. Furthermore, we have

$$(112) \quad q \geq q_1(x; \theta),$$

where the right-hand side is defined by (106). We select $T := q / \log q$, and note that for $v \leq T$ we have $|\theta v - a_v / q_v| \leq T / q Q_1(x) \leq 1 / q \log q$, with $a_v = av / (q, v)$, $q_v = q / (q, v)$. This implies $\|\theta v\| \geq (1/q) - 1/q \log q$, hence, for large x ,

$$1 - |\cos \pi v \theta| \geq \left(\frac{1}{2} \pi^2 + o(1)\right) / q^2 > (\log_3 x) / \log_2 x.$$

Inserting this into (105), we arrive at

$$(\log x)^{-1} \sum_{n \leq x} \frac{\Delta(n; \theta \Omega)}{n 2^{\Omega(n)}} \ll \frac{1}{T} + \frac{\log q}{\log_2 x} \ll q_1(x; \theta)^{-1+o(1)}.$$

Since $q_1(x; \theta)$ is a non-decreasing function of x , this implies (107) and the proof is thereby completed.

Proof of Corollary 11. Put $\tau(n, \theta) := \sum_{d|n} d^{i\theta}$. When $f = \log$, we have that $g_v(n) = \tau(n, 2\pi v)$. By lemma 30.2 of [14] we infer that, uniformly for $1 \leq |\theta| \leq \exp \sqrt{\log x}$,

$$\sum_{p \leq x} \frac{|\tau(p, \theta)|}{p} = \sum_{p \leq x} \frac{|\cos(\frac{1}{2} \theta \log p)|}{p} = \frac{2}{\pi} \log x + O(1).$$

This is proved by partial summation from a strong form of the prime number theorem. Thus we obtain that we have uniformly for $1 \leq v \leq \log x$

$$C_v(x; \log) = (1 - 2/\pi) \log_2 x + O(1).$$

Inserting this into (105) with $T = \log x$ and choosing optimally $y = \pi/2$, we obtain

$$\Delta(n; f) < \xi(n) (\log_2 n) \left(\frac{4}{\pi}\right)^{\Omega(n)} \text{ ppl}$$

for all $\xi(n) \rightarrow \infty$. This implies the required result by a now standard device.

6. METRIC RESULTS

In this last section, we investigate the problem of uniform distribution on divisors from a further statistical point of view, regarding as random not only the integer n but also the function f . Thus, we define a measure μ on the set \mathbf{A} of all real valued arithmetical function as the inverse image of the Haar measure on the compact group $(\mathbf{R}/\mathbf{Z})^{\mathbf{N}}$ by the canonical mapping $f \mapsto \langle f \rangle$. In other words, μ is characterised by the property that for all finite families $\{E_j : 1 \leq j \leq k\}$ of measurable subsets of the torus \mathbf{R}/\mathbf{Z} and for all integers n_1, n_2, \dots, n_k , we have

$$\mu \{f \in \mathbf{A} : \langle f(n_j) \rangle \in E_j \ (1 \leq j \leq k)\} = \prod_{j=1}^k \lambda(E_j),$$