

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: UNIFORM DISTRIBUTION ON DIVISORS AND BEHREND SEQUENCES
Autor: Tenenbaum, Gérald
Kapitel: 6. Metric results
DOI: <https://doi.org/10.5169/seals-87875>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

Inserting this into (105), we arrive at

$$(\log x)^{-1} \sum_{n \leq x} \frac{\Delta(n; \theta \Omega)}{n 2^{\Omega(n)}} \ll \frac{1}{T} + \frac{\log q}{\log_2 x} \ll q_1(x; \theta)^{-1+o(1)}.$$

Since $q_1(x; \theta)$ is a non-decreasing function of x , this implies (107) and the proof is thereby completed.

Proof of Corollary 11. Put $\tau(n, \theta) := \sum_{d|n} d^{i\theta}$. When $f = \log$, we have that $g_v(n) = \tau(n, 2\pi v)$. By lemma 30.2 of [14] we infer that, uniformly for $1 \leq |\theta| \leq \exp \sqrt{\log x}$,

$$\sum_{p \leq x} \frac{|\tau(p, \theta)|}{p} = \sum_{p \leq x} \frac{|\cos(\frac{1}{2} \theta \log p)|}{p} = \frac{2}{\pi} \log x + O(1).$$

This is proved by partial summation from a strong form of the prime number theorem. Thus we obtain that we have uniformly for $1 \leq v \leq \log x$

$$C_v(x; \log) = (1 - 2/\pi) \log_2 x + O(1).$$

Inserting this into (105) with $T = \log x$ and choosing optimally $y = \pi/2$, we obtain

$$\Delta(n; f) < \xi(n) (\log_2 n) \left(\frac{4}{\pi}\right)^{\Omega(n)} \text{ ppl}$$

for all $\xi(n) \rightarrow \infty$. This implies the required result by a now standard device.

6. METRIC RESULTS

In this last section, we investigate the problem of uniform distribution on divisors from a further statistical point of view, regarding as random not only the integer n but also the function f . Thus, we define a measure μ on the set \mathbf{A} of all real valued arithmetical function as the inverse image of the Haar measure on the compact group $(\mathbf{R}/\mathbf{Z})^{\mathbf{N}}$ by the canonical mapping $f \mapsto \langle f \rangle$. In other words, μ is characterised by the property that for all finite families $\{E_j : 1 \leq j \leq k\}$ of measurable subsets of the torus \mathbf{R}/\mathbf{Z} and for all integers n_1, n_2, \dots, n_k , we have

$$\mu \{f \in \mathbf{A} : \langle f(n_j) \rangle \in E_j \ (1 \leq j \leq k)\} = \prod_{j=1}^k \lambda(E_j),$$

where λ stands for the Lebesgue measure on \mathbf{R}/\mathbf{Z} . The only basic property of μ that we shall use is the orthogonality relation

$$(113) \quad \int_{\mathbf{A}} e(\nu f(n) - \nu f(m)) d\mu(f) = \begin{cases} 1 & \text{if } m = n, \\ 0 & \text{if } m \neq n. \end{cases}$$

If an arithmetic function behaves statistically, then one expects that the Weyl sums

$$g_\nu(n) := \sum_{d|n} e(\nu f(d)),$$

and hence the discrepancy $\Delta(n; f)$, will normally have size roughly $\sqrt{\tau(n)}$. The purpose of the next theorem is to establish that this is indeed the case.

THEOREM 14. *Let $\xi(n) \rightarrow \infty$. For μ -almost all arithmetic functions f , we have*

$$(114) \quad \Delta(n; f) < \xi(n) (\log_2 n)^3 (\tau(n))^{1/2} \quad \text{ppl.}$$

Moreover, the exponent $\frac{1}{2}$ is sharp in this statement.

Proof. The upper bound follows from (24) with $y = 2$, $T = (\log x)^2$, namely

$$(115) \quad S(x; f) := \frac{1}{\log x} \sum_{n \leq x} \frac{\Delta(n; f)^2}{2^{\Omega(n)} n} \\ \ll \frac{1}{(\log x)^3} + \frac{\log_2 x}{\sqrt{\log x}} \sum_{1 \leq \nu \leq T} \frac{1}{\nu} H_\nu(x; f),$$

with

$$H_\nu(x; f) := \sum_{k=1}^{\infty} \left(\frac{1}{2} \right)^{\Omega(k)} \frac{1}{k^{1+\sigma}} \left| \sum_{m=1}^{\infty} \frac{e(\nu f(km))}{m^{1+\sigma} 2^{\Omega(m)}} \right|^2 \quad (\sigma := 1/\log x).$$

We have $H_\nu(x; f) \leq 2H_\nu^\dagger(x; f) + 2H_\nu^\ddagger(x; f)$ where the m -sum is restricted to $m \leq (\log x)^3$ in H_ν^\dagger and to $m > (\log x)^3$ in H_ν^\ddagger . We note right away the trivial estimate

$$H_\nu^\dagger(x; f) \ll \sqrt{\log x} \log_2 x,$$

which follows from (6) by partial summation. We deduce from this and (115) that

$$(116) \quad S(x; f) \ll (\log_2 x)^3 + R(x; f),$$

with

$$R(x; f) := \frac{\log_2 x}{\sqrt{\log x}} \sum_{1 \leq \nu \leq T} \frac{1}{\nu} H_\nu^\ddagger(x; f).$$

Expanding the square and integrating over f with respect to μ we get from (113) that

$$\int_{\mathbf{A}} H_v^\dagger(x; f) d\mu(f) = \sum_{k=1}^{\infty} \frac{1}{2^{\Omega(k)} k^{1+\sigma}} \sum_{m > (\log x)^3} \frac{1}{4^{\Omega(m)} m^{2+2\sigma}} \ll 1/(\log x)^2 .$$

Hence

$$\int_{\mathbf{A}} R(x; f) d\mu(f) \ll \frac{(\log_2 x)^2}{(\log x)^{5/2}}$$

Markov's inequality thus implies that

$$\mu\{f \in \mathbf{A} : R(2^l; f) \geq 1\} \ll 1/l^2 \quad (l = 1, 2, \dots) ,$$

so it follows by the Borel-Cantelli theorem, that for μ -almost all f we have

$$R(2^l; f) \ll 1 \quad (l = 1, 2, \dots) .$$

In view of (116), we see that the estimate $S(2^l; f) \ll (\log 2l)^3$ holds μ -almost surely in f and uniformly for $l \geq 1$. However, using the trivial bound $\Delta(n; f) \leq \tau(n)$, we readily see that

$$S(x; f) - S(2^l; f) \ll 1 \quad (2^l \leq x < 2^{l+1}) .$$

This yields that for μ -almost all f and uniformly in $x \geq 3$, we have

$$S(x; f) \ll (\log_2 x)^3 ,$$

which in turn implies (114).

To show that the exponent $\frac{1}{2}$ is sharp, we simply use (109) with $v = 1$ in the form

$$4\pi^2 \int_{\mathbf{A}} \Delta(n; f)^2 d\mu(f) \geq \int_{\mathbf{A}} |g_1(n)|^2 d\mu(f) = \tau(n) ,$$

where the equality follows from (113). This plainly implies that there is no $\alpha < \frac{1}{2}$ such that $\Delta(n; f) \ll \tau(n)^\alpha$ pp l for μ -almost all f : such a bound is actually false as soon as $\tau(n)$ is large enough.

The same quadratic mean approach that we used for Theorem 14 yields metric results for more restricted classes of arithmetic functions. We quote without proof the following theorem.

THEOREM 15. *The function $d \mapsto \theta^d$ is erd for almost all $\theta > 1$ and the function $d \mapsto \lambda \theta^d$ is erd for all $\theta > 1$ and almost all λ .*

More precisely, the corresponding discrepancies satisfy

$$(117) \quad \Delta(n; f) \ll \tau(n)^{1/2+o(1)} \quad \text{ppl},$$

under the indicated hypotheses, and the exponent $\frac{1}{2}$ is sharp.

Theorems 14 and 15 together provide an optimal strengthening of theorem 5 of Dupain, Hall & Tenenbaum [4].

REFERENCES

- [1] DAVENPORT, H. *Multiplicative number theory*, second edition. Springer, New York, Heidelberg, Berlin, 1980.
- [2] DAVENPORT, H. and P. ERDŐS. On sequences of positive integers. *Acta Arith.* 2 (1937), 147-151.
- [3] DAVENPORT, H. and P. ERDŐS. On sequences of positive integers. *J. Indian Math. Soc.* 15 (1951), 19-24.
- [4] DUPAIN, Y., R. R. HALL and G. TENENBAUM. Sur l'équirépartition modulo 1 de certaines fonctions de diviseurs. *J. London Math. Soc. (2)* 26 (1982), 397-411.
- [5] ERDŐS, P. Some unconventional problems in number theory. *Astérisque* 61 (1979), 73-82.
- [6] ERDŐS, P., R. R. HALL and G. TENENBAUM. On the densities of sets of multiples. *J. reine angew. Math.* 454 (1994), 119-141.
- [7] GRAHAM, S. W. and G. KOLESNIK. *Van der Corput's method of exponential sums*. London Math. Soc. Lecture Notes 126, Cambridge University Press, 1991.
- [8] HALL, R. R. Sums of imaginary powers of the divisors of integers. *J. London math. Soc. (2)* 9 (1975), 571-580.
- [9] ——— The distribution of $f(d) \pmod{1}$. *Acta Arith.* 31 (1976), 91-97.
- [10] ——— A new definition of the density of an integer sequence. *J. Austral. Math. Soc. Ser. A* 26 (1978), 487-500.
- [11] ——— The divisor density of integer sequences. *J. London Math. Soc. (2)* 24 (1981), 41-53.
- [12] ——— Sets of multiples and Behrend sequences. In: *A tribute to Paul Erdős* (editors A. Baker, B. Bollobás, A. Hajnal), Cambridge University Press, 1990, 249-258.
- [13] HALL, R. R. and G. TENENBAUM. Les ensembles de multiples et la densité divisorielle. *J. Number Theory* 22 (1986), 308-333.
- [14] HALL, R. R. and G. TENENBAUM. *Divisors*. Cambridge University Press, 1988.
- [15] HALL, R. R. and G. TENENBAUM. On Behrend sequences. *Math. Proc. Camb. Phil. Soc.* 112 (1992), 467-482.
- [16] KARATSUBA, A. A. Estimates for trigonometric sums by Vinogradov's method and some applications. *Proc. Steklov Inst. Math.* 112 (1971), 251-265.
- [17] KÁTAI, I. The distribution of divisors (mod 1). *Acta Math. Acad. Sci. Hungar.* 27 (1976), 149-152.
- [18] ——— Distribution mod 1 of additive functions on the set of divisors. *Acta Arith.* 30 (1976), 209-212.