

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON THE GAUSS-BONNET FORMULA FOR LOCALLY SYMMETRIC SPACES OF NONCOMPACT TYPE
Autor: Leuzinger, Enrico
Kapitel: 4. A NEW PROOF OF THE GAUSS-BONNET FORMULA
DOI: <https://doi.org/10.5169/seals-87876>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

$\pi_A : Y \rightarrow A$ we set $A_\Theta(s) := \pi_A \circ \mu^{-1}(\mathcal{H}_{1\Theta}(s)) \subset A$. The set $A_\Theta(s)$ is contained in an "affine" subspace of A of the form $a_1 a_*(s) A^{q-k}$ where $a_1 a_*(s) \in A$ and A^{q-k} is a $q-k$ -dimensional subgroup of A (see Sections 3 and 4 of [L2]). We denote the restriction of dv_A to A^{q-k} by $dv_{A^{q-k}}$; for $k = q$ we have $A^0 = e$ and we set $dv_{A^0} \equiv 1$. By Lemma 3.3 we have (for k equal to the number of elements of Θ)

$$\text{Vol}(V_E^{n-k}(s)) \prec \int_{\mu^{-1}(\mathcal{H}_{1\Theta}(s))} \rho(a)^{-1} dv_U \wedge dv_Z \wedge dv_{A^{q-k}}.$$

Since the horospherical piece $\mathcal{H}_{1\Theta}(s)$ is part of a Siegel set $\mathcal{S}_{\omega, \tau}$ with ω relatively compact (and hence of finite volume) in UM we get

$$\begin{aligned} \int_{\mu^{-1}(\mathcal{H}_{1\Theta}(s))} \rho(a)^{-1} dv_U \wedge dv_Z \wedge dv_{A^{q-k}} &\prec \\ &\prec \int_{\omega} dv_U \wedge dv_Z \int_{A_\Theta(s)} \rho(a)^{-1} dv_{A^{q-k}} \prec \int_{A_\Theta(s)} \rho(a)^{-1} dv_{A^{q-k}}. \end{aligned}$$

Also by definition of a Siegel set we have $\alpha(a) \geq \tau \succ 1$ for all $\alpha \in \Delta$. Moreover, the computations in the proof of Lemma 4.1 (and Lemma 3.5) in [L2] show that for all $\alpha \in \Theta$ one has $\alpha(a_1 a_*(s)) \succ e^{\mu_\alpha s}$ with $\mu_\alpha > 0$. Hence, as $\Theta \subset \Delta$ is not empty and since $\rho = \sum_{\alpha \in \Delta} c_\alpha \alpha$ ($c_\alpha > 0$), there is a uniform constant $c > 0$ such that $\rho(a)^{-1} \prec e^{-cs}$ for all $a \in A_\Theta(s)$. As noted above the set $A_\Theta(s)$ is contained in a $(q-k)$ -dimensional affine cone in A . It is similar (in the sense of Euclidean geometry) to $A_\Theta(0)$ with similarity factor s (see the proof of Lemma 4.1 in [L2]). Hence we eventually get $\int_{A_\Theta(s)} dv_{A^{q-k}} \prec s^{q-k}$ and the Lemma follows. \square

4. A NEW PROOF OF THE GAUSS-BONNET FORMULA

In this section we present a new simplified proof of the Gauss-Bonnet theorem for higher rank locally symmetric spaces.

THEOREM 4.1. *Let X be a Riemannian symmetric space of noncompact type and \mathbb{R} -rank ≥ 2 and let Γ be an irreducible, torsion-free (non-uniform) lattice in the group of isometries of X . Then for the locally symmetric space $V = \Gamma \backslash X$ the Gauss-Bonnet formula holds:*

$$\chi(V) = \int_V \Psi dv.$$

Proof. By Proposition 2.2 there is an exhaustion $V = \bigcup_{s \geq 0} V(s)$ of V by Riemannian polyhedra $V(s)$. Each polyhedron $V(s)$ in this exhaustion is equipped with the Riemannian metric induced by the one of V . Proposition 1.1 applied to $V(s)$ yields

$$\left| (-1)^n \chi'(V(s)) - \int_{V(s)} \Psi dv \right| \prec \sum_{k=1}^q \sum_E \int_{V_E^{n-k}(s)} \int_{O(p)} \|\Psi_{E,k}\| d\omega_{k-1} dv_E(p)$$

where $q = \dim A$ is the \mathbb{Q} -rank of \mathbf{G} (see Section 2.1) and where the index E runs through a finite set. As we remarked in Section 1 the function $\Psi_{E,k}$ is locally computable from the components of the metric and the curvature tensor of $V(s)$ and from the components of the second fundamental form of $V_E^{n-k}(s)$ in $V(s)$. The fact that V is locally symmetric together with Lemma 3.2 thus implies that $\|\Psi_{E,k}\| \prec 1$ for all E, k . Using Lemma 3.4 we conclude that

$$\left| (-1)^n \chi'(V(s)) - \int_{V(s)} \Psi dv \right| \prec \sum_{k,E} \text{Vol}(V_E^{n-k}(s)) \prec e^{-cs} \sum_{k=1}^q s^{q-k}.$$

By Proposition 2.3 we have $\chi'(V(s)) = \chi(V)$. The polyhedra $V(s)$ exhaust V and $\chi(V)$ is an integer; hence $(-1)^n \chi(V) = \int_{V(s)} \Psi dv$ for sufficiently large s . Finally, for n odd $\Psi \equiv 0$ by definition (see [AW]) and the claimed formula follows. \square

REFERENCES

- [AW] ALLENDOERFER, C. B. and A. WEIL. The Gauss-Bonnet theorem for Riemannian polyhedra. *Trans. Amer. Math. Soc.* 53 (1943), 101–129.
- [BGS] BALLMANN, W., M. GROMOV and V. SCHROEDER. *Manifolds of Nonpositive Curvature*. Boston, 1985.
- [B1] BOREL, A. *Introduction aux groupes arithmétiques*. Paris, 1969.
- [B2] ——— Stable and real cohomology of arithmetic groups. *Ann. scient. Éc. Norm. Sup.* 4^e série 7 (1974), 235–272.
- [B3] ——— *Linear Algebraic Groups*. Second edition, New York, 1991.
- [BS] BOREL, A. and J.-P. SERRE. Corners and arithmetic groups. *Comment. Math. Helv.* 48 (1973), 436–491.
- [BT] BOREL, A. and J. TITS. Groupes réductifs. *I.H.E.S. Publ. Math.* 27 (1965), 55–150.
- [C] CHERN, S. S. A simple intrinsic proof of the Gauss-Bonnet formula for closed Riemannian manifolds. *Ann. of Math.* 45 (1944), 747–752.
- [CG1] CHEEGER, J. and M. GROMOV. On the characteristic numbers of complete manifolds of bounded curvature and finite volume. *Differential Geometry and Complex Analysis*. H. E. Rauch Memorial Volume (I. Chavel, H. M. Farkas, eds.), Berlin, 1985, 115–154.