

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: INTRODUCTORY NOTES ON RICHARD THOMPSON'S GROUPS
Autor: Cannon, J. W. / Floyd, W. J. / Parry, W. R.
Kapitel: §3. Presentations for F
DOI: <https://doi.org/10.5169/seals-87877>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

§3. PRESENTATIONS FOR F

Two presentations for F will be given in this section.

Now two groups F_1 and F_2 will be defined by generators and relations. The generators $A, B, X_0, X_1, X_2, \dots$ will be referred to as *formal symbols*, as opposed to the functions defined above. Given elements x, y in a group, $[x, y] = xyx^{-1}y^{-1}$.

$$F_1 = \langle A, B : [AB^{-1}, A^{-1}BA], [AB^{-1}, A^{-2}BA^2] \rangle$$

$$F_2 = \langle X_0, X_1, X_2, \dots : X_k^{-1}X_nX_k = X_{n+1} \quad \text{for } k < n \rangle$$

THEOREM 3.1. *There exists a group isomorphism from F_1 to F_2 which maps A to X_0 and B to X_1 .*

Proof. There is a group homomorphism from the free group generated by the formal symbols A and B to F_2 such that A maps to X_0 and B maps to X_1 . This homomorphism is surjective because $X_n = X_0^{-(n-1)}X_1X_0^{n-1}$ for $n \geq 2$. To see that the defining relations of F_1 are in the kernel of this homomorphism, note that

$$X_1^{-1}X_2X_1 = X_0^{-1}X_2X_0 \quad \text{and} \quad X_1^{-1}X_3X_1 = X_0^{-1}X_3X_0,$$

hence

$$[X_0X_1^{-1}, X_2] = 1 \quad \text{and} \quad [X_0X_1^{-1}, X_3] = 1,$$

hence

$$[X_0X_1^{-1}, X_0^{-1}X_1X_0] = 1 \quad \text{and} \quad [X_0X_1^{-1}, X_0^{-2}X_1X_0^2] = 1.$$

Thus to complete the proof of Theorem 3.1 it suffices to prove that there exists a group homomorphism from F_2 to F_1 which maps X_0 to A and X_1 to B . To prove this it in turn suffices, after setting $Y_0 = A$ and $Y_n = A^{-(n-1)}BA^{n-1}$ for $n \geq 1$, to prove that

$$(3.2) \quad Y_k^{-1}Y_nY_k = Y_{n+1} \quad \text{for } k < n.$$

A closely related statement is that

$$(3.3) \quad [A^{-1}B, Y_m] = 1 \quad \text{for } m \geq 3.$$

Lines (3.2) and (3.3) will be proved in this paragraph. To see that line (3.3) is true for $m = 3$ note that

$$[AB^{-1}, A^{-1}BA] = 1 \Rightarrow A^{-1}[AB^{-1}, A^{-1}BA]A = 1 \Rightarrow [B^{-1}A, A^{-2}BA^2] = 1$$

$$\Rightarrow [A^{-1}B, A^{-2}BA^2] = 1 \Rightarrow [A^{-1}B, Y_3] = 1.$$

The same argument gives line (3.3) for $m = 4$. The following equations show that line (3.2) is true if line (3.3) is true for $m = n - k + 2$.

$$\begin{aligned} Y_n Y_k &= A^{-n+1} B A^{n-1} A^{-k+1} B A^{k-1} = A^{-k+2} A^{-(n-k+1)} B A^{n-k+1} A^{-1} B A^{k-1} \\ &= A^{-k+2} Y_{n-k+2} A^{-1} B A^{k-1} = A^{-k+2} A^{-1} B Y_{n-k+2} A^{k-1} \\ &= A^{-k+1} B A^{k-1} A^{-k+1} Y_{n-k+2} A^{k-1} = Y_k Y_{n+1} \end{aligned}$$

Thus line (3.2) is true for every positive integer n and $k = n - 1$. In particular, $Y_3^{-1} Y_4 Y_3 = Y_5$. Because line (3.3) is true for $m = 3$ and $m = 4$, it follows that line (3.3) is true for $m = 5$. An obvious induction argument now gives line (3.3) for every $m \geq 3$. This proves lines (3.2) and (3.3).

The proof of Theorem 3.1 is now complete. \square

THEOREM 3.4. *There exist group isomorphisms from F_1 and F_2 to F which map the formal symbols $A, B, X_0, X_1, X_2, \dots$ to the corresponding functions in F .*

Proof. Example 1.2 shows that the interior of the support of the function AB^{-1} in F is disjoint from the supports of the functions $A^{-1}BA$, $A^{-2}BA^2$ in F , and so the functions A, B in F satisfy the defining relations of F_1 . Thus there exists a group homomorphism from F_1 to F which maps the formal symbols A, B to the corresponding functions in F . Corollary 2.6 shows that this group homomorphism is surjective. Theorem 3.1 shows that this surjective group homomorphism induces a surjective group homomorphism from F_2 to F which maps the formal symbols X_0, X_1, X_2, \dots to the corresponding functions in F . To prove Theorem 3.4 it suffices to prove that this latter group homomorphism is injective.

It will be proved that this latter group homomorphism is injective in this paragraph. The defining relations of F_2 imply that

$$X_k^{-1} X_n = X_{n+1} X_k^{-1}, \quad X_n^{-1} X_k = X_k X_{n+1}^{-1}, \quad X_n X_k = X_k X_{n+1} \quad \text{for } k < n.$$

It follows that every nontrivial element x of F_2 can be expressed as a positive element times a negative element as in Corollary-Definition 2.7. If X_k occurs in both the positive and negative part of x but X_{k+1} occurs in neither, then because $X_k X_{n+1} X_k^{-1} = X_n$ for $n > k$, it is possible to simplify x by deleting one occurrence of X_k from both the positive and negative part of x and replacing every occurrence of X_{n+1} in x by X_n for $n > k$. Thus every nontrivial element of F_2 can be put in normal form as in Corollary-Definition 2.7. It follows from Corollary-Definition 2.7 that every nontrivial element of F_2 maps to a nontrivial element of F , as desired.

This proves Theorem 3.4. \square