

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 42 (1996)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: SQUARE-FREE TOWER OF HANOI SEQUENCES
Autor: HINZ, Andreas M.
Kapitel: 2. Smaller alphabets
DOI: <https://doi.org/10.5169/seals-87878>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. SMALLER ALPHABETS

The six-letter alphabet \mathcal{A} can still be reduced by building blocks of three moves. From Corollary (i) we learn that they must be of the form $(\tilde{\alpha}, \tilde{\gamma}, \tilde{\beta})$ with $\tilde{\xi} \in \{\xi, \bar{\xi}\}$. Only five of these actually do occur:

THEOREM 2. *Triples of elements of c form a square-free sequence h over the five-letter alphabet $\{A, B, \Gamma, \Delta, E\}$ with*

$$\begin{aligned} A &:= (\alpha, \bar{\gamma}, \beta), & B &:= (\alpha, \gamma, \bar{\beta}), \\ \Gamma &:= (\bar{\alpha}, \gamma, \beta), & \Delta &:= (\alpha, \gamma, \beta), & E &:= (\bar{\alpha}, \gamma, \bar{\beta}). \quad \square \end{aligned}$$

Proof. From Corollary (o) we know: if γ occurs in c with a bar, its neighbors must be in odd positions and consequently unbarred. All the other triples turn up, the sequence starting

$$h = (A, B, A, \Gamma, A, B, \Delta, E, A, B, A, \Gamma, A, E, \Delta, \Gamma, A, B, A, \Gamma, A, B, \Delta, E, \dots).$$

Clearly, h is square-free, since any square would lead to a square in c as well, contradicting Theorem 1. \square

REMARK. h (and consequently c) is *not* strongly square-free; can you spot an abelian square? (The existence of a strongly square-free string over a five-letter alphabet has been established by P.A.B. Pleasants [17, Theorem 2].) \square

Let me finally mention another instance of the TH to emerge as a microcosmos: it is known that the number of *states*, i.e. distributions of the discs among the three pegs, of the TH which can be reached from the initial state with all discs on peg 0, say, in and in no less than $\mu \in \mathbf{N}_0$ moves, is a power of 2, namely $2^{\beta(\mu)}$, where $\beta(\mu)$ is the number of non-zero bits of μ (see [13, Proposition 5]). ($2^{\beta(\mu)}$ also happens to be the number of odd entries in the μ th row of Pascal's arithmetical triangle, as was realized by J.W.L. Glaisher [10, second § 14]; cf. [14, formula (4)].) Denoting $\beta(\mu) \bmod 2$ by m_μ , we obtain the *Thue-Morse sequence*

$$m := (0, 1, 1, 0, 1, 0, 0, 1, 1, 0, 0, 1, 0, 1, 1, \dots),$$

which by the subsequent substitution of α for $(0, 1, 1)$, β for $(0, 1)$, and γ for (0) leads to the square-free sequence

$$t := (\alpha, \beta, \gamma, \alpha, \gamma, \beta, \alpha, \beta, \gamma, \beta, \alpha, \dots)$$

over the three-letter alphabet $\{\alpha, \beta, \gamma\}$. This is, of course, the smallest possible alphabet with an infinite square-free string (clearly, a square-free word over a two-letter alphabet will come to an end after three entries) with which the whole theory started in the work of Axel Thue [19, Satz 3], [20, Sätze 6, 7, 20].

Obviously, t (as in fact any word with more than 7 elements over a three-letter alphabet) is not strongly square-free. Maybe TH sequences hold a clue for a more direct approach to the question (cf. [6]), if there is an infinite strongly square-free string over a four-letter alphabet, which has been answered positively by V. Keränen [16] employing a computer-aided proof. (An abelian square of length $2 \cdot 6$ in h starts after position 6.)

ACKNOWLEDGEMENTS. I gratefully acknowledge the liberty to work I enjoyed during my visit as a Research Fellow at King's College London, where this note was finished, made possible by EPSRC grant N° GR/K10362. I am indebted to Professor E. B. Davies FRS for his support and hospitality.

Thanks go to the referee for some valuable comments and in particular for bringing reference [16] to my notice.

REFERENCES

- [1] AFRIAT, S. N. *The Ring of Linked Rings*. Duckworth (London), 1982.
- [2] ALLOUCHE, J.-P., D. ASTOORIAN, J. RANDALL and J. SHALLIT. Morphisms, Squarefree Strings, and the Tower of Hanoi Puzzle. *Amer. Math. Monthly* 101 (1994), 651–658.
- [3] ALLOUCHE, J.-P. et F. DRESS. Tours de Hanoi et automates. *RAIRO Inform. Théor. Appl.* 24 (1990), 1–15.
- [4] BERSTEL, J. Some recent results on squarefree words. In *STACS 84 (Lecture Notes in Comput. Sci. 166)*, G. Goos, J. Hartmanis (eds.), Springer (Berlin), 1984, 14–25.
- [5] BOYER, C. B. *A History of Mathematics*. Princeton University Press (Princeton), 1985.
- [6] BROWN, T. C. Is there a sequence on four symbols in which no two adjacent segments are permutations of one another? *Amer. Math. Monthly* 78 (1971), 886–888.
- [7] CLAUS, N. (= E. LUCAS). *La Tour d'Hanoi, Véritable casse-tête annamite*. P. Bousrez (Tours), 1883.
- [8] CLAUS, N. (= E. LUCAS). La Tour d'Hanoi, Jeu de calcul. *Science et Nature*, Vol. I, N° 8 (1884), 127–128.
- [9] CUMMINGS, L. J. Gray codes and strongly square-free strings. In *Sequences II, Methods in Communication, Security, and Computer Science*, R. Capocelli, A. De Santis, U. Vaccaro (eds.), Springer (New York), 1993, 439–446.