Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 43 (1997)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: POLYGON SPACES AND GRASSMANNIANS

Autor: Hausmann, Jean-Claude / Knutson, Allen

Kapitel: 6. Toric manifold structures on $mP_+^3(\alpha)$ for m = 4,5,6

DOI: https://doi.org/10.5169/seals-63276

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 02.12.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

$$M = \begin{pmatrix} a_2 & b_2 \\ a_3 & b_3 \\ a_4 & b_4 \end{pmatrix}$$

for the diagonal $\partial_{2,4} := \rho(2) + \rho(3) + \rho(4)$. The bending flows around two diagonals $\partial_{p,q}$ and $\partial_{p',q'}$ commute if and only if the pairs $\{p,q\}$ and $\{p',q'\}$ intersect or are unlinked in $\mathbf{R}/m\mathbf{Z}$.

6. Toric manifold structures on ${}^m\mathcal{P}^3_+(\alpha)$ for m=4.5.6

In this section, we study examples of $\mathcal{P}_+^3(\alpha) \subset {}^m\mathcal{P}^3$ such that the m-3 diagonal functions $d_2,\ldots,d_{m-2}:\mathcal{P}_+^3(\alpha)\longrightarrow \mathbf{R}$ never vanish. The whole space $\mathcal{P}_+^3(\alpha)$ consists of prodigal polygons and, by §5, the bending flows give an action of a big (i.e. half-dimensional) torus on $\mathcal{P}_+^3(\alpha)$. By Delzant's theorem (see [De], or [Gu, §1]), we can construct from the moment polytope Δ_α alone a toric manifold which is equivariantly symplectomorphic to the space $\mathcal{P}_+^3(\alpha)$. This can be achieved also by [DJ,§1.5], though only up to equivariant diffeomorphism. The latter also gives the real part, the planar polygon space $\mathcal{P}^2(\alpha)$, as a 2^{m-3} -sheeted branched cover of Δ_α . We sum up below some results of these constructions without writing all the details.

Without explicit mention of the contrary, α is supposed to be generic. Contrary to the previous sections, we do not require that the perimeter of our polygons is 2. It was necessary to fix the perimeter in order to define the map ℓ and the value 2 is the natural choice to deal with the map $\Phi: \mathbf{V}_2(\mathbf{C}^m) \longrightarrow {}^m \widetilde{\mathcal{P}}^k$. But ${}^m \mathcal{F}^k(\alpha)$ makes sense for any $\alpha \in \mathbf{R}^m_{\geq 0}$ and so do the various moduli spaces ${}^m \mathcal{P}^k(\alpha)$, etc. When $\sum \alpha_i = 2$, the polytope Δ_α is a slice through the Gel'fand-Cetlin moment polytope Γ_m of §5; for general α it is a homothetic copy of this section.

(6.1) m=4: The condition which guarantees that d_2 never vanishes is $\alpha_1 \neq \alpha_2$ or $\alpha_3 \neq \alpha_4$. The space of quadrilaterals ${}^4\mathcal{P}^3_+(\alpha)$ is then a compact toric manifold of dimension 2, therefore diffeomorphic to $\mathbb{C}P^1$. The moment map d_2 has image the interval $\Delta_{\alpha} := I_1 \cap I_2$ where

$$I_1 := [|\alpha_1 - \alpha_2|, \alpha_1 + \alpha_2]$$
 and $I_2 := [|\alpha_4 - \alpha_3|, \alpha_4 + \alpha_3]$.

The space ${}^4\mathcal{P}^2(\alpha)$ is $\mathbf{R}P^1$. The quadrilateral spaces ${}^4\mathcal{P}^2(\alpha)_+$ have long since been classified (see for instance [Ha]). One has

$${}^{4}\mathcal{P}^{2}(\alpha)_{+} = \begin{cases} S^{1} \sqcup S^{1} & \text{when } I_{1} \subset I_{2} \text{ or } I_{2} \subset I_{1} \\ S^{1} & \text{otherwise} \end{cases}$$

Observe also that α is generic if and only if the boundaries of the intervals I_1 and I_2 do not meet.

By the Duistermaat-Heckman Theorem [Gu, §2], the symplectic volume of ${}^4\mathcal{P}^3(\alpha)$ is equal to the length of Δ_{α} . We would then obtain the same length if we had used the other diagonal $|\rho(2) + \rho(3)|$. This produces a statement of elementary Euclidean geometry: the variation intervals of the two diagonals of a quadrilateral with given sides in \mathbf{R}^3 are the same length.

(6.2) m=5: Conditions for which both d_2 and d_3 never vanish are for instance $\alpha_1 \neq \alpha_2$ and $\alpha_4 \neq \alpha_5$. The space of pentagons ${}^5\mathcal{P}^3_+(\alpha)$ is then a toric manifold of dimension 4. The moment polytope $\Delta_{\alpha} \in \mathbf{R}^2$ for (d_2, d_3) is the intersection of the rectangle I_{α}

$$I_{\alpha} := [|\alpha_1 - \alpha_2|, \alpha_1 + \alpha_2] \times [|\alpha_5 - \alpha_4|, \alpha_5 + \alpha_4]$$

with the non-compact rectangular region

$$\Omega_{\alpha} := \{(x, y) \in (\mathbb{R}_{>0})^2 \mid x + y \ge \alpha_3 \text{ and } y \ge x - \alpha_3 \text{ and } y \le x + \alpha_3 \}.$$




FIGURE 2: The moment polytope Δ_{α}

(see Figure 2). One sees that Δ_{α} has at most 7 sides. The generic α are exactly those for which the boundary of Ω_{α} contains no corner of I_{α} and ${}^5\mathcal{P}^3_+(\alpha)$ is then obtained by symplectic blowings up from $\mathbb{C}P^2$ or $S^2 \times S^2$. The space of planar polygons ${}^5\mathcal{P}^2_+(\alpha)$ is a closed surface obtained by gluing 4 copies of Δ_{α} and its Euler characteristic is given by the formula

$$\chi(^5\mathcal{P}^2(\alpha)) = 4 - \# \text{ (sides of } \Delta_{\alpha})$$

(see [DJ], Example 1.20) and is orientable if and only if $I_{\alpha} \subset \omega_{\alpha}$. One has of course $\chi({}^5\mathcal{P}^2_+(\alpha)) = 2\chi({}^5\mathcal{P}^2_-(\alpha))$ and ${}^5\mathcal{P}^2_+(\alpha)$ is an orientable surface $({}^m\mathcal{P}^k_+(\alpha))$ is always orientable). The possible cases, depending on the number of sides of Δ_{α} , are summed up in the following table.

# of sides	$\mathcal{P}^3_+(\alpha)$	$\mathcal{P}^2(\alpha)$	$\mathcal{P}^2_+(\alpha)$	Ex. of α
3	$\mathbb{C}P^2$	$\mathbb{R}P^2$	S^2	(2,1,5,1,2)
-	a) $\mathbb{C}P^2 \# \overline{\mathbb{C}P^2}$	Klein bottle	T^2	(3,2,5,1,2)
4	or			
	b) $S^2 \times S^2$	T^2	$T^2 \sqcup T^2$	(3,1,3,1,3)
5	$(S^2 \times S^2) \# \overline{\mathbb{C}P^2}$	$T^2 \# \mathbf{R} P^2$	Σ_2	(2,1,3,1,2)
L				
6	$(S^2 \times S^2) \# 2\overline{\mathbf{C}P^2}$	$T^2 # 2\mathbf{R}P^2$	Σ_3	(2,1,1,1,2)




FIGURE 3: $\Delta_{(1,1,1,1,1)}$

(6.3) Some embeddings of the regular pentagon $\alpha=(1,1,1,1,1)$ are not prodigal. However none are lined and thus the moduli space $V_0:={}^5\mathcal{P}^3(\alpha)$ is diffeomorphic for small ε to V_ε where $V_\varepsilon:={}^5\mathcal{P}^3(\alpha_\varepsilon)$ and

 $\alpha_{\varepsilon} := (1 + \varepsilon, 1, 1, 1, 1 + \varepsilon)$. The moment polytope for α_{ε} has then 7 sides and thus $V_0 \simeq V_{\varepsilon}$ is diffeomorphic to $(S^2 \times S^2) \# 3\overline{\mathbb{C}P^2}$ (if k = 2, ${}^5\mathcal{P}^2(\alpha)_+ \simeq \Sigma_4$). The "limit moment polytope" $\Delta_{(1,1,1,1,1)}$ is shown in Figure 3.

The pre-image in V_{ε} of the segments $\{x = \varepsilon\} \cap \Delta'_{\alpha}$ and $\{y = \varepsilon\} \cap \Delta'_{\alpha}$ are 2-spheres of symplectic volume proportional to ε , by the Duistermaat-Heckman Theorem. Passing to the limit V_0 , these spheres become Lagrangian, and so cannot be complex. This shows that the action of the bending torus is not complex — these polygon spaces are only equivariantly symplectomorphic, not equivariantly isometric, to toric varieties.

(6.4) Any class $r \in {}^5\mathcal{P}^{k=2,3}(\alpha)$ has a unique representative in $\rho \in {}^5\widetilde{\mathcal{P}}^k(\alpha)$ with $\rho(5) = (-\alpha_5, 0, 0)$ and $\gamma(r) := \rho(1) + \rho(2)$ in the half-plane $\mathcal{H} = \{z = 0, y \geq 0\}$. This provides a map $\gamma : {}^5\mathcal{P}^3(\alpha) \longrightarrow \mathcal{H}$ whose image $\widetilde{\Delta}_{\alpha}$ is the intersection $R_1 \cap R_2 \cap \mathcal{H}$ where R_1 and R_2 are the rings

$$R_1 := \{ v \in \mathbf{R}^2 \mid |\alpha_1 - \alpha_2| \le |v| \le \alpha_1 + \alpha_2 \},$$

$$R_2 := \{ v \in \mathbf{R}^2 \mid |\alpha_4 - \alpha_3| \le |v| \le \alpha_4 + \alpha_3 \}.$$




FIGURE 4: $\widetilde{\Delta}_{\alpha}$

The idea of reconstructing ${}^5\mathcal{P}^2(\alpha)$ by gluing copies of $\widetilde{\Delta}_{\alpha}$ goes back to the early works of W. Thurston on planar linkages (see [TW, p.100]). The relationship with our theory is the following: the domain $\widetilde{\Delta}_{\alpha}$ is straightened up into a PL-polytope Δ_{α} in \mathbf{R}^2 by the map $v \mapsto (|v|, |v - (0, \alpha_5)|)$ and Δ_{α} is just the moment polytope for the bending Hamiltonians $\partial_1(\rho) = |\rho(1) + \rho(2)|$ and $\partial_2(\rho) = |\rho(3) + \rho(4)|$.

(6.5) m=6: The conditions $\alpha_1 \neq \alpha_2$ and $\alpha_5 \neq \alpha_6$ imply that d_2 and d_4 never vanish. However, one cannot guarantee generically $d_3 \neq 0$. But we can replace the $d=(d_1,d_2,d_3)$ by $\delta:=(\partial_1,\partial_2,\partial_3)$ where

$$\partial_1 := d_1 = |\rho(1) + \rho(2)|$$
, $\partial_2 := |\rho(3) + \rho(4)|$, $\partial_3 := d_3 = |\rho(5) + \rho(6)|$

and guarantee non-vanishing of the δ_i 's by the generic condition $\alpha_{2i-1} \neq \alpha_{2i}$. Observe that $\partial_i \circ \Phi : \mathbf{V}_2(\mathbf{C}^m) \longrightarrow \mathbf{R}$ (i = 1, 2, 3) are the functions on $\mathbf{V}_2(\mathbf{C}^m)$ given (on $(a, b) \in \mathbf{V}_2(\mathbf{C}^m)$) by the difference of the eigenvalues of the (2×2) -matrices $M_i^* M_i$, where

$$M_1 := \begin{pmatrix} a_1 & b_1 \\ a_2 & b_2 \end{pmatrix} \qquad M_2 := \begin{pmatrix} a_3 & b_3 \\ a_4 & b_4 \end{pmatrix} \qquad M_3 := \begin{pmatrix} a_5 & b_5 \\ a_6 & b_6 \end{pmatrix}.$$

The moment polytope in \mathbb{R}^3 is the intersection of the rectangular parallelepiped

$$I_{\alpha} := [|\alpha_1 - \alpha_2|, \alpha_1 + \alpha_2] \times [|\alpha_4 - \alpha_3|, \alpha_4 + \alpha_3] \times [|\alpha_6 - \alpha_5|, \alpha_6 + \alpha_5]$$
 with the region

$$\Omega := \{(x, y, z) \in \mathbf{R}^3 \mid 0 \le z \le x + y , 0 \le x \le y + z \text{ and } 0 \le y \le x + z\}.$$

The domain Ω can be described as the convex hull of the three half-lines

$$\{0 \le x = y \text{ and } z = 0\}$$
, $\{0 \le y = z \text{ and } x = 0\}$, $\{0 \le z = x \text{ and } y = 0\}$

or the cone $\mathbf{R}_+ \cdot \Xi_3$ on the hypersimplex Ξ_3 . The polytope Δ_{α} has then at most 9 facets. The length-system α is generic when the boundary of Ω does not contain corners of I_{α} . As 6 is even, the regular hexagon is not generic: ${}^6\mathcal{P}^1(1,\ldots,1)$ contains 10 elements.

(6.6) The bending flows ∂ occurring in (6.4) and 6 admit the following generalization. For m=2n-1 or 2n, we define the *even-step* map $e: {}^m\mathcal{F}^k \longrightarrow {}^n\mathcal{F}^k$ by $e(\rho)(i):=\rho(2i-1)+\rho(2i)$ taking $e(\rho)(n):=\rho(m)$ if m is odd. We also call e the induced maps ${}^m\widetilde{\mathcal{P}}^k \stackrel{e}{\longrightarrow} {}^n\widetilde{\mathcal{P}}^k, {}^m\mathcal{P}^k_+ \stackrel{e}{\longrightarrow} {}^n\mathcal{P}^k_+$ and ${}^m\mathcal{P}^k \stackrel{e}{\longrightarrow} {}^n\mathcal{P}^k$. We call $\rho \in {}^m\mathcal{F}^k$ even generic if $e(\rho)$ is a proper polygon. Above the space of proper polygons, the map e is a smooth locally trivial bundle whose fiber is a product of (k-1)-spheres. Define $\partial = (\partial_1, \ldots, \partial_n): {}^m\mathcal{F}^k \longrightarrow \mathbf{R}^n$ by $\partial := \ell \circ e$. The map ∂ gives the side lengths of the new polygon $e(\rho)$. It is always continuous and smooth when $e(\rho)$ is a proper polygon. As the map e is a submersion on even-generic polygons, the critical values of ∂ are the same as those of ℓ , the walls of 4.3. As for the map ℓ , the map ∂ can be defined on each ${}^m\mathcal{P}^k(\alpha)$. Call $\alpha \in \mathbf{R}^m$ even generic if ${}^m\mathcal{P}^k(\alpha)$ only consists of even-generic polygons. For instance, α is even-generic if $\alpha_{2i-1} \neq \alpha_{2i}$ for all i. When k=3, ∂ is a moment map for the corresponding bending action of T^n defined on even-generic polygons.

Restrict to ${}^m\mathcal{P}^3(\alpha)_+$ for an even-generic α . Define the right-angled polytope

$$I_{\alpha} := \prod_{i=1}^{n} [|\alpha_{2i} - \alpha_{2i-1}|, \alpha_{2i} + \alpha_{2i-1}]$$

and consider the convex polytope $\Delta_{\alpha} \subset \mathbf{R}^n$

$$\Delta_{\alpha} := \begin{cases} I_{\alpha} \cap (\mathbf{R}_{+} \cdot \Xi_{n}) & \text{when } m = 2n \\ I_{\alpha} \cap (\mathbf{R}_{+} \cdot \Xi_{n}) \cap \{x_{n} = |\rho(m)|\} & \text{when } m = 2n - 1 \end{cases}$$

PROPOSITION 6.7. 1) The image of $\partial: {}^m\mathcal{P}^k(\alpha)_+ \longrightarrow \mathbf{R}^n$ is the whole polytope Δ_{α} .

2) If $x \in \Delta_{\alpha}$ is a regular value of ∂ , the even-step map e induces, for m = 3, a symplectomorphism from the symplectic reduction $T^n \setminus \partial^{-1}(x)$ onto ${}^n\mathcal{P}^k_+(x)$. \square

7. REMARKS AND OPEN PROBLEMS

- (7.1) Is there an octonionic version of Section 3? Alternately, are there $U_1(\mathbf{H})$ bendings in dimension 5 (like the $U_1(\mathbf{C})$ bending flows in dimension 3 and $U_1(R)$ flippings in dimension 2)?
- (7.2) Observe that the inclusion ${}^m\mathcal{P}^k \subset {}^m\mathcal{P}^{k+1}$ becomes a bijection when $k \geq m-1$ (triangles are always planar, etc.). In what ways are these spaces ${}^m\mathcal{P}^{m-1}$ more natural than the unstable ones?
- (7.3) The m-polygons whose first diagonal is of a given length forms a sphere bundle over a space of (m-1)-polygons. (For k=3 this is just symplectic reduction by the first bending circle.) This gives an inductive way to construct the space of m-polygons by gluing together (sphere bundles over) the spaces of (m-1)-polygons; it would require identification of these sphere bundles, which in k=3 might be done using the Duistermaat-Heckman theorem (where the circle bundle is determined by its Euler class).

Alternately one might work out the fibers of the whole map d of section 5. Unfortunately in dimensions above 3 these are always singular (at, in particular, the planar polygons).

(7.4) In [KM1] and [Wa] there are presented "wall-crossing arguments" for identifying the spaces ${}^m\mathcal{P}^2(\alpha)$. It would be nice to relate these to a combination of [Du] and the paper [GS2], which presents its own wall-crossing arguments for any symplectic reduction by a torus.