

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN ALGORITHM FOR CELLULAR MAPS OF CLOSED SURFACES
Autor: DICKS, Warren / Glover, H. H.
Kurzfassung
DOI: <https://doi.org/10.5169/seals-63277>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

ETH-ZÜRICH

19. Jan. 1998

BIBLIOTHEK

AN ALGORITHM FOR CELLULAR MAPS OF CLOSED SURFACES

by Warren DICKS and H. H. GLOVER

ABSTRACT. The purpose of this article is to use diagrammatic methods to give proofs, accessible to algebraists, of some important topological results of H. Kneser, A. L. Edmonds, and R. Skora; we then describe some consequences for homomorphisms between surface groups. Cellular maps between two-dimensional CW-complexes can be represented by diagrams which, in turn, can be interpreted algebraically in terms of fundamental groupoids. For diagrams representing cellular maps between closed surfaces, we show how to apply certain homotopy equivalences algorithmically to obtain a normal-form map, which is a branched covering, or a pinching followed by a covering, or a map which collapses a graph of punctured spheres to a graph immersed in the one-skeleton of the target surface. We then indicate how the algorithm can be expressed entirely in terms of formal manipulations with presentations of surface groupoids, yielding algebraic proofs of results about homomorphisms between surface groups.

1. INTRODUCTION

We begin by recalling some basic concepts.

1.1. DEFINITIONS. Let β be a map between closed surfaces (without boundary).

Then β is a *branched covering* if deleting finitely many points from the source and from the target yields a covering.

We say that β is a (possibly trivial) *pinching* if it is obtained by collapsing, to a point, a compact subsurface with a single boundary component.

The (geometric) *degree* of β , denoted $\mathcal{G}(\beta)$, is the least non-negative integer d such that there is a map β' homotopic to β , such that the inverse image under β' of some 2-disk consists of d 2-discs, each mapped homeomorphically by β' to the chosen disk.