

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	43 (1997)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	QUATERNARY CUBIC FORMS AND PROJECTIVE ALGEBRAIC THREEFOLDS
Autor:	SCHMITT, Alexander
Kapitel:	2. The Invariant Theory of Quaternary Cubic Forms
DOI:	https://doi.org/10.5169/seals-63278

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

2. THE INVARIANT THEORY OF QUATERNARY CUBIC FORMS

2.1. *Stable, Semistable and Nullforms.* The stable and semistable quaternary cubic forms and the quaternary cubic nullforms were determined by Hilbert [Hi] (for the definition of semistable and stable see [Ne], *nullform* means non-semistable form):

THEOREM 3. i) A quaternary cubic form f is stable (resp. semistable) if and only if the surface $\{f = 0\}$ has at most singularities of type A_1 (resp. A_2).

ii) A quaternary cubic form f is a nullform if and only if the surface $\{f = 0\}$ has isolated singularities of type A_k ($k \geq 3$), D_4 , D_5 , E_6 , or \tilde{E}_6 , or if it has non-isolated singularities.

2.2. *Degenerations of Orbits of Semistable Forms.* First, one observes that the semistable forms with closed orbit are precisely the forms whose associated cubic surfaces have three A_2 -singularities. Applying Luna's slice theorem, one then computes the following table of degenerations where we characterize a form by the configuration of singularities on the corresponding cubic surface:

$$\begin{array}{ccc} A_2 & A_2A_1 & 2A_1A_2 \\ \searrow & \searrow & \downarrow \\ 2A_2 & 2A_2A_1 & \\ \searrow & \downarrow & \\ 3A_2 & & \end{array}$$

The details can be found in [Sch1], 58ff.

2.3. *The Ring of Invariants.* Proofs of the following results can be found in the paper [Be]. We want to describe the ring $A := \mathbf{C}[S^3(\mathbf{C}^{4^\vee})]^{\mathrm{SL}_4(\mathbf{C})}$. This is the coordinate ring of the categorical quotient $S^3(\mathbf{C}^{4^\vee}) // \mathrm{SL}_4(\mathbf{C})$. It is the ring of polynomial expressions in the coefficients of cubic polynomials which are constant on all $\mathrm{SL}_4(\mathbf{C})$ -orbits. In order to describe the ring A , we first introduce the following vector space

$$S := \left\{ r_1x_1^3 + r_2x_2^3 + r_3x_3^3 + r_4x_4^3 + r_5x_5^3 \mid \sum x_i = 0 \right\}.$$

On S , there is a natural action of the alternating group \mathfrak{A}_5 , and $A \subset \mathbf{C}[S]^{\mathfrak{A}_5}$. This inclusion is constructed as follows: The group of automorphisms H of the Sylvester pentrahedron naturally acts on S , and it can be shown that the natural

morphism $S//H \longrightarrow S^3(\mathbf{C}^4^\vee)//\mathrm{SL}_4(\mathbf{C})$ is birational. This induces the inclusion $A \subset \mathbf{C}[S]^H$. Now, H is a finite group of order 480 obviously containing \mathfrak{A}_5 . Denote by σ_i , $i = 1, 2, 3, 4, 5$, and v the i -th symmetric function and the Vandermonde determinant in the r_i . Then $\mathbf{C}[S]^{\mathfrak{A}_5} = \mathbf{C}[\sigma_1, \dots, \sigma_5, v]$.

THEOREM 4. *The ring of invariants A is the subring of $\mathbf{C}[S]^{\mathfrak{A}_5}$ generated by the following invariant polynomials*

$$\begin{aligned} I_8 &:= \sigma_4^2 - 4\sigma_3\sigma_5, & I_{16} &:= \sigma_5^3\sigma_1, & I_{24} &:= \sigma_5^4\sigma_4, \\ I_{32} &:= \sigma_5^6\sigma_2, & I_{40} &:= \sigma_5^8, & I_{100} &:= \sigma_5^{18}v, \end{aligned}$$

which satisfy a relation

$$I_{100}^2 = P(I_8, I_{16}, I_{24}, I_{32}, I_{40}).$$

2.4. The Discriminant. Using techniques from the paper [BC], one obtains the following

PROPOSITION 3. *The discriminant of quaternary cubic forms is given by the formula*

$$\Delta = (I_8^2 - 64I_{16})^2 - 2^{11}(I_8I_{24} + 8I_{32}).$$

2.5. Moduli Spaces of Cubic Surfaces. Define $\overline{\mathcal{M}}$ to be the hyper-surface $\{I_{100}^2 - P(I_8, I_{16}, I_{24}, I_{32}, I_{40}) = 0\}$ in the weighted projective space $\mathbf{P}(8, 16, 24, 32, 40) = \mathbf{P}(1, 2, 3, 4, 5)$. Then $\mathcal{M} := \overline{\mathcal{M}} \setminus \{\Delta = 0\}$ is a moduli space for non-singular cubic surfaces. On the other hand, every non-singular cubic surface can be obtained as the blow up of \mathbf{P}_2 in six points in general position. The sextuples of points in general position form an open subset $\mathcal{U} \subset S^6\mathbf{P}_2$ of the sixth symmetric power of \mathbf{P}_2 . Furthermore, there is an action of $\mathrm{PGL}_3(\mathbf{C})$ on \mathcal{U} , and the geometric quotient $\mathcal{N} := \mathcal{U} // \mathrm{PGL}_3(\mathbf{C})$ does exist [Is]. By [Is], §6, \mathcal{N} is a coarse moduli space for pairs (X, L) consisting of a cubic surface X and a globally generated line bundle L which defines a blow down $X \longrightarrow \mathbf{P}_2$. Forgetting the line bundle L provides us with a morphism $\mathcal{N} \longrightarrow \mathcal{M}$, so that there is a surjection $f: \mathcal{U} \longrightarrow \mathcal{M}$. Hence, we can view the invariants of quaternary cubic forms as regular functions on \mathcal{U} . This relates the geometry of the cubic surface to the set of six points. One obtains, e.g.,

PROPOSITION 4. *The set of sextuples in \mathcal{U} whose associated cubic surface is given by an equation which is not a (nondegenerate) Sylvestrian pentahedral form is the Zariski-closed subset $\{f^*I_{40} = 0\}$.*

Of course, a better understanding of the geometric meaning of the other invariants should allow to extend this result.

II. CUBIC FORMS OF PROJECTIVE THREEFOLDS

1. PRELIMINARIES

For the convenience of the reader, we have collected the crucial theorems which we will use in the construction of our examples.

1.1. The Lefschetz Theorem on Hyperplane Sections. We summarize Bertini's Theorem and Lefschetz' Theorem in :

THEOREM 5. *Let Y be a projective manifold, L a very ample line bundle on Y , and $X := Z(s)$ the zero-set of a general section $s \in H^0(X, L)$. Then X is a manifold (connected if $\dim Y \geq 2$), and the inclusion $\iota: X \hookrightarrow Y$ induces isomorphisms*

$$\begin{aligned}\iota^*: H^i(Y, \mathbf{Z}) &\longrightarrow H^i(X, \mathbf{Z}), & i &= 1, \dots, \dim Y - 2; \\ \iota_*: \pi_i(X) &\longrightarrow \pi_i(Y), & i &= 1, \dots, \dim Y - 2.\end{aligned}$$

Proof. [La], Th. 3.6.7 & Th. 8.1.1. \square

1.2. Formulas for Blow Ups. A very simple way to obtain a new manifold from a given one is the blow up in a point or along a smooth curve. The cup form behaves as follows (we will suppose for simplicity that $H^2(Y, \mathbf{Z})$ is without torsion) :

THEOREM 6. i) *Let $\sigma: X \longrightarrow Y$ be the blow up of Y in a point. Let $q(x_1, \dots, x_n)$ be the cubic polynomial which describes the cup form of Y w. r. t. the basis $(\kappa_1, \dots, \kappa_n)$ of $H^2(Y, \mathbf{Z})$. If $h_0 \in H^2(X, \mathbf{Z})$ is the cohomology class of the exceptional divisor, then $(h_0, \sigma^*\kappa_1, \dots, \sigma^*\kappa_n)$ is a basis of $H^2(Y, \mathbf{Z})$ w. r. t. which the cup form of X is given by*

$$x_0^3 + q(x_1, \dots, x_n).$$