

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: QUATERNARY CUBIC FORMS AND PROJECTIVE ALGEBRAIC THREEFOLDS
Autor: SCHMITT, Alexander
Kapitel: 4. Real Cubic Forms which are not Cup Forms of Projective Algebraic Manifolds
DOI: <https://doi.org/10.5169/seals-63278>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3.12. $(3A_2)$. Let $X = (\mathbf{P}_1 \times \mathbf{P}_1 \times \mathbf{P}_1)^\wedge(p)$ be the blow up of $\mathbf{P}_1 \times \mathbf{P}_1 \times \mathbf{P}_1$ in the point p . The cup form of X is

$$x_4^3 + 6x_1x_2x_3.$$

3.13. $(2A_1A_2)$. Consider the curve $C = Z(s) \subset \mathbf{P}_1 \times \mathbf{P}_1 \times \mathbf{P}_1$ where $s \in H^0(\mathcal{O}(1, 1, 0) \oplus \mathcal{O}(0, 0, 1))$ is a general section, and let X be the blow up of $\mathbf{P}_1 \times \mathbf{P}_1 \times \mathbf{P}_1$ along C . The cup form

$$6x_1x_2x_3 - 3x_1x_4^2 - 3x_2x_4^2 - 2x_4^3$$

of X defines a cubic surface with A_1 -singularities in $[1 : 0 : 0 : 0]$ and $[0 : 1 : 0 : 0]$ and an A_2 -singularity in $[0 : 0 : 1 : 0]$.

3.14. (D_4'') . Let $X := \widehat{\mathbf{P}_1 \times \mathbf{P}_2}(p_1, p_2)$ be the blow up of $\mathbf{P}_1 \times \mathbf{P}_2$ in the points p_1 and p_2 . Its cup form is described by the polynomial

$$3x_1x_2^2 + x_3^3 + x_4^3.$$

This polynomial is the equation of a cubic surface with a D_4 -singularity in $[1 : 0 : 0 : 0]$.

3.15. *A Non-Singular Quadric with a Transversal Plane.* Manifolds with such cup forms may be obtained as suitable \mathbf{P}_1 -bundles over surfaces. Indeed, let Y be a smooth surface with $b_2 = 3$. W. r. t. a suitable basis (h_1, h_2, h_3) of $H^2(Y, \mathbf{Z})$, its cup form is given by $x_1^2 + x_2^2 + x_3^2$. Now, let E be a vector bundle of rank 2 such that $c_1^2(E) - c_2(E) \neq 0$. Let $X := \mathbf{P}(E) \xrightarrow{\pi} Y$ and choose $(\pi^*h_1, \pi^*h_2, \pi^*h_3, c_1(\mathcal{O}_X(1)))$ as a basis of $H^2(X, \mathbf{Z})$. Then, by [OV], Prop. 15, the cup form of X is given by

$$(c_1^2(E) - c_2(E))x_4^3 + x_4(x_1^2 + x_2^2 + x_3^2).$$

3.16. *A Quadric Cone with a Transversal Plane.* Let Y be a simply connected surface with $b_2 = 3$ and torsion free homology. The cup form of Y is given by a quadratic polynomial $q(x_1, x_2, x_3)$ defining a smooth conic. Thus, the cup form of $Y \times \mathbf{P}_1$ is given by

$$x_4 q(x_1, x_2, x_3).$$

4. REAL CUBIC FORMS WHICH ARE NOT CUP FORMS OF PROJECTIVE ALGEBRAIC MANIFOLDS

In the paper [Sch2], the author investigated the restrictions on the real cubic forms of projective manifolds imposed by the so called Hodge-Riemann bilinear relations:

THEOREM 8. *Let X be a Kählerian threefold and $h \in H^2(X, \mathbf{R})$ be a Kähler class. Then the map*

$$\begin{aligned} \langle \cdot, \cdot \rangle: \quad H^2(X, \mathbf{R}) \times H^2(X, \mathbf{R}) &\longrightarrow \mathbf{R} \\ (a, b) &\longmapsto a \cup b \cup h \end{aligned}$$

is a non-degenerate, symmetric bilinear form of signature $(2h^{2,0} + 1, h^{1,1} - 1)$.

One can restate this theorem in such a form as to obtain – at least in theory – some explicit inequalities in the coefficients of cubic polynomials which are satisfied by the cup forms of Kählerian and hence projective algebraic threefolds. The main result of [Sch2] is

THEOREM 9. *For $n \geq 4$, the polynomial*

$$x_0 \left(\frac{4-n}{4} x_0^2 - 3x_1^2 - \dots - 3x_n^2 \right)$$

cannot occur as the (real) cup form of a projective algebraic threefold with $b_1 = 0$ and $b_3 = 0$.

As a corollary, one obtains the following generalization of a result of Campana and Peternell [CP]:

THEOREM 10. *For $n \geq 4$, twistor spaces over $\mathbb{H}_{i=1}^n \mathbf{P}_2$ are not homeomorphic to projective algebraic threefolds.*

REFERENCES

- [AGV] ARNOLD, V.I., S.M. GUSEIN-ZADE and A.N. VARCHENKO. *Singularities of Differentiable Maps, Vol. I.* Birkhäuser, 1985.
- [BC] BARDELLI, F. and A. DEL CENTINA. Nodal cubic surfaces and the rationality of the moduli space of curves of genus two. *Math. Ann.* 270 (1985), 599–602.
- [Be] BEKLEMISHEV, N.D. Invariants of cubic forms in four variables. *Vestnik Mosk. Univ. Mat.* 37 (1982), 42–9.
- [BW] BRUCE, J.W. and C.T.C. WALL. On the classification of cubic surfaces. *J. London Math. Soc. (2)* 19 (1979), 245–56.
- [CP] CAMPANA, F. and T. PETERNELL. Rigidity of Fano 3-folds. *Comm. Anal. Geom.* 2 (1994), 173–201.
- [GH] GRIFFITHS, Ph. and J. HARRIS. *Principles of Algebraic Geometry.* Wiley Interscience, 1978.