

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THEOREM OF INGHAM IMPLYING THAT DIRICHLET'S L-FUNCTIONS HAVE NO ZEROS WITH REAL PART ONE
Autor: Bateman, Paul T.
DOI: <https://doi.org/10.5169/seals-63280>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

A THEOREM OF INGHAM
IMPLYING THAT DIRICHLET'S L -FUNCTIONS
HAVE NO ZEROS WITH REAL PART ONE

by Paul T. BATEMAN

§ 1. INTRODUCTION

Using Landau's lemma on Dirichlet series with non-negative coefficients, A. E. Ingham in [1] proved the following theorem.

INGHAM'S THEOREM. *Let*

$$g(s) = g(s, \epsilon) = \prod_p \left(1 - \frac{\epsilon(p)}{p^s} \right)^{-1} = \sum_{n=1}^{\infty} \frac{\epsilon(n)}{n^s} \quad (\operatorname{Re} s > 1),$$

where the product is extended over all prime numbers p and ϵ is a bounded completely multiplicative arithmetic function. Suppose that g can be analytically continued into some domain containing the closed interval $[\frac{1}{2}, 1]$ of the real axis. Then

$$g(1) \neq 0.$$

We recall that an arithmetic function ϵ is said to be completely multiplicative if $\epsilon(mn) = \epsilon(m)\epsilon(n)$ for all positive integers m and n . Since a completely multiplicative arithmetic function ϵ is determined by the values $\epsilon(p)$ for primes p , it is immediate that ϵ is bounded if and only if $|\epsilon(p)| \leq 1$ for all primes p . Actually Ingham assumed that $|\epsilon(p)|$ is either 0 or 1 for any prime p , but his proof can easily be modified to require only that $|\epsilon(p)| \leq 1$. (Cf. [6]).

The most interesting application of Ingham's theorem is that obtained by taking $\epsilon(n) = \chi(n)n^{-i\alpha}$, where χ is a residue character modulo k and α is a real number which is different from zero if χ is the principal character. Then

$g(s) = L(s + i\alpha, \chi)$ and the theorem gives the assertion $L(1 + i\alpha, \chi) \neq 0$. The main interest in Ingham's theorem is its breadth of applicability. In contrast the familiar use of a trigonometric inequality does not cover the case in which $\alpha = 0$ and χ is a real non-principal character; that case is the very one to which Landau's lemma applies most easily (cf. §2 of [3]).

Ingham proved the theorem by establishing and using the identity

$$(*) \quad \zeta(s) g(s, \epsilon) g(s, \eta) g(s, \epsilon\eta) = g(2s, \epsilon\eta) \sum_{n=1}^{\infty} E(n) H(n) n^{-s},$$

where ζ denotes the Riemann zeta function, η is another completely multiplicative arithmetic function, $\epsilon\eta$ is the pointwise product of ϵ and η , and

$$E(n) = \sum_{d|n} \epsilon(d), \quad H(n) = \sum_{d|n} \eta(d).$$

This is a generalization of a result of Ramanujan for the case $\epsilon(n) = n^a$, $\eta(n) = n^b$, where a and b are fixed complex numbers.

While the identity (*) is of some interest in itself, we show that for the purpose of proving Ingham's theorem there is no reason to make the detour needed to prove (*). Of course we still require Landau's lemma, which we state in the following form.

LANDAU'S LEMMA. *Suppose β and γ are real numbers with $\beta < \gamma$. Suppose that $c_n \geq 0$ for $n = 1, 2, 3, \dots$ and that the series $\sum c_n n^{-s}$ converges for $\operatorname{Re} s > \gamma$. Put*

$$f(s) = \sum_{n=1}^{\infty} c_n n^{-s} \quad (\operatorname{Re} s > \gamma).$$

If f can be analytically continued into some domain containing the closed interval $[\beta, \gamma]$ of the real axis, then the series $\sum c_n n^{-\beta}$ converges.

For a proof of Landau's lemma see [5], [4], or §2 of [2].

The proof of Ingham's theorem which we give here uses an argument similar to that used in [4] and [5], except that the argument in those two papers was phrased in such a way as to require analytic continuation into a domain containing the interval $[0, 1]$ of the real axis instead of the interval $[\frac{1}{2}, 1]$. The interval $[\frac{1}{2}, 1]$ could not be replaced in the hypothesis of Ingham's theorem by a shorter interval, i.e., one of the form $[\theta, 1]$, where $\theta > \frac{1}{2}$; this is shown by the example in which ϵ is the Liouville function λ and $g(s) = \zeta(2s)/\zeta(s) = \sum \lambda(n) n^{-s}$.

§2. PROOF OF INGHAM'S THEOREM

Suppose that $g(1) = 0$. We show that this assumption leads to a contradiction. We consider the function

$$F(s) = \zeta(s)^2 g(s) g^*(s) \quad (\text{Re } s > 1),$$

where $g^*(s) = g(s, \bar{\epsilon})$ and $\bar{\epsilon}$ is the arithmetic function which is the complex conjugate of ϵ . Clearly $g^*(1) = \overline{g(1)} = 0$. By hypothesis g is regular along the stretch $[\frac{1}{2}, 1]$ of the real axis and so therefore is g^* , since $g^*(s) = \overline{g(\bar{s})}$. Hence F is regular on $[\frac{1}{2}, 1]$, since the double pole of ζ^2 at $s = 1$ is canceled by the zeros of g and g^* there.

Using the identity

$$(1 - z)^{-1} = \exp\left(\sum_{k=1}^{\infty} z^k/k\right) \quad (|z| < 1),$$

we obtain (for $\text{Re } s > 1$)

$$\begin{aligned} F(s) &= \prod_p \left(1 - \frac{1}{p^s}\right)^{-2} \left(1 - \frac{\epsilon(p)}{p^s}\right)^{-1} \left(1 - \frac{\bar{\epsilon}(p)}{p^s}\right)^{-1} \\ &= \prod_p \exp\left(\sum_{k=1}^{\infty} \frac{2 + \epsilon(p)^k + \bar{\epsilon}(p)^k}{kp^{ks}}\right) \\ &= \prod_p \left\{ 1 + \left(\sum_{k=1}^{\infty} \frac{2 + \epsilon(p)^k + \bar{\epsilon}(p)^k}{kp^{ks}}\right) + \frac{1}{2} \left(\sum_{k=1}^{\infty} \frac{2 + \epsilon(p)^k + \bar{\epsilon}(p)^k}{kp^{ks}}\right)^2 + \dots \right\}. \end{aligned}$$

Thus F has a Dirichlet series expansion

$$F(s) = \sum_{n=1}^{\infty} a(n) n^{-s} \quad (\text{Re } s > 1).$$

Furthermore, since

$$2 + \epsilon(p)^k + \bar{\epsilon}(p)^k = 2 + 2 \text{Re}\{\epsilon(p)^k\} \geq 0,$$

we have $a(n) \geq 0$ for all n .

At this point we deviate from the approach used in [4] and [5] by noting that $a(p^2) \geq 1$ for each prime p . For, since

$$F(s) = \prod_p \left(1 + \frac{2}{p^s} + \frac{3}{p^{2s}} + \dots\right) \left(1 + \frac{\epsilon(p)}{p^s} + \frac{\epsilon(p)^2}{p^{2s}} + \dots\right) \left(1 + \frac{\bar{\epsilon}(p)}{p^s} + \frac{\bar{\epsilon}(p)^2}{p^{2s}} + \dots\right),$$

we find that

$$\begin{aligned}
 a(p^2) &= 3 + 2\epsilon(p) + 2\bar{\epsilon}(p) + \epsilon(p)^2 + \epsilon(p)\bar{\epsilon}(p) + \bar{\epsilon}(p)^2 \\
 &= 2 - \epsilon(p)\bar{\epsilon}(p) + \{1 + \epsilon(p) + \bar{\epsilon}(p)\}^2 \\
 &\geq 2 - |\epsilon(p)|^2 \geq 1.
 \end{aligned}$$

Thus

$$\sum_{n=1}^{\infty} \frac{a(n)}{n^{1/2}} \geq \sum_p \frac{a(p^2)}{p} \geq \sum_p \frac{1}{p}.$$

In view of the divergence of $\sum p^{-1}$, it follows that $\sum a(n)n^{-1/2}$ diverges.

On the other hand, applying Landau's lemma with $c_n = a(n)$, $\beta = \frac{1}{2}$, $\gamma = 1$, we find that $\sum a(n)n^{-1/2}$ converges. This contradiction shows that the assumption $g(1) = 0$ is untenable and so the proof is complete.

REFERENCES

- [1] INGHAM, A.E. Note on Riemann's ζ -function and Dirichlet's L -functions. *J. London Math. Soc.* 5 (1930), 107–112.
- [2] LANDAU, Edmund. Über einen Satz von Tschebyschef. *Math. Ann.* 61 (1905), 527–550, or *Collected Works vol. 2* (Thales Verlag, 1986), 206–229.
- [3] ——— Über das Nichtverschwinden einer Dirichletschen Reihe. *Sitzungsberichte der Preussischen Akad. Wiss. Berlin* (1906), 314–320, or *Collected Works vol. 2* (Thales Verlag, 1986), 230–236.
- [4] NARASIMHAN, Raghavan. Une remarque sur $\zeta(1 + it)$. *L'Enseignement Math.* (2) 14 (1969), 189–191.
- [5] NEWMAN, D.J. A “natural” proof of the non-vanishing of L -series. *Contemp. Math.* 143 (1993), 495–498.
- [6] SHAPIRO, George. On the non-vanishing at $s = 1$ of certain Dirichlet series. *Amer. J. Math.* 71 (1949), 621–626.

(Reçu le 16 avril 1997)

Paul T. Bateman

Department of Mathematics
 University of Illinois
 Urbana, IL 61801
 U. S. A.
 e-mail: bateman@math.uiuc.edu