Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	43 (1997)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
Artikel:	ON CYCLOTOMIC POLYNOMIALS, POWER RESIDUES, AND RECIPROCITY LAWS
Autor:	Sharifi, Romyar T.
Kapitel:	3. Background
DOI:	https://doi.org/10.5169/seals-63283

### Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. <u>Mehr erfahren</u>

#### **Conditions d'utilisation**

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. <u>En savoir plus</u>

#### Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. <u>Find out more</u>

## Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

THEOREM 3. Let p be as in Theorem 2. Then if q > 3 and  $s \ge 2$ , or q = 3 and  $s \ge 3$ , we have that q is a  $q^s$  th power modulo p.

We will also derive an analogue of Theorem 1 for the always tricky case of q = 2. The proof is found in Section 7 and requires the use of several valuable properties of norm residue symbols.

THEOREM 4. Let n be a positive multiple of 4, and let s be the largest integer such that  $2^s$  divides n. Let  $p = \Phi_n(2x)$  if s > 2, and let  $p = \Phi_n(4x)$ if s = 2. If p is a prime number, then every integer dividing x is a  $2^s$  th power modulo p.

We shall use the following notation throughout the remainder of the paper. Lower case Roman letters will denote rational integers unless otherwise noted. In particular, we shall use *m* as a generic positive integer. Furthermore,  $\zeta_m$  will denote a primitive *m*th root of unity in an appropriate cyclotomic extension of the rationals **Q**, and for such a choice of  $\zeta_m$  we set  $\lambda_m = 1 - \zeta_m$ . For a Galois extension *K* of a field *F*, we will denote its Galois group by  $G_{K/F}$ and its norm by  $N_{K/F}$ . If the ground field *F* is **Q**, it shall be left out of the notation. For example, the Galois group of *K* over **Q** is denoted by  $G_K$ .

## 3. BACKGROUND

We now recall the formalism of the power residue and norm residue symbols and list the general reciprocity laws that relate them. This section is designed for those not yet familiar with this material and may be skipped by others. Most of the bibliographical references give a more thorough treatment of one or more aspects of the material we present below. This section requires only knowledge of algebraic concepts such as the integral closure and Galois theory, but it will help to have some knowledge of local and global fields.

By an *algebraic number field* F we mean a finite extension of the rationals. Its *ring of integers* A is the integral closure of  $\mathbb{Z}$  in F. The set of *fractional ideals* of F is the set of finitely generated non-zero A-submodules of F. Any fractional ideal can be uniquely factored into integral powers of a finite number of prime ideals, and hence the fractional ideals form a group by taking formal products of the prime ideals. A non-zero element  $\alpha$  of F will be treated as a fractional ideal by considering the fractional ideal  $\alpha A$  that it generates. The additive p-adic valuation  $v_p$  corresponding to the prime ideal p returns the power of this prime ideal occurring in a factorization of the fractional ideal. Two fractional ideals a and b of A are said to be *relatively prime* if for every prime ideal p such that  $v_p(a) \neq 0$  we have  $v_p(b) = 0$ . For  $\alpha_1, \dots, \alpha_r \in F^*$ , denote by  $I(\alpha_1, \dots, \alpha_r)$  the group of fractional ideals of A which are relatively prime to  $\alpha_1, \dots, \alpha_r$ .

Let F be an algebraic number field containing the set  $\mu_m$  of mth roots of unity. Then for  $\alpha \in F^*$  and  $b \in I(m, \alpha)$ , the mth power residue symbol takes on a value in  $\mu_m$  and is denoted

$$(\alpha/\mathfrak{b})_{m,F}$$
 or  $\left(\frac{\alpha}{\mathfrak{b}}\right)_{m,F}$ .

When usage is clear, we will leave F out of the notation.

For an ideal  $\mathfrak{b}$  of A, we let  $N\mathfrak{b} = [A : \mathfrak{b}]$ . As a result,  $N\mathfrak{b} = [\mathbf{Z} : N_K\mathfrak{b}]$ and  $N_K\mathfrak{b} = (N\mathfrak{b})$ . For a prime ideal  $\mathfrak{p}$  and  $\alpha \in F^*$  relatively prime to  $\mathfrak{p}$  we have the following formula:

(3) 
$$\qquad \qquad \left(\frac{\alpha}{\mathfrak{p}}\right)_m \equiv \alpha^{(N\mathfrak{p}-1)/m} \mod \mathfrak{p}.$$

We also have:

THEOREM 5. The power residue symbol  $(\cdot / \cdot)_m$  has the following properties:

- (a)  $(\alpha \alpha'/\mathfrak{b})_m = (\alpha/\mathfrak{b})_m (\alpha'/\mathfrak{b})_m$  for  $\alpha, \alpha' \in F^*$  and  $\mathfrak{b} \in I(m, \alpha, \alpha')$ ;
- (b)  $(\alpha/\mathfrak{b}\mathfrak{b}')_m = (\alpha/\mathfrak{b})_m (\alpha/\mathfrak{b}')_m$  for  $\alpha \in F^*$  and  $\mathfrak{b}, \mathfrak{b}' \in I(m, \alpha)$ ;
- (c)  $(\alpha/\mathfrak{b})_m = (\alpha'/\mathfrak{b})_m$  if  $\alpha \equiv \alpha' \mod \mathfrak{b}$  for  $\alpha, \alpha' \in F^*$  and  $\mathfrak{b} \in I(m, \alpha)$ ,  $\mathfrak{b}$  an ideal;
- (d)  $\sigma(\alpha/\mathfrak{b})_m = (\sigma\alpha/\sigma\mathfrak{b})_m$  for  $\alpha \in F^*$ ,  $\mathfrak{b} \in I(m, \alpha)$  and  $\sigma$  an automorphism of F.

Finally, when  $\alpha$  is an *m*th root of unity, the power residue symbol can be evaluated by using (3) and Theorem 5(b). For  $\xi \in \mu_m$  and  $\mathfrak{b} \in I(m)$ , one sees that

(4) 
$$\left(\frac{\xi}{\mathfrak{b}}\right)_m = \xi^{(N\mathfrak{b}-1)/m}$$

By a *local field* we mean  $\mathbf{R}$ ,  $\mathbf{C}$  or a finite extension of  $\mathbf{Q}_p$ , the *p*-adic numbers, for some prime number *p*. In the latter case, the field is called *non-archimedean* and in the others, *archimedean*. Archimedean local fields

arise through completion of an algebraic number field with respect to an embedding of it in **R** or **C**. A non-archimedean local field over  $\mathbf{Q}_p$  arises through completion of an algebraic number field F with respect to a metric determined by the additive valuation associated to a prime ideal  $\mathfrak{p}$  of A which lies over the prime ideal p of  $\mathbf{Z}$ .

Both the absolute values defined by archimedean embeddings and the prime ideals of the ring of integers of the number field F are referred to as *primes* of F. Let  $\mathfrak{p}$  be a prime of F, and take the completion of F as described above. We say that the resulting local field  $F_{\mathfrak{p}}$  is the *completion of* F *at the prime*  $\mathfrak{p}$ . It is a topological field under the topology induced by the completion. In the non-archimedean case, the subring of  $F_{\mathfrak{p}}$  which is the completion of the ring of integers of F is a local ring called the *valuation ring*  $\mathcal{O}_{F_{\mathfrak{p}}}$  of  $F_{\mathfrak{p}}$ .

Let K be a local field which contains the *m*th roots of unity. Then

$$(\cdot, \cdot)_{m,K} \colon K^* \times K^* \to \mu_m$$

will denote the *m*th norm residue symbol of a field K with multiplicative group  $K^*$ . We use the definition of the norm residue symbol coinciding with that of [CF], [H], and [Se], which is the inverse of the symbol defined in [AT], [FV], [Iy], and [Ne]. As with the power residue symbol, K will usually be left out of the notation.

The norm residue symbol has many important and useful properties. We list several of them in the following theorem.

THEOREM 6. The norm residue symbol  $(\cdot, \cdot)_{m,K}$  has the following properties:

- (a)  $(\cdot, \cdot)_m$  is bimultiplicative;
- (b)  $(\alpha, \beta)_m = (\beta, \alpha)_m^{-1}$  for  $\alpha, \beta \in K^*$ ;
- (c)  $(\alpha, \beta)_m = 1$  for  $\alpha, \beta \in K^*$  if and only if  $\beta$  is the norm of an element of  $K(\sqrt[m]{\alpha})$ ;
- (d)  $(\alpha, \beta)_{m,L} = (N_{L/K}(\alpha), \beta)_{m,K}$  for  $\alpha \in L^*, \beta \in K^*$ , where L is a finite separable extension of K;

(e) 
$$(\alpha, \beta)_{mn}^n = (\alpha, \beta)_m$$
 for  $\alpha, \beta \in K^*$  if  $\mu_{mn} \subseteq K$ ;

(f) 
$$(\alpha, 1-\alpha)_m = 1$$
 for  $\alpha \in K^*, \alpha \neq 1$ ,

- (g)  $(\alpha, -\alpha)_m = 1$  for  $\alpha \in K^*$ ;
- (h)  $\sigma(\alpha, \beta)_m = (\sigma\alpha, \sigma\beta)_m$  for  $\alpha, \beta \in K^*$  and  $\sigma$  a continuous automorphism of K.

Now assume that *K* is non-archimedean. For  $\beta \in K^*$ , the *conductor* of the norm residue symbol  $(\cdot, \beta)_{m,K}$  is an ideal  $\mathfrak{f} = \mathfrak{f}(\beta)$  of the valuation ring  $\mathcal{O}_K$  and hence a power of the unique maximal ideal of this ring. The conductor is the largest ideal having the property that if  $\alpha \in \mathcal{O}_K^*$  is such that  $\alpha \equiv 1 \mod \mathfrak{f}$ , then  $(\alpha, \beta)_K = 1$ .

Again let F be an algebraic number field containing the *m*th roots of unity. Let  $\infty$  denote the formal product of the real primes (embeddings) of the field F, and let  $F_p$  denote the completion of F at a prime p. Then we have the following *law of reciprocity* for  $\alpha$ ,  $\beta \in F^*$  relatively prime to each other and to m:

(5) 
$$\left(\frac{\alpha}{\beta}\right)_{m,F} \left(\frac{\beta}{\alpha}\right)_{m,F}^{-1} = \prod_{\mathfrak{p}\mid m\infty} (\beta, \alpha)_{m,F\mathfrak{p}} ,$$

where  $\mathfrak{p} \mid m\infty$  indicates that  $\mathfrak{p}$  appears in the decomposition of  $m\infty$  into a product of primes. (That is, the product is taken over all prime ideals dividing m and all real primes.) Furthermore, if  $\gamma \in F^*$  is such that  $\mathfrak{p}$  divides m for all prime ideals  $\mathfrak{p}$  satisfying  $\upsilon_{\mathfrak{p}}(\gamma) \neq 0$  and  $\beta \in F^*$  is again relatively prime to m, we have

(6) 
$$\left(\frac{\gamma}{\beta}\right)_{m,F} = \prod_{\mathfrak{p}\mid m\infty} (\beta,\gamma)_{m,F_{\mathfrak{p}}}.$$

# 4. The odd case

For an odd prime q and a positive integer s, we now set  $l = q^s$ . If  $\alpha \in \mathbf{Q}_q^*$ , then  $\alpha$  may be written uniquely as  $\alpha = \xi q^b (1-q)^c$  where  $\xi \in \mu_{q-1}$ ,  $b \in \mathbf{Z}$ , and  $c \in \mathbf{Z}_q$ . Note that  $b = \upsilon_q(\alpha)$ , where  $\upsilon_q$  is the q-adic valuation. Denote by  $f_l(\alpha)$  the conductor of the norm residue character  $(\cdot, \alpha)_l$  for the *l*th cyclotomic field  $\mathbf{Q}_q(\zeta_l)$  over the q-adic rationals  $\mathbf{Q}_q$ . Robert Coleman and William McCallum have computed these conductors for all  $\alpha \in \mathbf{Q}_q^*$  in [CM]. We state the result here, though we shall use only its corollary. Recall that  $\lambda_m = 1 - \zeta_m$  for all positive integers m.