

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ON CYCLOTOMIC POLYNOMIALS, POWER RESIDUES, AND RECIPROCITY LAWS
Autor: Sharifi, Romyar T.
Kapitel: 8. The second proof
DOI: <https://doi.org/10.5169/seals-63283>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 18.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

We have shown that $(\pi, a)_l = 1$, and we get the desired result if $(a/\pi)_l = 1$. This is easily seen. Let $a = a'2^k$ where a' is odd. Then

$$\left(\frac{a}{\pi}\right)_l = \left(\frac{a'}{\pi}\right)_l \left(\frac{2}{\pi}\right)_l^k = \left(\frac{\pi}{a'}\right)_l (\pi, a')_l (\pi, 2)_l^k = \left(\frac{1}{a'}\right)_l (\pi, a)_l = 1,$$

where we have used the reciprocity laws (5) and (6) in the second equality and Theorem 5(c) in the third. \square

8. THE SECOND PROOF

This second proof is in many ways preferable to the first. It is much less dependent upon machinery (i.e., knowledge of the conductors), and it is specific to the case of cyclotomic polynomials.

Second proof of Theorem 1. We keep the notation of the first proof. The beginning of the proof runs along the lines of the first. Via the reciprocity laws, we therefore conclude that

$$\left(\frac{a}{\pi}\right)_l = (\pi, a)_l.$$

As in the proof of Theorem 4, it suffices to show that $(1 - qx\zeta_l, a)_l = 1$ for any multiple x of a and a primitive l th root of unity ζ_l .

By Theorem 6(f), we have

$$(1 - qx\zeta_l, a)_l = (1 - qx\zeta_l, qxa^{-1}\zeta_l)_l^{-1} = (1 - a\alpha, \alpha)_l,$$

where we have set $\alpha = qxa^{-1}\zeta_l$.

Now note that if we are given a power series $f_i \in \mathbf{Z}_q[[X]]$ with $f_i(0) = \gamma_i$ and a symbol $(1 - \alpha^i f_i(\alpha), \alpha)_l$, we can use multiplicativity on the left to manipulate the symbol into

$$\left(\frac{1 - \alpha^i f_i(\alpha)}{(1 - \alpha^i)^{\gamma_i}}, \alpha\right)_l (1 - \alpha^i, \alpha)_l^{\gamma_i} = (1 - \alpha^{i+1} f_{i+1}(\alpha), \alpha)_l (1 - \alpha^i, \alpha)_l^{\gamma_i},$$

where f_{i+1} is another power series over \mathbf{Z}_q . Since α has positive valuation, large enough powers of it will be congruent to 0 modulo the conductor of α . Therefore the symbol $(1 - \alpha^i f_i(\alpha), \alpha)_l$ will be 1 for large i . Taking $f_1 = a$, we see recursively that $(1 - a\alpha, \alpha)_l$ can be expressed as a finite product of powers of symbols of the form $(1 - \alpha^i, \alpha)_l$ with $i \geq 1$.

Let us fix an i and set $i = i'q^r$ with i' not divisible by q . Then i' is invertible mod l , and so by multiplicativity of the norm residue symbol we have

$$(1 - \alpha^i, \alpha)_l = (1 - (\alpha^{i'})^{q^r}, \alpha^{i'})_l^{i'-1}.$$

Now note that $\beta = \alpha^{i'}$ has the same form as α . That is, β is an integer multiple of q times a primitive n th root of unity. It will therefore suffice to show that $(1 - \alpha^{q^r}, \alpha)_l = 1$ for all $r \geq 0$. If $r = 0$, then Theorem 6(f) tells us already that this symbol is 1.

Now assume $1 \leq r < s$ (so $s \geq 2$). Note that

$$1 - (qx\zeta_l)^{q^r} = \prod_{j=1}^{q^r} (1 - qx\zeta_l \zeta_{q^r}^j).$$

So we need only show that $(1 - qx\zeta_l \xi, qx\zeta_l)_l = 1$ for every q^{s-1} th root of unity ξ . In this case,

$$(1 - qx\zeta_l \xi, qx\zeta_l)_l = (1 - qx\zeta_l \xi, \xi)_l^{-1}$$

by Theorem 6(f). As in (9), we can apply reciprocity law (5) and equation (4) to obtain

$$(10) \quad (1 - qx\zeta_l \xi, \xi)_l = \xi^{(N_K(1 - qx\zeta_l) - 1)/l}.$$

Here we have used the fact that ζ_l is a Galois conjugate of $\zeta_l \xi$. Note that

$$N_K(1 - qx\zeta_l) = \Phi_l(qx) \equiv 1 \pmod{q^{q^{s-1}}}.$$

As $q^{s-1} \geq 2s - 1$ for $s \geq 2$ and $q \geq 3$, we conclude that the symbol in (10) is 1.

Finally, assume that $r \geq s$. We then have

$$(1 - (qx\zeta_l)^{q^r}, qx\zeta_l)_l = (1 - (qx)^{q^r}, qx)_l (1 - (qx)^{q^r}, \zeta_l)_l.$$

As both entries are rational, we have that $(1 - (qx)^{q^r}, qx)_l$ is an l th root of unity which, by Theorem 6(h), is invariant under the action of $G_{\mathbf{Q}_q(\zeta_l)/\mathbf{Q}_q}$ and so must be 1. Furthermore, $(1 - (qx)^{q^r}, \zeta_l)_l$ can be evaluated as in (10). Since $[K : \mathbf{Q}] = q^{s-1}(q - 1)$, we have

$$N_K(1 - (qx)^{q^r}) = (1 - (qx)^{q^r})^{q^{s-1}(q-1)} \equiv 1 \pmod{q^{q^r+s-1}}.$$

Now we need only note that $q^r + s - 1 \geq 2s$ for all $r \geq s$ to finish the proof. \square

This method is easily used to deal with the case of $q = 2$, as most of the proof carries over. We leave the proof to the reader. Extending this method, the author has been able to compute the conductors which were used in the first proof of the theorems (for all q) [Sh2].

ACKNOWLEDGMENTS. Hendrik Lenstra was of great help throughout the preparation of this paper. Robby Robson, along with Tom Schmidt, advised me at the 1993 NSF Research Experiences for Undergraduates program at Oregon State. Raghavan Narasimhan made many helpful comments. I thank them, and all those who offered me guidance, wholeheartedly.

REFERENCES

- [AT] ARTIN, E. and J. TATE. *Class Field Theory*. Harvard, 1961.
- [CF] CASSELS, J. W. S. and A. FRÖHLICH, eds. *Algebraic Number Theory*. Academic Press, New York, 1967.
- [CM] COLEMAN, R. and W. MCCALLUM. Stable reduction of Fermat curves and Jacobi sum Hecke characters. *J. Reine Angew. Math.* 385 (1988), 41–101.
- [C] COX, D. *Primes of the Form $x^2 + ny^2$* . John Wiley & Sons, New York, 1989.
- [FV] FESENKO, I. and S. VOSTOKOV. *Local Fields and Their Extensions: A Constructive Approach*. American Mathematical Society, Providence, 1993.
- [H] HASSE, H. *Bericht über neuere Untersuchungen und Probleme aus der Theorie der algebraischen Zahlkörper, Teil II: Reziprozitätsgesetz*. Physica-Verlag, Würzburg, Germany, 1965.
- [IR] IRELAND, K. and M. ROSEN. *A Classical Introduction to Modern Number Theory, 2nd. ed.* Springer-Verlag, New York, 1990.
- [Iw] IWASAWA, K. *Local Class Field Theory*. Oxford University Press, New York, 1986.
- [Iy] IYANAGA, S. *The Theory of Numbers*. American Elsevier Publishing, New York, 1975.
- [La] LANG, S. *Algebraic Number Theory*. Addison-Wesley, Reading, Mass., 1970.
- [N] NEUKIRCH, J. *Class Field Theory*. Springer-Verlag, New York, 1986.
- [P] PRAPAVESSI, D. On the conductor of 2-adic Hilbert norm residue symbols. *J. Algebra* 149 (1992), 85–101.
- [Se] SERRE, J.-P. *Local Fields*. Springer-Verlag, New York, 1979.