

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AMENABILITY AND GROWTH OF ONE-RELATOR GROUPS
Autor: Ceccherini-Silberstein, Tullio G. / GRIGORCHUK, Rostislav I.
Kapitel: 3. Uniformly exponential growth and growth of graded algebras
DOI: <https://doi.org/10.5169/seals-63284>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3. UNIFORMLY EXPONENTIAL GROWTH AND GROWTH OF GRADED ALGEBRAS

In this section we describe a method of estimating growth functions of a group in terms of its graded Lie, and associative algebras defined via dimension subgroups. We begin by recalling some concepts and notations.

As in [Gri] considerations were given with respect to a Galois field \mathbf{GF}_p , here we modify the arguments for a field of characteristic 0, namely \mathbf{Q} .

Let G be a group; denote by $\mathbf{Q}[G]$ the group algebra of G over \mathbf{Q} , and by $\Delta \subset \mathbf{Q}[G]$ the augmentation ideal, that is the ideal generated by the elements of the form $g - 1$, with $g \in G$. Recall that the *lower central series* of G is the sequence of subgroups $\{\gamma_n(G)\}_{n=1}^{\infty}$ of G defined by $\gamma_1(G) = G$ and, for $n \geq 2$, $\gamma_n(G) = [G, \gamma_{n-1}(G)]$.

The subgroup

$$G_n = \{g \in G : g - 1 \in \Delta^n\}$$

is called the *n-th dimension subgroup* of G over \mathbf{Q} and it has the following characterisation due to Jennings [J] (see also [P: IV, Thm. 1.5] or [Pm: 11, Thm. 1.10])

$$G_n = \sqrt{\gamma_n(G)} := \{g \in G : \exists k \in \mathbf{N}, g^k \in \gamma_n(G)\}.$$

For any group G one defines as usual an associative graded algebra $\mathcal{A}(G)$ and two graded Lie algebras $L(G)$ and $\mathcal{L}(G)$ by

$$\begin{aligned} \mathcal{A}(G) &= \bigoplus_{n=1}^{\infty} \Delta^n / \Delta^{n+1} \\ L(G) &= \bigoplus_{n=1}^{\infty} [(G_n / G_{n+1}) \otimes_{\mathbf{Z}} \mathbf{Q}] \\ \mathcal{L}(G) &= \bigoplus_{n=1}^{\infty} [(\gamma_n(G) / \gamma_{n+1}(G)) \otimes_{\mathbf{Z}} \mathbf{Q}] \end{aligned}$$

(see for instance [P], [Pm]). Quillen's Theorem [Q] states that $\mathcal{A}(G)$ is the universal enveloping algebra of $L(G)$.

Assume now that G is finitely generated and set

$$\begin{aligned} a_n(G) &= \dim(\Delta^n / \Delta^{n+1}) \\ b_n(G) &= \text{rank}(G_n / G_{n+1}) \\ c_n(G) &= \text{rank}(\gamma_n(G) / \gamma_{n+1}(G)) \end{aligned}$$

where, by rank, we mean the torsion free rank of the corresponding abelian group. Then the following relations hold

$$\sum_{n=0}^{\infty} a_n(G)z^n = \prod_{n=1}^{\infty} (1 - z^n)^{-b_n(G)} = \prod_{n=1}^{\infty} (1 - z^n)^{-c_n(G)}.$$

The first equality follows easily from Quillen's Theorem [Pm: Thm. 4.10, Chapter 3] and the second one follows from the equality $b_n(G) = c_n(G)$ as proved in [Be].

In [Be] it is also proved that

$$\limsup_{n \rightarrow \infty} \sqrt[n]{a_n} = \limsup_{n \rightarrow \infty} \sqrt[n]{c_n}.$$

3.1. LEMMA. *For any finite system of generators A of a group G the following inequality holds:*

$$a_n(G) \leq \gamma_A^G(n), \quad n \geq 1.$$

Proof. For $x, y \in G$ we have

$$\begin{aligned} xy - 1 &= (x - 1) + (y - 1) + (x - 1)(y - 1) \\ x^{-1} - 1 &= -(x - 1) - (x - 1)(x^{-1} - 1) \end{aligned}$$

so that

$$\begin{aligned} xy - 1 &\equiv (x - 1) + (y - 1) \pmod{\Delta^2} \\ x^{-1} - 1 &\equiv -(x - 1) \pmod{\Delta^2}. \end{aligned}$$

The ideal Δ^n is spanned, over \mathbf{Q} , by the elements of the form

$$y_1(x_1 - 1)y_2(x_2 - 1) \cdots y_n(x_n - 1)y_{n+1},$$

where $x_i \in G$ and $y_j \in \mathbf{Q}[G]$, $1 \leq i \leq n$, $1 \leq j \leq n + 1$. Since

$$y = \sum_{g \in G} k_g g \equiv \sum_{g \in G} k_g \pmod{\Delta}, \quad k_g \in \mathbf{Q}$$

a basis for the quotient space Δ^n/Δ^{n+1} can be chosen among the images modulo Δ^{n+1} of the elements of the form

$$(a_{i_1} - 1)(a_{i_2} - 1) \cdots (a_{i_n} - 1),$$

where $a_{i_j} \in A$. But $(a_{i_1} - 1)(a_{i_2} - 1) \cdots (a_{i_n} - 1) = \sum_{g \in G} k'_g g$, where the summation extends over elements g of length at most n with respect to the system of generators A . \square

3.2. COROLLARY. *Let G be a finitely generated group and suppose that the ranks of $\gamma_n(G)/\gamma_{n+1}(G)$ grow exponentially. Then G has uniformly exponential growth and the estimate*

$$\lambda_*(G) \geq \limsup_{n \rightarrow \infty} \sqrt[n]{\text{rank}(\gamma_n(G)/\gamma_{n+1}(G))}$$

holds.

Recall that a group G is *parafree of para-rank m* if it is residually nilpotent and the factors of consecutive groups in its lower central series equal the corresponding ones of a free group of rank m . There are parafree groups which are not isomorphic to free groups [B 2,3].

3.3. PROPOSITION. *A finitely generated parafree group G of para-rank $m \geq 2$ has uniformly exponential growth and*

$$\lambda_*(G) \geq m.$$

Proof. It is known (see for instance [MKS: Thms. 5.11 (Witt's Formulae) and 5.12]) that for a free group \mathbf{F}_m the rank of $(\gamma_n(\mathbf{F}_m)/\gamma_{n+1}(\mathbf{F}_m))$ equals the n -th coefficient of the Maclaurin power series of the function $U(z) = 1/(1-mz)$ and the previous corollary can be applied. \square

Given a parafree group G of para-rank $m \geq 2$ it would be interesting to compare $\lambda_*(G)$ with $\lambda_*(\mathbf{F}_m) = 2m - 1$.

3.4. PROBLEM. *Is it true that, for a finitely generated para-free group G of para-rank $m \geq 2$ which is not free, one has $\lambda_*(G) > 2m - 1$?*

In order to formulate the next statement we recall the following

3.5. DEFINITION. An element $R \in F$ is said to be *primitive with respect to the lower central series* if, for all $n \geq 2$, it is not an n -th power modulo $\gamma_{\omega(R)+1}(F)$ where $\omega(R)$ is the weight of R . (The latter is defined by $R \in \gamma_{\omega(R)}(F)$ but $R \notin \gamma_{\omega(R)+1}(F)$.)

3.6. THEOREM ([L 1,2]). *Let R be an element of the free group F of finite rank m which is primitive with respect to the lower central series. Denote by $k = \omega(R)$ its weight and by $\langle R \rangle$ the normal closure of R in F . Let $G = F/\langle R \rangle$ and let $\mathcal{L}(F)$ and $\mathcal{L}(G)$ be the corresponding Lie algebras. Let then r be the image of R in $\mathcal{L}_k(F)$, the k -th component of $\mathcal{L}(F)$ and denote by I the ideal of $\mathcal{L}(F)$ generated by r .*

Then I is the kernel of the canonical homomorphism of $\mathcal{L}(F)$ onto $\mathcal{L}(G)$, i.e.

$$\mathcal{L}(G) = \mathcal{L}(F)/I.$$

Moreover for all $n \geq 1$ the abelian group $\mathcal{L}_n(G)$ is a torsion free group whose rank is the n -th coefficient of the Maclaurin power series of the function

$$U(z) = \frac{1}{1 - mz + z^k}.$$

4. MORE ON UNIFORMLY EXPONENTIAL GROWTH OF ONE-RELATOR GROUPS

Any two-generated one-relator group G can be presented in the form $G = \langle a, b : a^k w(a, b) = 1 \rangle$ where $k \in \mathbf{Z}$ and $w(a, b)$ belongs to the commutator subgroup $[F, F]$ of the free group $F = F(a, b)$ freely generated by a and b (this follows from Lemma 1.1). Since a and b constitute a basis in $F/\gamma_2(F)$ and $[a, b]$ generates $\gamma_2(F)/\gamma_3(F)$, one can also present G in the form

$$G = \langle a, b : a^k [a, b]^l w(a, b) = 1 \rangle$$

where $k, l \in \mathbf{Z}$ and $w(a, b) \in \gamma_3(F)$.

In this section we shall see that, under suitable assumptions on k, l and $w(a, b)$, the corresponding group has uniformly exponential growth.

As an application of Labute's Theorem we get the following:

4.1. PROPOSITION. *Let $G = \langle a, b : R(a, b) = 1 \rangle$ be such that R is primitive with respect to $\{\gamma_n(F)\}_{n=1}^{\infty}$ and $R \in \gamma_3(F)$. Then G has uniformly exponential growth.*

Proof. If $\omega(R) \geq 3$, Theorem 3.6 shows that the corresponding function $U(z)$ has a pole z_0 with $0 < z_0 < 1$. It follows that the coefficients $c_n(G)$ grow exponentially. By Corollary 3.2, $\lambda_*(G) > 1$. \square