

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AMENABILITY AND GROWTH OF ONE-RELATOR GROUPS
Autor: Ceccherini-Silberstein, Tullio G. / GRIGORCHUK, Rostislav I.
Kapitel: 4. More on uniformly exponential growth of one-relator groups
DOI: <https://doi.org/10.5169/seals-63284>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 31.03.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

3.6. THEOREM ([L 1,2]). *Let R be an element of the free group F of finite rank m which is primitive with respect to the lower central series. Denote by $k = \omega(R)$ its weight and by $\langle R \rangle$ the normal closure of R in F . Let $G = F/\langle R \rangle$ and let $\mathcal{L}(F)$ and $\mathcal{L}(G)$ be the corresponding Lie algebras. Let then r be the image of R in $\mathcal{L}_k(F)$, the k -th component of $\mathcal{L}(F)$ and denote by I the ideal of $\mathcal{L}(F)$ generated by r .*

Then I is the kernel of the canonical homomorphism of $\mathcal{L}(F)$ onto $\mathcal{L}(G)$, i.e.

$$\mathcal{L}(G) = \mathcal{L}(F)/I.$$

Moreover for all $n \geq 1$ the abelian group $\mathcal{L}_n(G)$ is a torsion free group whose rank is the n -th coefficient of the Maclaurin power series of the function

$$U(z) = \frac{1}{1 - mz + z^k}.$$

4. MORE ON UNIFORMLY EXPONENTIAL GROWTH OF ONE-RELATOR GROUPS

Any two-generated one-relator group G can be presented in the form $G = \langle a, b : a^k w(a, b) = 1 \rangle$ where $k \in \mathbf{Z}$ and $w(a, b)$ belongs to the commutator subgroup $[F, F]$ of the free group $F = F(a, b)$ freely generated by a and b (this follows from Lemma 1.1). Since a and b constitute a basis in $F/\gamma_2(F)$ and $[a, b]$ generates $\gamma_2(F)/\gamma_3(F)$, one can also present G in the form

$$G = \langle a, b : a^k [a, b]^l w(a, b) = 1 \rangle$$

where $k, l \in \mathbf{Z}$ and $w(a, b) \in \gamma_3(F)$.

In this section we shall see that, under suitable assumptions on k, l and $w(a, b)$, the corresponding group has uniformly exponential growth.

As an application of Labute's Theorem we get the following:

4.1. PROPOSITION. *Let $G = \langle a, b : R(a, b) = 1 \rangle$ be such that R is primitive with respect to $\{\gamma_n(F)\}_{n=1}^{\infty}$ and $R \in \gamma_3(F)$. Then G has uniformly exponential growth.*

Proof. If $\omega(R) \geq 3$, Theorem 3.6 shows that the corresponding function $U(z)$ has a pole z_0 with $0 < z_0 < 1$. It follows that the coefficients $c_n(G)$ grow exponentially. By Corollary 3.2, $\lambda_*(G) > 1$. \square

For Proposition 4.3 we need the following notations. Let ξ be a positive rational number such that $\xi \neq 1$ and denote by Q_ξ the smallest subgroup of the additive group of the rationals, which contains 1 and is invariant under multiplication by ξ and ξ^{-1} . In other words if $\xi = \frac{p}{q}$ with $p, q \in \mathbf{Z}$ and $\gcd(p, q) = 1$ then $Q_\xi \equiv \mathbf{Z}[\frac{1}{p}, \frac{1}{q}]$. Consider now the automorphism α of Q_ξ defined by $\alpha(x) = \xi x$, $x \in Q_\xi$. Let \mathbf{Z} act on Q_ξ by powers of α . Denote by $G_\xi = Q_\xi \rtimes_\alpha \mathbf{Z}$ the corresponding semidirect product. The group G_ξ is a two-generated group with system of generators $\{\bar{a}, \bar{b}\}$, where $\bar{a} = 1 \in Q_\xi$ and the element \bar{b} implements the automorphism $\alpha: \bar{b}^{-1}x\bar{b} = \alpha(x)$, $x \in Q_\xi$.

Let now d be a natural number ≥ 2 and set $B_d = \prod_{\mathbf{Z}} \mathbf{Z}_d$. The group \mathbf{Z} acts on B_d by shifts. The corresponding semidirect product $\Gamma(d)$, also denoted by $\mathbf{Z}_d \wr \mathbf{Z}$, is called the *wreath product* of \mathbf{Z} and \mathbf{Z}_d . We shall consider $\Gamma(d)$ as generated by $\bar{a} = (\dots, 0, 0, 1, 0, 0, \dots)$ where 1 denotes a generator of \mathbf{Z}_d (in the expression of \bar{a} it appears at the 0-th coordinate place), and by \bar{b} , the element which implements the shift.

We have short exact sequences

$$\begin{aligned} 0 &\longrightarrow Q_\xi \longrightarrow G_\xi \longrightarrow \mathbf{Z} \longrightarrow 0 \\ 0 &\longrightarrow B_d \longrightarrow \Gamma(d) \longrightarrow \mathbf{Z} \longrightarrow 0 \end{aligned}$$

so that G_ξ and $\Gamma(d)$ are two-step solvable. Slightly modifying the proof of Proposition 2.6 one gets

4.2. LEMMA. *The groups G_ξ and $\Gamma(d)$ have uniformly exponential growth.*

Our last class of two-generated one-relator groups of uniformly exponential growth is determined in the following statement.

4.3. PROPOSITION. *Let $G = \langle a, b; a^k[a, b]^l w(a, b) = 1 \rangle$ with $k, l \in \mathbf{Z}$ and $w(a, b) \in F^{(2)}$ where $F^{(2)} = [[F, F], [F, F]]$ denotes the second commutator subgroup of the free group $F = F(a, b)$ on a and b . Suppose that $(k, l) \notin \{\pm(2, 1), \pm(1, 1), \pm(1, 0), \pm(0, 1)\}$. Then G has uniformly exponential growth.*

Proof. Set $G_{k,l} = \langle a, b; a^k[a, b]^l w(a, b) \rangle$. Set also

$$\bar{G}_{k,l} = \langle a, b; a^k[a, b]^l w(a, b), F^{(2)} \rangle = \langle a, b; a^k[a, b]^l, F^{(2)} \rangle$$

which is a 2-step solvable quotient group of $G_{k,l}$. We shall show that $\bar{G}_{k,l}$ can be mapped homomorphically onto either G_ξ or $\Gamma(d)$ for a suitable positive rational number $\xi \neq 1$ or natural number $d \geq 2$.

Suppose first that $k \neq l, 2l$ and $lk \neq 0$. These assumptions guarantee that $\xi := \left| \frac{l-k}{l} \right| \neq 0, 1$. Then the map $a \mapsto (\bar{a})^{\text{sgn}(\frac{l-k}{l})}, b \mapsto \bar{b}$ from F onto G_ξ factorizes through $\bar{G}_{k,l}$. Indeed if we suppose, for instance, that $\frac{l-k}{l} > 0$, then the image of $a^k[a, b]^l$ is the number $k + l(-1 + \xi) \in \mathbf{Q}_\xi$ which is zero. Thus $\bar{G}_{k,l}$ maps onto G_ξ .

Suppose now that $\text{gcd}(k, l) = d$ or $(k, l) \in \{\pm(d, 0), \pm(0, d)\}$ for some $d \geq 2$. Then, the same arguments as before show that $\bar{G}_{k,l}$ can be mapped onto $\Gamma(d)$ via the map $a \mapsto \bar{a}, b \mapsto \bar{b}$.

Finally observe that $\bar{G}_{0,0}$ is the free two-generated two-step solvable group $F/F^{(2)}$ and thus maps homomorphically onto $\Gamma(d)$ for any $d \geq 2$.

The proof follows from Lemma 4.2. \square

Remark that the two-generated one-relator groups that are not covered by our statements have their relator that can be reduced to one of the form $bw, [a, b]w$ or $ba^{-1}baw$, where $w = w(a, b) \in F^{(2)}$.

Let us finish the paper by the following observation.

In [GrLP] it was conjectured that if G is a group with m generators and p relations, then

$$\lambda_*(G) \geq 2(m - p) - 1.$$

For one-relator groups there is one case when Gromov's conjecture holds true.

4.4. PROPOSITION. *Let $G = \langle a_1, a_2, \dots, a_m : R(a_1, a_2, \dots, a_m) = 1 \rangle$, with $m \geq 2$, be a one-relator group such that the relator R does not belong to the commutator subgroup F' of the free group F of rank m freely generated by a_1, a_2, \dots, a_m . Then $\lambda_*(G) \geq 2m - 3$.*

Proof. We may assume that G is torsion-free. Indeed if $U, V \in F$ are such $U = V^k$ for some $k \in \mathbf{Z}$, then $U \in F'$ iff $V \in F'$. If the relator R is a proper power, say $R = W^k$, then G maps onto $G_1 = \langle a_1, a_2, \dots, a_m : W(a_1, a_2, \dots, a_m) = 1 \rangle$, which is torsion-free, and $\lambda_*(G) \geq \lambda_*(G_1)$.

Under our assumptions on R , $H_1(G, \mathbf{Q}) \cong \mathbf{Z}^{m-1}$ and the second rational homology group $H_2(G, \mathbf{Q})$ vanishes.

In [S] it is proven that if $H_2(G, \mathbf{k}) = 0$, where \mathbf{k} is a field, then any subset $\{x_j\} \in G$, whose image in $H_1(G, \mathbf{k})$ is linearly independent, freely generates a free group.

Let $X = \{x_1, x_2, \dots, x_n\}$ be a finite system of generators for G . Then $\bar{X} = \{\bar{x}_1, \dots, \bar{x}_n\}$, where \bar{x}_i denotes the image of x_i in $H_1(G, \mathbf{Q})$, generates

$H_1(G, \mathbf{Q})$. We can find an independent subsystem $\{\bar{x}_{i_1}, \dots, \bar{x}_{i_{m-1}}\}$ in $H_1(G, \mathbf{Q})$ such that its pre-image $\{x_{i_1}, \dots, x_{i_m}\}$ freely generates a free group. Therefore $\lambda_X(G) \geq 2(m-1) - 1 = 2m - 3$. \square

It seems to us that for a one-relator group G of rank $m \geq 3$ the inequality $\lambda_*(G) \geq 2m - 3$ cannot be deduced directly from Magnus' Theorem as it is claimed in [GrLP].

ACKNOWLEDGMENTS. This paper was written during our stay at the Section de Mathématiques de l'Université de Genève. We thank Professor Pierre de la Harpe for his invitation and kind hospitality. We also thank him for his encouragement, stimulating conversations and suggestions.

We are grateful for various supports: the CNR for the first author, the RFFI grant 96-01-00713 for the second author, and the "Fonds National Suisse de la Recherche Scientifique" for both of us.

REFERENCES

- [A] AVEZ, A. Entropie des groupes de type fini. *C.R. Acad. Sc. Paris* 275, Série A (1972), 1363–1366.
- [B 1] BAUMSLAG, G. A survey of groups with a single defining relation. Proceedings of groups—St. Andrews 1985, p. 30–58. *London Math. Soc. Lecture Note Ser. 121*. Cambridge Univ. Press, 1986.
- [B 2,3] — Groups with the same lower central sequence as a relatively free group I. The groups. *Trans. Amer. Math. Soc.* 129 (1967), 308–321; II. Properties. *Trans. Amer. Math. Soc.* 142 (1969), 507–538.
- [BBV] BÉGUIN, C., H. BETTAIEB and A. VALETTE. K-Theory for C^* -algebras of one-relator groups. Preprint of the Institut de Mathématiques de l'Université de Neuchâtel (1996).
- [BC] BARTHOLDI, L. and T. CECCHERINI-SILBERSTEIN. Growth series and random walks on some hyperbolic graphs. Preprint Université de Genève (1996).
- [BCCH] BARTHOLDI, L., S. CANTAT, T. CECCHERINI-SILBERSTEIN and P. DE LA HARPE. Estimates for simple random walks on fundamental groups of surfaces. *Colloq. Math.* 72 (1997), 173–194.
- [Be] BEREZNYI, A.E. Discrete sub-exponential groups. *Zap. Nauchn. Sem. Leningrad. Otdel. Mat. Inst. Steklov. (LOMI)* 123 (1983), 155–166; English transl. in *J. Soviet Math.* 28 (1985), no. 4.
- [BP 1] BAUMSLAG, B. and S.J. PRIDE. Groups with one more generator than relators. *Math. Z.* 167 (1979), 279–281.