

2. Invariant and symmetric functions

Objektyp: **Chapter**

Zeitschrift: **L'Enseignement Mathématique**

Band (Jahr): **43 (1997)**

Heft 1-2: **L'ENSEIGNEMENT MATHÉMATIQUE**

PDF erstellt am: **21.09.2024**

Nutzungsbedingungen

Die ETH-Bibliothek ist Anbieterin der digitalisierten Zeitschriften. Sie besitzt keine Urheberrechte an den Inhalten der Zeitschriften. Die Rechte liegen in der Regel bei den Herausgebern. Die auf der Plattform e-periodica veröffentlichten Dokumente stehen für nicht-kommerzielle Zwecke in Lehre und Forschung sowie für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und den korrekten Herkunftsbezeichnungen weitergegeben werden. Das Veröffentlichen von Bildern in Print- und Online-Publikationen ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Die systematische Speicherung von Teilen des elektronischen Angebots auf anderen Servern bedarf ebenfalls des schriftlichen Einverständnisses der Rechteinhaber.

Haftungsausschluss

Alle Angaben erfolgen ohne Gewähr für Vollständigkeit oder Richtigkeit. Es wird keine Haftung übernommen für Schäden durch die Verwendung von Informationen aus diesem Online-Angebot oder durch das Fehlen von Informationen. Dies gilt auch für Inhalte Dritter, die über dieses Angebot zugänglich sind.

by the *power sum forms* in the results; to our knowledge this phenomenon has not been observed before. The combinatorial identities involving harmonic numbers that we encounter are also interesting. Sections 2-6 contain results in hermitian complex geometry and may be read without prior knowledge of Arakelov theory. §8 applies our calculations to obtain a presentation of the Arakelov Chow ring of the arithmetic Grassmannian.

This should be regarded as a companion paper to [T]; both papers will be part of the author's 1997 University of Chicago thesis. I wish to thank my advisor William Fulton for many useful conversations and exchanges of ideas.

2. INVARIANT AND SYMMETRIC FUNCTIONS

The symmetric group S_n acts on the polynomial ring $\mathbf{Z}[x_1, x_2, \dots, x_n]$ by permuting the variables, and the ring of invariants $\Lambda(n) = \mathbf{Z}[x_1, x_2, \dots, x_n]^{S_n}$ is the ring of symmetric polynomials. For $B = \mathbf{Q}$ or \mathbf{C} , let $\Lambda(n, B) = \Lambda(n) \otimes_{\mathbf{Z}} B$.

Let $e_k(x_1, \dots, x_n)$ be the k -th elementary symmetric polynomial in the variables x_1, \dots, x_n and $p_k(x_1, \dots, x_n) = \sum_i x_i^k$ the k -th power sum. The fundamental theorem on symmetric functions states that $\Lambda(n) = \mathbf{Z}[e_1, \dots, e_n]$ and that e_1, \dots, e_n are algebraically independent. For λ a partition, i.e. a decreasing sequence $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_m$ of nonnegative integers, define $p_\lambda := \prod_{i=1}^m p_{\lambda_i}$. It is well known that the p_λ 's form an additive \mathbf{Q} -basis for the ring of symmetric polynomials (cf. [M], §I.2). The two bases are related by Newton's identity:

$$(2) \quad p_k - e_1 p_{k-1} + e_2 p_{k-2} - \dots + (-1)^k k e_k = 0.$$

Another important set of symmetric functions related to the cohomology ring of grassmannians are the Schur polynomials. For a partition λ as above, the Schur polynomial s_λ is defined by

$$s_\lambda(x_1, \dots, x_n) = \frac{1}{\Delta} \cdot \det(x_i^{\lambda_j + n - j})_{1 \leq i, j \leq n},$$

where $\Delta = \prod_{1 \leq i < j \leq n} (x_i - x_j)$ is the Vandermonde determinant. The s_λ for all λ of length $m \leq n$ form a \mathbf{Z} -basis of $\Lambda(n)$ (cf. [M], §I.3).

Let $\mathbf{C}[T_{ij}]$ ($1 \leq i, j \leq n$) be the coordinate ring of the space $M_n(\mathbf{C})$ of $n \times n$ matrices. $GL_n(\mathbf{C})$ acts on matrices by conjugation, and we let

$I(n) = \mathbf{C}[T_{ij}]^{GL_n(\mathbf{C})}$ denote the corresponding graded ring of invariants. There is an isomorphism $\tau: I(n) \rightarrow \Lambda(n, \mathbf{C})$ given by evaluating an invariant polynomial ϕ on the diagonal matrix $\text{diag}(x_1, \dots, x_n)$. We will often identify ϕ with the symmetric polynomial $\tau(\phi)$. We will need to consider invariant polynomials with rational coefficients; let $I(n, \mathbf{Q}) \simeq \mathbf{Q}[x_1, x_2, \dots, x_n]^{S_n}$ be the corresponding ring.

Given $\phi \in I(n)_k$, let ϕ' be a k -multilinear form on $M_n(\mathbf{C})$ such that

$$\phi'(gA_1g^{-1}, \dots, gA_kg^{-1}) = \phi'(A_1, \dots, A_k)$$

for $g \in GL(n, \mathbf{C})$ and $\phi(A) = \phi'(A, A, \dots, A)$. Such forms are most easily constructed for the power sums p_k by setting

$$p'_k(A_1, A_2, \dots, A_k) = \text{Tr}(A_1A_2 \cdots A_k).$$

For p_λ we can take $p'_\lambda = \prod p'_{\lambda_i}$. Since the p_λ 's are a basis of $\Lambda(n, \mathbf{Q})$, it follows that one can use the above constructions to find multilinear forms ϕ' for any $\phi \in I(n)_k$.

An explicit formula for ϕ' is given by polarizing ϕ :

$$\phi'(A_1, \dots, A_k) = \frac{(-1)^k}{k!} \sum_{j=1}^k \sum_{i_1 < \dots < i_j} (-1)^j \phi(A_{i_1} + \dots + A_{i_j}).$$

Although above formula for ϕ' is symmetric in A_1, \dots, A_k , this property is not needed for the applications that follow.

3. HERMITIAN DIFFERENTIAL GEOMETRY

Let X be a complex manifold, E a rank n holomorphic vector bundle over X . Denote by $A^k(X, E)$ the C^∞ sections of $\Lambda^k T^*X \otimes E$, where T^*X denotes the cotangent bundle of X . In particular $A^k(X)$ is the space of smooth complex k -forms on X . Let $A^{p,q}(X)$ the space of smooth complex forms of type (p, q) on X and $A(X) := \bigoplus_p A^{p,p}(X)$. The decomposition $A^1(X, E) = A^{1,0}(X, E) \oplus A^{0,1}(X, E)$ induces a decomposition $D = D^{1,0} + D^{0,1}$ of each connection D on E . Let $d = \partial + \bar{\partial}$ and $d^c = (\partial - \bar{\partial})/(4\pi i)$.

Assume now that E is equipped with a hermitian metric h . The pair (E, h) is called a *hermitian vector bundle*. The metric h induces a canonical connection $D = D(h)$ such that $D^{0,1} = \bar{\partial}_E$ and D is *unitary*, i.e.

$$dh(s, t) = h(Ds, t) + h(s, Dt), \quad \text{for all } s, t \in A^0(X, E).$$