Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: BOTT-CHERN FORMS AND ARITHMETIC INTERSECTIONS
Autor: Tamvakis, Harry
Kapitel: 2. Invariant and symmetric functions
DOI: https://doi.org/10.5169/seals-63270

Nutzungsbedingungen
Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation
L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use
The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 11.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

by the *power sum forms* in the results; to our knowledge this phenomenon has not been observed before. The combinatorial identities involving harmonic numbers that we encounter are also interesting. Sections 2-6 contain results in hermitian complex geometry and may be read without prior knowledge of Arakelov theory. §8 applies our calculations to obtain a presentation of the Arakelov Chow ring of the arithmetic Grassmannian.

This should be regarded as a companion paper to [T]; both papers will be part of the author's 1997 University of Chicago thesis. I wish to thank my advisor William Fulton for many useful conversations and exchanges of ideas.

2. Invariant and symmetric functions

The symmetric group S_n acts on the polynomial ring $\mathbf{Z}[x_1, x_2, \ldots, x_n]$ by permuting the variables, and the ring of invariants $\Lambda(n) = \mathbf{Z}[x_1, x_2, \ldots, x_n]^{S_n}$ is the ring of symmetric polynomials. For $B = \mathbf{Q}$ or $\mathbf{C}$, let $\Lambda(n, B) = \Lambda(n) \otimes_{\mathbf{Z}} B$.

Let $e_k(x_1, \ldots, x_n)$ be the k-th elementary symmetric polynomial in the variables $x_1, \ldots, x_n$ and $p_k(x_1, \ldots, x_n) = \sum_i x_i^k$ the k-th power sum. The fundamental theorem on symmetric functions states that $\Lambda(n) = \mathbf{Z}[e_1, \ldots, e_n]$ and that $e_1, \ldots, e_n$ are algebraically independent. For λ a partition, i.e. a decreasing sequence $\lambda_1 \geqslant \lambda_2 \geqslant \cdots \geqslant \lambda_m$ of nonnegative integers, define $p_\lambda := \prod_{i=1}^{m} p_{\lambda_i}$. It is well known that the p_λ's form an additive $\mathbf{Q}$-basis for the ring of symmetric polynomials (cf. [M], §I.2). The two bases are related by Newton's identity:

$$p_k - e_1 p_{k-1} + e_2 p_{k-2} - \cdots + (-1)^k k e_k = 0 . \tag{2}$$

Another important set of symmetric functions related to the cohomology ring of grassmannians are the Schur polynomials. For a partition λ as above, the Schur polynomial s_λ is defined by

$$s_\lambda(x_1, \ldots, x_n) = \frac{1}{\Delta} \cdot \det(x_i^{\lambda_j + n - j})_{1 \leqslant i, j \leqslant n} ,$$

where $\Delta = \prod_{1 \leqslant i < j \leqslant n} (x_i - x_j)$ is the Vandermonde determinant. The s_λ for all λ of length $m \leqslant n$ form a $\mathbf{Z}$-basis of $\Lambda(n)$ (cf. [M], §I.3).

Let $\mathbf{C}[T_{ij}]$ $(1 \leqslant i, j \leqslant n)$ be the coordinate ring of the space $M_n(\mathbf{C})$ of $n \times n$ matrices. $GL_n(\mathbf{C})$ acts on matrices by conjugation, and we let

$I(n) = \mathbf{C}[T_{ij}]^{GL_n(\mathbf{C})}$ denote the corresponding graded ring of invariants. There is an isomorphism $\tau\colon I(n) \to \Lambda(n, \mathbf{C})$ given by evaluating an invariant polynomial ϕ on the diagonal matrix $\operatorname{diag}(x_1, \ldots, x_n)$. We will often identify ϕ with the the symmetric polynomial $\tau(\phi)$. We will need to consider invariant polynomials with rational coefficients; let $I(n, \mathbf{Q}) \simeq \mathbf{Q}[x_1, x_2, \ldots, x_n]^{S_n}$ be the corresponding ring.

Given $\phi \in I(n)_k$, let ϕ' be a k-multilinear form on $M_n(\mathbf{C})$ such that

$$\phi'(gA_1g^{-1}, \ldots, gA_kg^{-1}) = \phi'(A_1, \ldots, A_k)$$

for $g \in GL(n, \mathbf{C})$ and $\phi(A) = \phi'(A, A, \ldots, A)$. Such forms are most easily constructed for the power sums p_k by setting

$$p'_k(A_1, A_2, \ldots, A_k) = \operatorname{Tr}(A_1A_2 \cdots A_k).$$

For p_λ we can take $p'_\lambda = \prod p'_{\lambda_i}$. Since the p_λ's are a basis of $\Lambda(n, \mathbf{Q})$, it follows that one can use the above constructions to find multilinear forms ϕ' for any $\phi \in I(n)_k$.

An explicit formula for ϕ' is given by polarizing ϕ:

$$\phi'(A_1 \ldots, A_k) = \frac{(-1)^k}{k!} \sum_{j=1}^{k} \sum_{i_1 < \ldots < i_j} (-1)^j \phi(A_{i_1} + \ldots + A_{i_j}).$$

Although above formula for ϕ' is symmetric in $A_1 \ldots, A_k$, this property is not needed for the applications that follow.

3. Hermitian differential geometry

Let X be a complex manifold, E a rank n holomorphic vector bundle over X. Denote by $A^k(X, E)$ the C^∞ sections of $\Lambda^k T^*X \otimes E$, where T^*X denotes the cotangent bundle of X. In particular $A^k(X)$ is the space of smooth complex k-forms on X. Let $A^{p,q}(X)$ the space of smooth complex forms of type (p, q) on X and $A(X) := \bigoplus_p A^{p,p}(X)$. The decomposition $A^1(X, E) = A^{1,0}(X, E) \bigoplus A^{0,1}(X, E)$ induces a decomposition $D = D^{1,0} + D^{0,1}$ of each connection D on E. Let $d = \partial + \overline{\partial}$ and $d^c = (\partial - \overline{\partial})/(4\pi i)$.

Assume now that E is equipped with a hermitian metric h. The pair (E, h) is called a *hermitian vector bundle*. The metric h induces a canonical connection $D = D(h)$ such that $D^{0,1} = \overline{\partial}_E$ and D is *unitary*, i.e.

$$d\,h(s, t) = h(Ds, t) + h(s, Dt), \qquad \text{for all } s, t \in A^0(X, E).$$