

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: BOTT-CHERN FORMS AND ARITHMETIC INTERSECTIONS
Autor: Tamvakis, Harry
Kapitel: 8. Arakelov Chow rings of grassmannians
DOI: <https://doi.org/10.5169/seals-63270>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

8. ARAKELOV CHOW RINGS OF GRASSMANNIANS

In this section $G = G(r, n)$ will denote the Grassmannian over $\text{Spec } \mathbf{Z}$. Over any field k , G parametrizes the r -dimensional linear subspaces of a vector space over k . Let

$$(13) \quad \bar{\mathcal{E}} : 0 \rightarrow \bar{S} \rightarrow \bar{E} \rightarrow \bar{Q} \rightarrow 0$$

denote the universal exact sequence of vector bundles over G . Here the trivial bundle $E(\mathbf{C})$ is given the trivial metric and the tautological subbundle $S(\mathbf{C})$ and quotient bundle $Q(\mathbf{C})$ the induced metrics. The homogeneous space $G(\mathbf{C}) \simeq U(n)/(U(r) \times U(n-r))$ is a complex manifold. $G(\mathbf{C})$ is endowed with a natural $U(n)$ -invariant metric coming from the Kähler form $\eta_G = c_1(\overline{Q(\mathbf{C})})$.

G is a smooth arithmetic scheme and $G(\mathbf{C})$ with the metric coming from η_G is a hermitian symmetric space, so we have an Arakelov Chow ring $CH(\bar{G})$. Note that since the hermitian vector bundles in (13) are invariant under the action of $U(n)$, their Chern forms are harmonic, and thus the arithmetic characteristic classes obtained are all elements of $CH(\bar{G})$. Maillot [Ma] found a presentation of $CH(\bar{G})$, using the above observation and the short exact sequence (11). We wish to offer another description of this ring, based on the calculations in this paper.

First recall the geometric picture: for the ordinary Chow ring we have

$$(14) \quad CH(G) = \frac{\mathbf{Z}[c(S), c(Q)]}{\langle c(S)c(Q) = 1 \rangle}.$$

(see for instance [F], Example 14.6.6). If x_1, \dots, x_r are the Chern roots of S , y_1, \dots, y_s are the Chern roots of Q , $H = S_r \times S_{n-r}$ is the product of two symmetric groups, and t is a formal variable, then (14) can be rewritten

$$(15) \quad CH(G) = \frac{\mathbf{Z}[x_1, \dots, x_r, y_1, \dots, y_s]^H}{\langle \prod_i (1 + x_i t) \prod_j (1 + y_j t) = 1 \rangle}.$$

Maillot's presentation of $CH(\bar{G})$ is an analogue of (14); ours will be an analogue of (15). We introduce $2n$ variables

$$\hat{x}_1, \dots, \hat{x}_r, \hat{y}_1, \dots, \hat{y}_s, x_1, \dots, x_r, y_1, \dots, y_s$$

and consider the rings

$$A = \mathbf{Z}[\hat{x}_1, \dots, \hat{x}_r, \hat{y}_1, \dots, \hat{y}_s]^H \quad \text{and} \quad B = \mathbf{R}[x_1, \dots, x_r, y_1, \dots, y_s]^H$$

and the ring homomorphism $\omega : A \rightarrow B$ defined by $\omega(\hat{x}_i) = x_i$ and $\omega(\hat{y}_j) = y_j$. A ring structure is defined on the abelian group $A \oplus B$ by setting

$$(\widehat{x}, x') * (\widehat{y}, y') = (\widehat{xy}, \omega(\widehat{x})y' + x'\omega(\widehat{y})).$$

We will adopt the convention that $\widehat{\alpha}$ denotes $(\widehat{\alpha}, 0)$, β denotes $(0, \beta)$, and any product $\prod x_i y_j$ denotes $(0, \prod x_i y_j)$; the multiplication $*$ is thus characterized by the properties $\widehat{\alpha} * \beta = \alpha\beta$ and $\beta_1 * \beta_2 = 0$. We now define two sets of relations in $(A \oplus B)[t]$:

$$\mathcal{R}_1 : \prod_i (1 + x_i t) \prod_j (1 + y_j t) = 1,$$

$$\mathcal{R}_2 : \prod_i (1 + \widehat{x}_i t) * \prod_j (1 + \widehat{y}_j t) * \left(1 + t \sum_j \frac{\log(1 + y_j t)}{1 + y_j t} \right) = 1,$$

and let \mathcal{A} denote the quotient of the graded ring $A \oplus B$ by these relations. Using this notation we can state

THEOREM 6. *There is a unique ring isomorphism $\Phi : \mathcal{A} \rightarrow CH(\overline{G})$ such that*

$$\begin{aligned} \Phi\left(\prod_i (1 + \widehat{x}_i t^i)\right) &= \sum_i \widehat{c}_i(\overline{S})t^i, & \Phi\left(\prod_j (1 + \widehat{y}_j t^j)\right) &= \sum_j \widehat{c}_j(\overline{Q})t^j, \\ \Phi\left(\prod_i (1 + x_i t^i)\right) &= \sum_i a(c_i(\overline{S}))t^i, & \Phi\left(\prod_j (1 + y_j t^j)\right) &= \sum_j a(c_j(\overline{Q}))t^j. \end{aligned}$$

Proof. The isomorphism Φ of \mathcal{A} with $CH(\overline{G})$ is obtained exactly as in [Ma], Theorem 4.0.5. The key fact is that since G has a cellular decomposition (in the sense of [F], Ex. 1.9.1), it follows that $CH^{p,p-1}(G) = 0$ for all p (using the excision exact sequence for the groups $CH^{*,*}(G)$; cf. [G], §8). Summing the sequence (11) over all p gives

$$(16) \quad 0 \longrightarrow \mathcal{H}(G_{\mathbf{R}}) \xrightarrow{a} CH(\overline{G}) \xrightarrow{\zeta} CH(G) \longrightarrow 0.$$

We can now use our knowledge of the rings $\mathcal{H}(G_{\mathbf{R}})$ and $CH(G)$ together with the five lemma, as in loc. cit. The multiplication $*$ is a consequence of the general identity $a(x)y = a(x\omega(y))$ in $\widehat{CH}(G)$. To complete the argument we must show that the relation $\widehat{c}(\overline{S})\widehat{c}(\overline{Q}) = 1 + a(\widehat{c}(\overline{E}))$ translates to the relation \mathcal{R}_2 above.

Let $p_i(y)$ be the i -th power sum in the variables y_1, \dots, y_s , identified under Φ with the class $a(p_i(\overline{Q}))$ in $CH(\overline{G})$ (we will use such identifications freely in the sequel). We also define $p_a(t) = \sum_{i=0}^{\infty} (-1)^{i+1} \mathcal{H}_i p_i(y) t^{i+1}$. Proposition 3 implies that

$$(17) \quad \widehat{c}_t(\bar{S})\widehat{c}_t(\bar{Q}) = 1 + a(\widehat{c}_t(\bar{E})) = 1 - p_a(t),$$

where the subscript t denotes the corresponding Chern polynomial. Multiplying both sides of (17) by $1 + p_a(t)$ and using the properties of multiplication in \mathcal{A} gives the equivalent form

$$(18) \quad \widehat{c}_t(\bar{S}) * \widehat{c}_t(\bar{Q}) * (1 + p_a(t)) = 1.$$

We now note that the *harmonic number generating function*

$$\sum_{i=0}^{\infty} \mathcal{H}_i t^i = \frac{t}{1-t} + \frac{1}{2} \frac{t^2}{1-t} + \frac{1}{3} \frac{t^3}{1-t} + \cdots = \frac{\log(1-t)}{t-1}.$$

It follows that

$$p_a(-t) = \sum_{i=0}^{\infty} \mathcal{H}_i p_i(y) t^{i+1} = t \sum_{j=1}^s \sum_{i=0}^{\infty} \mathcal{H}_i (y_j t)^i = -t \sum_{j=1}^s \frac{\log(1 - y_j t)}{1 - y_j t}$$

and thus

$$p_a(t) = t \sum_{j=1}^s \frac{\log(1 + y_j t)}{1 + y_j t}.$$

Substituting this in equation (18) gives relation \mathcal{R}_2 . \square

Theorem 6 shows that the relations in the Arakelov Chow ring of G are the classical geometric ones perturbed by a new “arithmetic factor” of $1 + p_a(t)$. While this factor is closely related to the power sums $p_i(\bar{Q})$, the most natural basis of symmetric functions for doing calculations in $CH(G)$ is the basis of Schur polynomials (corresponding to the Schubert classes; see for example [F], §14.7). The arithmetic analogues of the special Schubert classes involve the power sum perturbation above; multiplication formulas are thus quite complicated (see [Ma]).

In geometry the Chern roots x_i and y_j all “live” on the complete flag variety above G . There are certainly natural line bundles on the flag variety whose first Chern classes correspond to the roots in Theorem 6. However on flag varieties the situation is more complicated and our knowledge is not as complete. We refer the reader to [T] for more details.

REFERENCES

- [A] ARAKELOV, S. J. Intersection theory of divisors on an arithmetic surface. *Math. USSR Izvestiya* 8 (1974), 1167–1180.
- [BiGS] BISMUT, J.-M., H. GILLET and C. SOULÉ. Analytic torsion and holomorphic determinant bundles I. Bott-Chern forms and analytic torsion. *Comm. Math. Phys.* 115 (1988), 49–78.