

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 43 (1997)
Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: ATKIN-LEHNER EIGENFORMS AND STRONGLY MODULAR LATTICES
Autor: QUEBBEMANN, H.-G.
Kapitel: 2. ATKIN-LEHNER ACTION ON THETA FUNCTIONS
DOI: <https://doi.org/10.5169/seals-63271>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 27.04.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

EXAMPLE 4. Similarly, $K = \mathbf{Q}\left(e\left(\frac{1}{8}\right), \sqrt{\pm q}\right)$ may be used to obtain strongly modular lattices Λ in $G_{8r}(2^+q^+)$ from rank r unimodular hermitian lattices over \mathcal{O}_K , setting now

$$x \cdot y = \frac{1}{2} \operatorname{Tr}_{K/\mathbf{Q}}(h(x, y)/(2 - \sqrt{2})).$$

Again, we always have $\min \Lambda \geq 4$. E.g., \mathcal{O}_K itself gives the tensor product of D_4 and the binary lattice with Gram matrix $\begin{pmatrix} 2 & 1 \\ 1 & (q+1)/2 \end{pmatrix}$.

2. ATKIN-LEHNER ACTION ON THETA FUNCTIONS

The subject treated in this section is not new, but appears to be difficult to cite from the literature (in the form we need it). For convenience, I give a rather detailed account, starting from a classical formula (due to Jacobi and others). Let Λ be an even lattice. The theta function of a coset $\bar{v} = v + \Lambda$ in Λ^* (and, in particular, Θ_Λ for $v = 0$) is that function defined on the upper half-plane by

$$\Theta_{\bar{v}}(z) = \sum_{x \in \bar{v}} e\left(\frac{1}{2}(x \cdot x)z\right).$$

Now let $n = 2k$ (k integral), and recall that $SL_2(\mathbf{R})$ acts on functions f on the upper half-plane by

$$(f |_k S)(z) = (cz + d)^{-k} f\left(\frac{az + b}{cz + d}\right), \quad S = \begin{pmatrix} a & b \\ c & d \end{pmatrix}.$$

Let S be in $SL_2(\mathbf{Z})$, with $c > 0$. For $u, v \in \Lambda^*$ define

$$\phi_S(u, v) = \sum_x e((ax \cdot x + 2x \cdot v + dv \cdot v)/2c)$$

where x runs through a system of representatives of those elements of $\Lambda^*/c\Lambda$ which reduce to \bar{u} in $D = \Lambda^*/\Lambda$. Each summand clearly depends only on the class $x + c\Lambda$, and the whole sum depends only on \bar{u} and \bar{v} . The latter statement is trivial for \bar{u} , while for \bar{v} it is proved (using $1 = ad - bc$) by

$$\begin{aligned} \phi_S(u, v) &= \sum_x e(a(x + dv) \cdot (x + dv)/2c) e(-b(2x \cdot v + dv \cdot v)/2) \\ (2.1) \quad &= \phi_S(u + dv, 0) e(-b(2u \cdot v + dv \cdot v)/2). \end{aligned}$$

So we may write $\phi_S(u, v) = \phi_S(\bar{u}, \bar{v})$. Then the formula we need is (see [Mi], p. 189)

$$(2.2) \quad \Theta_{\bar{u}}|_k S = (\det \Lambda)^{-\frac{1}{2}} (ic)^{-k} \sum_{\bar{v} \in D} \phi_S(\bar{u}, \bar{v}) \Theta_{\bar{v}}.$$

Let Λ be of level ℓ . Then it is a well-known consequence of (2.2) that Θ_{Λ} is a modular form for the subgroup $\Gamma_0(\ell)$ of $SL_2(\mathbf{Z})$ defined by $c \equiv 0 \pmod{\ell}$. Now let $m|\ell$, $m' = \ell/m$. After [AL] the m -th Atkin-Lehner involution $\Gamma_0(\ell)W_m$, a coset of $\Gamma_0(\ell)$ in its normalizer in $SL_2(\mathbf{R})$, is given by any matrix of the form

$$W_m = S \begin{pmatrix} \sqrt{m} & 0 \\ 0 & 1/\sqrt{m} \end{pmatrix}, \quad S = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma_0(m'), \quad d \equiv 0 \pmod{m}.$$

We specifically choose $a = 1$, $c = m'$, solve $tm + t'm' = 1$, and put $b = -t'$, $d = tm$. In any case,

$$W_m^2 \equiv 1, \quad W_m W_{m'} \equiv W_{\ell} \pmod{\Gamma_0(\ell)}.$$

Now these involutions are connected, via Gaussian sums, to those in Section 1 by the following important relation which seems to have been proved first by Kitaoka ([Ki] where, however, the constant factor is not worked out; cf. also [BS], p. 77, for the statement of a generalization).

ATKIN-LEHNER IDENTITY:

$$(2.3) \quad \Theta_{\Lambda}|_k W_m = \left(\frac{\det \Lambda_m}{\det \Lambda} \right)^{\frac{1}{4}} i^{-k} g_{m'}(\Lambda_{m'}) \Theta_{\Lambda_m}.$$

How this identity will be used here is readily explained. Suppose that Λ (and then also Λ_m) has determinant ℓ^k . We will be interested in the following gradually stronger conditions

- (1) $\Theta_{\Lambda} = \Theta_{\Lambda_m}$ for all $m|\ell$
- (2) Λ is strongly modular.

So condition (1) says that Θ_{Λ} is an eigenform of all the Atkin-Lehner involutions; Section 3 deals with such modular forms.

Proof of (2.3). Recall that here S is chosen such that $a = 1$, $c = m'$ and $m|d$. Write $\bar{v} \in D$ in the form $\bar{v} = \bar{w} + \bar{y}$ where $\bar{w} \in D(m)$, $\bar{y} \in D(m')$. Then $\phi_S(0, \bar{v}) = \phi_S(0, \bar{y})$ by (2.1), and (2.2) gives

$$\begin{aligned} (\det \Lambda)^{\frac{1}{2}} (im')^k \Theta_{\Lambda}|_k S &= \sum_{\bar{v} \in D} \phi_S(0, \bar{v}) \Theta_{\bar{v}} \\ &= \sum_{\bar{y} \in D(m')} \phi_S(0, \bar{y}) \sum_{\bar{w} \in D(m)} \Theta_{\bar{w} + \bar{y}}. \end{aligned}$$

We first must know that $\phi_S(0, \bar{y}) = 0$ for $\bar{y} \neq 0$. To see this, write $x \in \Lambda$ in the form $x = x_1 + m'x'$ with x_1 from some system of representatives of $\Lambda/\sqrt{m'}\Lambda_{m'}$ and $\sqrt{m'}x' \in \Lambda_{m'}$. Then

$$\phi_S(0, y) e(-dy \cdot y/2m') = \sum_{x_1} e((x_1 \cdot x_1 + 2x_1 \cdot y)/2m') \sum_{\bar{x}' \in D(m')} e(x' \cdot y).$$

But the sum over $D(m')$ vanishes for $\bar{y} \neq 0$ (since $D(m')$ is regular with respect to the discriminant form). It remains to determine $\phi_S(0, 0)$. Put $E = \Lambda_{m'}^*/\Lambda_{m'}$. Since $(1/\sqrt{m'})\Lambda/\Lambda_{m'}$ is $E(m')$, the last formula for $y = 0$ gives

$$\begin{aligned} \phi_S(0, 0) &= \#D(m') \sum_{\bar{v} \in E(m')} e\left(\frac{1}{2}v \cdot v\right) \\ &= \#D(m') (\#E(m'))^{\frac{1}{2}} g_{m'}(\Lambda_{m'}) \end{aligned}$$

where

$$\begin{aligned} \#D(m') (\#E(m'))^{\frac{1}{2}} &= (\Lambda_{m'} : \sqrt{m'}\Lambda) (\sqrt{m'}\Lambda : m'\Lambda_{m'})^{\frac{1}{2}} \\ &= (\#D(m'))^{\frac{1}{2}} (m')^k \\ &= (\#D(m))^{-\frac{1}{2}} (\det \Lambda)^{\frac{1}{2}} (m')^k \\ &= m^{-\frac{k}{2}} \left(\frac{\det \Lambda_m}{\det \Lambda}\right)^{\frac{1}{4}} (\det \Lambda)^{\frac{1}{2}} (m')^k \quad \square \end{aligned}$$

Note that, by Propositions 1 and 2, (2.3) in the case $\det \Lambda = \ell^k$ becomes

$$(2.4) \quad \Theta_\Lambda|_k W_m = g_m(\Lambda_{m'})^{-1} \Theta_{\Lambda_m}.$$

3. SOME USE OF MODULAR FORMS

Let $W(\ell)$ be the elementary abelian 2-group formed by the Atkin-Lehner involutions $w_m = \Gamma_0(\ell)W_m$ for all $m \parallel \ell$. Let k be even, and let $\mathcal{S}_k(\ell)$ denote the space of cusp forms of weight k on $\Gamma_0(\ell)$. Then $W(\ell)$ acts on this space. For a character χ of $W(\ell)$ we let $\mathcal{S}_k(\ell)_\chi$ denote the subspace on which $W(\ell)$ acts by χ . If Λ and M are lattices of dimension $2k$ and level ℓ belonging to the same genus, then $f = \Theta_\Lambda - \Theta_M$ is known to be in $\mathcal{S}_k(\ell)$, and when both lattices are strongly modular identity (2.4) implies that f is in $\mathcal{S}_k(\ell)_\chi$ for the character $\chi(w_m) = g_m(\Lambda)$. So we are interested in such spaces now.

Fortunately, the dimension of $\mathcal{S}_k(\ell)_\chi$ is known; I am indebted to N.-P. Skoruppa for pointing out the reference. Let $s(\ell)$ denote the number of prime