

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	43 (1997)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	THE NORMALISER ACTION AND STRONGLY MODULAR LATTICES
Autor:	Nebe, Gabriele
Kapitel:	2. PRELIMINAIRES AND NOTATION
DOI:	https://doi.org/10.5169/seals-63272

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 19.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

of G in $GL_d(\mathbf{Q})$ to test strong modularity of L . In the next section we derive some methods for explicitly constructing elements of $N_{GL_d(\mathbf{Q})}(G)$.

Every finite subgroup of $GL_d(\mathbf{Q})$ is a subgroup of the automorphism group of an integral lattice. In particular the maximal finite subgroups of $GL_d(\mathbf{Q})$ are automorphism groups of distinguished lattices. A subgroup of $GL_d(\mathbf{Q})$ is called rational irreducible if it does not preserve a proper subspace $\neq \{0\}$ of \mathbf{Q}^d . The rational irreducible maximal finite, abbreviated to *r.i.m.f.*, subgroups of $GL_d(\mathbf{Q})$ are classified for $d < 32$ (cf. [PIN 95], [NeP 95], [Neb 95], [Neb 96], [Neb 96a]). Their invariant lattices provide many examples of strongly modular lattices. The following theorem is proved by applying the methods derived in Section 4.

THEOREM. *In dimension $d < 32$, all even lattices $L \subseteq \mathbf{R}^d$ that are preserved by a r.i.m.f. group and satisfy $L^\# / L \cong (\mathbf{Z}/l\mathbf{Z})^{d/2}$ for some $l \in \mathbf{N}$ are strongly modular, except for the lattices of the r.i.m.f. group $[\pm \text{Alt}_6 . 2^2]_{16}$ in $GL_{16}(\mathbf{Q})$ (cf. [NeP 95]).*

2. PRELIMINARIES AND NOTATION

The main strategy in this paper is the application of the following *normaliser principle*.

Let G be a group acting on a set S , H a subgroup of the group of transformations of S . Then the normaliser of G in H acts on the set of G -orbits.

In our situation $G = \text{Aut}(L)$ is the automorphism group of an integral lattice L in the Euclidean space $\mathbf{R}L \cong \mathbf{R}^d$. By writing the action of G on $\mathbf{R}L$ with respect to a \mathbf{Z} -basis (b_1, \dots, b_d) of L , G becomes a finite subgroup of $GL_d(\mathbf{Z})$. Then $G = \text{Aut}(F) = \{g \in GL_d(\mathbf{Z}) \mid gFg^{tr} = F\}$ where F is the Gram matrix $F = ((b_i, b_j))_{i,j=1}^d$ of L .

For the rest of this article let $H = GL_d(\mathbf{Q})$, $G \leq H$, be a finite subgroup of H , and let $N := N_H(G)$ be its normaliser. We also assume that G contains the negative unit matrix, $-I_d \in G$.

We apply the normaliser principle to the following three situations.

- (i) $S = \{L \subseteq \mathbf{Q}^d \mid L = \sum_{i=1}^d \mathbf{Z}b_i \text{ for a basis } (b_1, \dots, b_d) \text{ of } \mathbf{Q}^d\}$, the set of \mathbf{Z} -lattices of rank d in \mathbf{Q}^d , and the action of H on S is right multiplication: $S \times H \rightarrow S$, $(L, h) \mapsto Lh := \{lh \mid l \in L\}$. Then the set of G -fixed points is

$$\mathcal{Z}(G) := \{L \in S \mid Lg = L \text{ for all } g \in G\},$$

the set of G -invariant lattices.

- (ii) $S = \{F \in M_d(\mathbf{Q}) \mid F = F^{tr}, F \text{ positive definite}\}$, the set of positive definite symmetric matrices, where x^{tr} denotes the transposed matrix of $x \in M_d(\mathbf{Q})$ and the action of H on S is $S \times H \rightarrow S$, $(F, h) \mapsto hFh^{tr}$. Then the set of G -fixed points is

$$\mathcal{F}_{>0}(G) := \{F \in S \mid gFg^{tr} = F \text{ for all } g \in G\}.$$

Note that $(\mathbf{R}_{>0})\mathcal{F}_{>0}(G)$ is the set of G -invariant Euclidean scalar products on \mathbf{R}^d . G is called *uniform*, if there is essentially one G -invariant Euclidean structure on \mathbf{R}^d , that is if $\mathcal{F}_{>0}(G) = \{aF \mid 0 < a \in \mathbf{Q}\}$ for some $F \in M_d(\mathbf{Q})$.

- (iii) $S = M_d(\mathbf{Q})$, and the action of H is conjugation: $S \times H \rightarrow S$, $(c, h) \mapsto h^{-1}ch$. Then the set of G -fixed points is the *commuting algebra* of G

$$C_{M_d(\mathbf{Q})}(G) := \{c \in M_d(\mathbf{Q}) \mid cg = gc \text{ for all } g \in G\}.$$

The following two remarks follow immediately from the normaliser principle.

REMARK 1. Assume that G is uniform and let $F \in \mathcal{F}_{>0}(G)$. Then for each $n \in N$, the matrix nFn^{tr} is also G -invariant and therefore $nFn^{tr} = (\det(n))^{2/d}F$. Hence n induces a similarity of F .

REMARK 2. For $n \in N$ and $L \in \mathcal{Z}(G)$, the lattice $Ln \in \mathcal{Z}(G)$ is also G -invariant.

3. SIMILARITIES NORMALISE

In this section we show that if G is the automorphism group of a (strongly modular) lattice L then the similarities between L and $L' \in \pi(L)$ are elements of N .

PROPOSITION 3. Let $G = \text{Aut}(F) \leq GL_d(\mathbf{Z})$ be the full automorphism group of a lattice L . Assume that L is an integral lattice. Let $L' \in \pi(L)$ and $n \in GL_d(\mathbf{Q})$ which induces a similarity from L' to L , i.e. $L'n = L$ and $nFn^{tr} = aF$, ($a \in \mathbf{N}$). Then $a^{-1}n^2 \in G$ and $n \in N$.

Proof. The matrix $a^{-1}n^2$ is clearly orthogonal with respect to F . Therefore to prove that $a^{-1}n^2 \in G$ we only have to show that $La^{-1}n^2 = L$. Now $L' = Ln^{-1}$, hence its dual lattice is

$$(L')^\# = \{v \in \mathbf{Q}^d \mid vF(ln^{-1})^{tr} \in \mathbf{Z} \text{ for all } l \in L\}.$$