

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	43 (1997)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	THE NORMALISER ACTION AND STRONGLY MODULAR LATTICES
Autor:	Nebe, Gabriele
Kapitel:	5. Proof of the Theorem
DOI:	https://doi.org/10.5169/seals-63272

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 13.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

5. PROOF OF THE THEOREM

In this section we prove the theorem stated in the Introduction. The principle of the proof is given in the following remark.

REMARK 7. *Let $G = \text{Aut}(F) \leq GL_d(\mathbf{Z})$ be a uniform automorphism group of the lattice $L = \mathbf{Z}^d$ and let $n \in N$, where N is the normaliser of G . Then by Remark 1, n induces a similarity $L \rightarrow Ln$. Remark 2 says that the lattice Ln is also G -invariant. Let $L' \in \pi(L)$ such that $\det(n) = [L : (L \cap Ln)] [Ln : (L \cap Ln)]^{-1}$ equals $[L' : L]^{-1}$. Then $[L' : (L' \cap Ln)] = [Ln : (L' \cap Ln)]$. So one may conclude that, if there is no other G -invariant lattice M in the layer of L' (i.e. with $[L' : (M \cap L')] = [M : (M \cap L')]$), then $Ln = L'$.*

The last uniqueness condition is fulfilled if $C_{M_d(\mathbf{Q})}(G) \cong \mathbf{Q}$, all lattices in $\mathcal{Z}(G)$ are even, and G is lattice sparse according to the following definition. In this case $\mathcal{Z}(G) = \{aL' \mid L' \in \pi(L), a \in \mathbf{Q}^*\}$ for any G -invariant lattice L . Note that this implies that the exponent $\exp(L^\# / L)$ is square free.

DEFINITION 8. *If p is a prime then a finite group $G \leq GL_d(\mathbf{Q})$ is called p -lattice sparse if any lattice $L \in \mathcal{Z}(G)$ can be obtained from any other lattice in $\mathcal{Z}(G)$ that contains L of p -power index by a combination of the five operations: taking sums, intersections, the dual lattice or the even sublattice with respect to some $F \in \mathcal{F}_{>0}(G)$, or multiplying by invertible elements of $C_{M_d(\mathbf{Q})}(G)$. The group G is called lattice sparse if G is p -lattice sparse for all primes p .*

Since the proof of the Theorem is similar for all r.i.m.f. groups, we only deal with the most interesting cases $d = 16$ and 24 .

The r.i.m.f. groups of degree 16 and 24 fixing strongly modular lattices, which are not proper tensor products, are displayed in the table. The first column gives the number of the group under which it is referred to in [NeP 95] or [Neb 95] and [Neb 96]. The second column contains a name for the matrix group as partially explained in the paragraph preceding Proposition 5. In the notation there we additional make the following abbreviations. If $z = d_1$ or d_2 , we omit \times and Q in the symbols. Also (1) is omitted if $p = 1$. The division algebra Q is abbreviated as α , if $Q = \mathbf{Q}[\alpha]$, by the set of ramified primes, if Q is a quaternion algebra over \mathbf{Q} , and omitted if $Q = \mathbf{Q}$. For the finite simple and quasisimple groups we use the notation of [CCNPW 85]

except that the alternating group is denoted by Alt_n to avoid confusion with A_n , which also denotes the automorphism group of the root lattice A_n .

The next three columns give information about the invariant strongly modular lattice L . First the determinant $\det(L)$ is given as the product of the abelian invariants of the Sylow subgroups of $L^\# / L$. The next column contains $\min(L)$ the minimum of the square lengths of the non zero vectors in L . From these two columns one may see whether L is an extremal lattice as defined in [Que 96]. The number of vectors of square length $\min(L)$ decomposed as a sum of the orbit lengths under $\text{Aut}(L)$ is displayed in the fifth column. The last column contains the primes p for which $\text{Aut}(L)$ is p -lattice sparse. A + indicates that $\text{Aut}(L)$ is lattice sparse.

Proof of the Theorem. The commuting algebras of the groups in the table are all isomorphic to \mathbf{Q} except for the one of $[6.\text{Alt}_7 : 2]_{24}$ which is $\mathbf{Q}[\sqrt{-6}]$. So all these groups are uniform. Since the arguments are similar for all groups G we only deal with $G = [SL_2(5) \underset{\infty, 3}{\boxtimes} SL_2(9)]_{16}^{2(3)}$ extensively. Let $L \in \mathcal{Z}(G)$ be a G -invariant lattice. There is a unique $F \in \mathcal{F}_{>0}(L)$ such that $\{l_1 Fl_2^{tr} \mid l_1, l_2 \in L\} = \mathbf{Z}$. The determinant of L with respect to F is $|L^\# / L| = 3^8 \cdot 5^8$ and its minimum is 10. If this lattice is strongly modular, then it is an extremal strongly modular lattice ([Que 96]).

Since G is of the form described in Proposition 5 with $p = 3$, there is an element $a_1 \in N = N_{GL_{16}(\mathbf{Q})}(G)$ with $\frac{1}{3}a_1^2 \in G$. Since G is lattice sparse and $\det(a_1) = \frac{1}{3}^8$ one has $La_1 = 3L^\# \cap L \in \mathcal{Z}(G)$ and $a_1Fa_1^{tr} = 3F$ (by Remarks 2 and 1). Hence a_1 induces a similarity between L and $3L^\# \cap L$.

Next consider the normal subgroup $U := SL_2(5) \underset{\infty, 3}{\boxtimes} SL_2(9) \trianglelefteq G$. The commuting algebra $C_{M_{16}(\mathbf{Q})}(U) =: K$ is isomorphic to $\mathbf{Q}[\sqrt{5}]$. From Proposition 4 one obtains an element $c \in N$ with $c^2 = 5$. As above one concludes that c yields a similarity between L and $5L^\# \cap L$. The product $a_1c \in N$ is of determinant $\pm 15^8$ and gives a similarity between L and $15L^\#$. Therefore L is strongly modular.

Most of the other groups can be dealt with similarly. One has to use Proposition 6 to construct an additional element of N for the r.i.m.f. groups 4 and 14 of $GL_{16}(\mathbf{Q})$. For $G = [2.\text{Alt}_{10}]_{16}$ (number 14), one obtains $n \in N$ of determinant $\pm 5^8$, since the character extends to $2.S_{10}$ with character field $\mathbf{Q}[\sqrt{\pm 5}]$ (cf. [CCNPW 85]). Analogously, for $F_4 \tilde{\otimes} F_4 = 2_+^{1+8} \cdot O_8^+(2)$ (number 4) the character extends to $2_+^{1+8} \cdot O_8^+(2).2$ with character field $\mathbf{Q}[\sqrt{\pm 2}]$.

The strong modularity for the lattices of the r.i.m.f. groups 9 and 21 of $GL_{16}(\mathbf{Q})$ (in particular the similarity of L with the lattice corresponding to

the Sylow-2-subgroup of $L^\# / L$) may be derived from the equality

$$[(Sp_4(3) \circ C_3) \overset{2}{\boxtimes}_{\sqrt{-3}} SL_2(3)]_{16} = [(Sp_4(3) \circ C_3) \overset{2(2)}{\boxtimes}_{\sqrt{-3}} SL_2(3)]_{16}$$

and

$$[SL_2(5) \overset{2(3)}{\boxtimes}_{\infty, 3} (SL_2(3) \overset{2}{\square} C_3)]_{16} = [(SL_2(5).2 \circ C_3) \overset{2(2)}{\boxtimes}_{\sqrt{-3}} SL_2(3)]_{16}$$

using Proposition 5.

Similarly one uses Proposition 5 to show the 2-modularity of the lattices of the r.i.m.f. group 6 in $GL_{24}(\mathbf{Q})$ using the description

$$[6.U_4(3).2 \overset{2}{\boxtimes}_{\sqrt{-3}} SL_2(3)]_{24} = [6.U_4(3).2 \overset{2(2)}{\circ} SL_2(3)]_{24}.$$

For the groups 44 and 64, which are the only groups which are not p -lattice sparse for a relevant prime p ($=2$), one has to note that the invariant sublattice of index 2^{12} in L is unique.

The theorem now follows from the next lemma. \square

LEMMA 9. *The lattices (of determinant $3^8 \cdot 5^8$) of the r.i.m.f. subgroup $G := [\pm \text{Alt}_6.2^2]_{16} \leq GL_{16}(\mathbf{Q})$ (number 20 of [NeP 95]) are not (strongly) modular.*

Proof. Let L be such a G -invariant lattice and $L' \in \pi(L)$. Assume that there is a similarity $s : L' \rightarrow L$. By Proposition 3, this similarity s normalises G . Let $U \cong \text{Alt}_6$ be the characteristic subgroup $\cong \text{Alt}_6$ of G . Since the full automorphism group of U is already induced by conjugation with elements of G , there exists $g \in G$, such that $n := gs \in GL_{16}(\mathbf{Q})$ centralises U . Hence $n \in C_{M_{16}(\mathbf{Q})}(U) \cong \mathbf{Q}[\sqrt{5}]$. Since this number field does not contain an element of norm 3, one concludes that $[L' : L] = 5^8$. So the lattice L is neither similar to $L^\#$ nor to the lattice $L' \in \pi(L)$ corresponding to the 3-Sylow subgroup of $L^\# / L$. Note that if $[L' : L] = 5^8$, an element $x \in C_{M_{16}(\mathbf{Q})}(U)$ with $x^2 = 5$, induces a similarity by Proposition 4. \square

REFERENCES

[CCNPW 85] CONWAY, J. H., R. T. CURTIS, S. P. NORTON, R. A. PARKER and R. A. WILSON. *Atlas of Finite Groups*. Oxford University Press, 1985.

[Neb 95] NEBE, G. Endliche rationale Matrixgruppen vom Grad 24. Dissertation RWTH Aachen. *Aachener Beiträge zur Mathematik* 12 (1995).