Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 43 (1997)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE LOCAL LINEARIZATION PROBLEM FOR SMOOTH SL(n) -

ACTIONS

Autor: CAIRNS, Grant / Ghys, Étienne

Kapitel: 4. SL(n,R)-ACTIONS ON \$R^n\$ FOR n 3

DOI: https://doi.org/10.5169/seals-63275

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 28.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

4. $SL(n, \mathbf{R})$ -actions on \mathbf{R}^n for $n \ge 3$

Let $n \geq 3$. We first give examples of C^0 -actions of $SL(n, \mathbf{R})$ on \mathbf{R}^n . Consider the canonical projective action of $SL(n, \mathbf{R})$ on S^{n-1} . Let Δ^+ be the radial half-line through the first basis element e_1 and let H denote the subgroup of $SL(n, \mathbf{R})$ that fixes Δ^+ . So $SL(n, \mathbf{R})/H \cong S^{n-1}$. Consider the homomorphism

$$\psi \colon (A_{ij}) \in H \mapsto \ln A_{11} \in \mathbf{R}$$
.

Notice that one obtains a linear action of H on $\mathbf{R}_*^+ = (0, \infty)$ by setting $h(x) = e^{\psi(h)}x$, for all $h \in H$, $x \in \mathbf{R}_*^+$. Obviously this is conjugate to the H-action on Δ^+ . It follows from Lemma 3.7 that the action of $SL(n, \mathbf{R})$ obtained by suspension of this action of H on \mathbf{R}_*^+ is the canonical linear action of $SL(n, \mathbf{R})$ on $\mathbf{R}^n \setminus \{0\}$. In fact, the map

$$\psi \colon [g.x] \in \left(SL(n,\mathbf{R}) \times \mathbf{R}_*^+\right)/H \mapsto g(x e_1) \in \mathbf{R}^n \setminus \{0\}$$

is an isomorphism. We now deform the action of H. Choose a topological flow $(\phi^I)_{I\in\mathbf{R}}$ on $\mathbf{R}^+=[0,\infty)$, fixing 0. This defines an action of H on \mathbf{R}^+_* by setting $h(x)=\phi^{\psi(h)}(x)$, for all $h\in H$, $x\in\mathbf{R}^+_*$. Now suspend this action of H and let Φ denote the resulting action of $SL(n,\mathbf{R})$ on the space $M=(SL(n,\mathbf{R})\times\mathbf{R}^+_*)/H$. The space M fibres over S^{n-1} , with fibre \mathbf{R}^+_* , and the structure group is orientation preserving. So topologically, M is $\mathbf{R}^+_*\times S^{n-1}$. Thus, identifying $S^{n-1}\times\{0\}$ to a point, we obtain an $SL(n,\mathbf{R})$ -action on \mathbf{R}^n . The fixed points of the flow ϕ correspond to orbits in \mathbf{R}^n which are spheres of dimension n-1. In general, an n-dimensional orbit is either all of $\mathbf{R}^n\setminus\{0\}$, as in the linear case, or it is a spherical shell, bounded by S^{n-1} orbits, or a punctured ball bounded by an S^{n-1} orbit, or the exterior of an S^{n-1} orbit. In all cases, the n-dimensional orbits are conjugate to the canonical linear one on $\mathbf{R}^n\setminus\{0\}$, by Theorem 3.5(c).

THEOREM 4.1. For all $n \geq 3$, every non-trivial C^0 -action of $SL(n, \mathbf{R})$ on $(\mathbf{R}^n, 0)$ is conjugate to one of the above actions Φ .

Proof. Suppose that we have a non-trivial C^0 -action of $SL(n, \mathbf{R})$ on $(\mathbf{R}^n, 0)$. First use Proposition 3.8 to linearize the SO(n)-action. Then by Lemma 3.9, the $SL(n, \mathbf{R})$ -action preserves the radial lines. Hence the radial projection $\mathbf{R}^n \setminus \{0\} \to S^{n-1}$ is equivariant, where the action of $SL(n, \mathbf{R})$ on S^{n-1} is the canonical projective one. Let H be the stabilizer of the radial half-line Δ^+ through e_1 , as above. So the action of $SL(n, \mathbf{R})$ on $\mathbf{R}^n \setminus \{0\}$ is induced by some action of H on \mathbf{R} . Notice that this action is trivial when restricted to

SO(n-1). It remains to consider all actions of H on \mathbb{R} which are trivial on SO(n-1). Again, by Lie [23, ibid.], these are given by homomorphisms from H to \mathbb{R} , Aff, or (some cover of) $PSL(2,\mathbb{R})$. We have the homomorphism $\psi\colon (A_{ij})\in H\mapsto \ln A_{11}\in \mathbb{R}$. Note that $\ker \psi=SL(n-1,\mathbb{R})\ltimes \mathbb{R}^{n-1}$. But it is easy to see that there are no non-trivial homomorphisms of $\ker \psi$ to \mathbb{R} or Aff. There are no non-trivial homomorphisms of $\ker \psi$ to $SL(2,\mathbb{R})$, except in the case n=3, and in this case there are no such homomorphisms which are trivial on SO(n-1). So the only possibility left is that H acts on \mathbb{R} by some flow. Finally, we put back the origin, as in the proof of Proposition 3.8. This completes the proof of the theorem.

We now prove Theorem 1.1 for $n \ge 3$.

THEOREM 4.2. For all $n \ge 3$ and $k = 1, ..., \infty$, every C^k -action of $SL(n, \mathbf{R})$ on $(\mathbf{R}^n, 0)$ is C^k -linearizable.

Proof. Let $n \ge 3$ and $k = 1, ..., \infty$ and suppose that we have a non-trivial C^k -action of $SL(n, \mathbf{R})$ on $(\mathbf{R}^n, 0)$. By Remark 3.4, we may assume that the differential of the action at the origin is either the identity or the map $g \mapsto (g^{-1})^t$. We will assume that it is the identity; the other possibility can be handled using the same argument.

Linearizing the SO(n)-action, using the Bochner-Cartan theorem, one may assume that the SO(n)-action is the canonical one. Then by Lemma 3.9, the $SL(n, \mathbf{R})$ -action preserves the radial lines. Let Δ denote the radial line through the first of the canonical basis elements, e_1 . Consider $H = \operatorname{Stab}_{SL(n,\mathbf{R})}(\Delta)$, as before. So, as we saw in the proof of Theorem 4.1, H defines a C^k -flow on Δ . This flow is hyperbolic, by the first paragraph. Hence by Theorem 2.5, this flow is linearizable by some local C^k -diffeomorphism f of $\Delta (\cong \mathbf{R})$. So, after conjugacy, we may assume that H acts linearly on Δ . Now define the local C^k -diffeomorphism F of \mathbf{R}^n by the formula:

(2)
$$F(x) = \begin{cases} \frac{f(||x||)}{||x||} x, & x \neq 0 \\ 0, & x = 0. \end{cases}$$

To see that F is of class C^k , the key point is to verify that f is a C^k odd function on \mathbf{R} . This follows easily from the fact that the flow on Δ commutes with $\operatorname{Stab}_{SO(n,\mathbf{R})}(\Delta)$, and the SO(n)-action is linear.

Now notice that F agrees with f on $\Delta^+ = \{t e_1 \in \Delta : t \geq 0\}$, and as F commutes with the SO(n)-action, the SO(n)-action is unchanged by conjugation by F. In particular, the SO(n)-action still commutes with dilations.

It follows that after conjugation by F, the $SL(n, \mathbf{R})$ -action commutes with dilations. Indeed, consider the conjugated $SL(n, \mathbf{R})$ -action. If $f \in SL(n, \mathbf{R})$, $x \in \mathbf{R}^n$ and $\lambda > 0$, then choose $a, b \in SO(n)$ such that $ax \in \Delta^+$ and $bf(\lambda x) \in \Delta^+$. Provided x is sufficiently close to 0, ax and $bf(\lambda x)$ will lie in the domain of f. Then $bfa^{-1} \in H$ and so

$$f(\lambda x) = b^{-1}bfa^{-1}a(\lambda x) = b^{-1}(bfg^{-1})\lambda a(x)$$

= $b^{-1}\lambda(bfa^{-1})a(x) = \lambda b^{-1}(bfa^{-1})a(x)$
= $\lambda f(x)$.

The proof of the theorem is then completed by the following well known result (cf. [17, Lemma 2.1.4]). \Box

LEMMA 4.3. Every C^1 map commuting with dilations is linear.

Proof. Suppose that f is a C^1 -diffeomorphism of \mathbb{R}^n which commutes with dilations. By comparing the differential of $\lambda \cdot f$ and $f \circ \lambda$ at x we have $\lambda df|_x = \lambda df|_{\lambda x}$, for each $\lambda > 0$ and every $x \in \mathbb{R}^n$. Hence $df|_x = df|_{\lambda x}$ and so df is constant on the radial lines. Thus $df|_x = df|_0$ for all x and so f is linear. \square

5. The adjoint representation of $SL(2, \mathbf{R})$

Let us recall some facts concerning the linear representations of $SL(2, \mathbf{R})$. Let $P_l(\mathbf{R}^2)$ denote the space of real valued homogeneous polynomials, of two variables, of degree l. As a vector space, $P_l(\mathbf{R}^2) \cong \mathbf{R}^{l+1}$, and the action of $SL(2,\mathbf{R})$ on \mathbf{R}^2 defines a linear action on $P_l(\mathbf{R}^2)$: up to isomorphism, this is the (unique) irreducible representation of $SL(2,\mathbf{R})$ in dimension l+1. In dimension 3, there is another useful realization of the polynomial representation, called the *adjoint representation*. Notice that the group $SL(2,\mathbf{R})$ acts by the adjoint representation on its Lie algebra $\mathfrak{sl}(2,\mathbf{R})$. Of course, $\mathfrak{sl}(2,\mathbf{R})$ is the space of 2×2 real traceless matrices; so as a vector space, $\mathfrak{sl}(2,\mathbf{R})\cong \mathbf{R}^3$. The adjoint representation $Ad:SL(2,\mathbf{R})\to GL(3,\mathbf{R})$, defined by

$$Ad(g): h \mapsto ghg^{-1}, \quad \forall g \in SL(2, \mathbf{R}), \ h \in \mathfrak{sl}(2, \mathbf{R}),$$

is an irreducible linear representation. In fact, an explicit equivariant isomorphism $\psi \colon \mathfrak{sl}(2, \mathbf{R}) \to P_2(\mathbf{R}^2)$ is obtained by taking $\psi(h)$, as a function of variables x and y, to be the area of the parallelogram spanned by (x, y) and h(x, y). That is,