Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 44 (1998)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN ASYMPTOTIC FREIHEITSSATZ FOR FINITELY GENERATED GROUPS

Autor: Cherix, Pierre-Alain / SCHAEFFER, Gilles

Kapitel: 3. About genericity

DOI: https://doi.org/10.5169/seals-63893

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 15.01.2026

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

3. About genericity

LEMMA 3.1. *Let* $X = \{x_1, \ldots, x_k, y\}$. *For every* $0 < \epsilon < 1/(k+1)$, *the ratio*

$$\frac{\#\{r \in \mathbf{F}_X \mid |r| = n, r \text{ is } (\epsilon, y)\text{-balanced}\}}{\#\{r \in \mathbf{F}_X \mid |r| = n\}}$$

tends to 1 *when* n *tends* ∞.

Proof. First we want to rephrase the Lemma in terms of generating functions. Let K be any fixed subset of $\mathbf{F}_X$ and $F_K(z,u)$ be the generating function defined by

$$F_K(z,u) = \sum_{r \in K} z^{|r|} u^{n_y(r)}.$$

$F_K(z,u)$ strongly depends on the choice of the generator y. However, as y is fixed throughout the proof and to lighten the notation, we write $F_K(z,u)$ instead of $F_{y,K}(z,u)$.

Defining $c_{n,l}$ and $p_n(l)$ by

$$F_{\mathbf{F}_X}(z,u) = \sum_{r \in \mathbf{F}_X} z^{|r|} u^{n_y(r)} = \sum_{n,l} c_{n,l} z^n u^l \quad \text{and} \quad p_n(l) = \frac{c_{n,l}}{\sum_m c_{n,m}},$$

we have to prove that for every $0 < \epsilon < 1/(k+1)$,

$$\lim_{n \to \infty} \sum_{0 \le l < \epsilon n} p_n(l) = 0.$$

We want to find an analytical form for $F_{\mathbf{F}_X}(z,u)$.

It is clear that if K_1 and K_2 are disjoint subsets of $\mathbf{F}_X$ then $F_{K_1 \cup K_2}(z,u) = F_{K_1}(z,u) + F_{K_2}(z,u)$.

Let K_1, K_2 be two subsets of F_X; assume that the map $K_1 \times K_2 \to K_1K_2$ defined by $(\omega_1, \omega_2) \mapsto \omega_1\omega_2$ is one to one and satisfies $|\omega_1\omega_2| = |\omega_1| + |\omega_2|$ for $\omega_i \in K_i$ (where $K_1K_2 = \{\omega_1\omega_2 | \omega_i \in K_i\}$); it is also clear that $F_{K_1K_2}(z,u) = F_{K_1}(z,u)F_{K_2}(z,u)$. This can be extended to a finite product of such K_i's.

First we compute the generating functions of some subsets K of $\mathbf{F}_X$.

- $F_{\{e\}}(z,u) = 1$.
- Denote by $X' = X - \{y\}$. As there are exactly $2k(2k-1)^{n-1}$ reduced words of length $n \ge 1$ in $\mathbf{F}_{X'}$, we obtain $F_{[\mathbf{F}_{X'} - \{e\}]}(z,u) = \frac{2kz}{1-(2k-1)z}$. Set $f(z,u) = F_{[\mathbf{F}_{X'} - \{e\}]}(z,u)$.
- For $\langle y \rangle = \{y^i \mid i \in \mathbf{Z} - \{0\}\}$, we have $F_{\langle y \rangle}(z,u) = \frac{2uz}{1-uz}$, because there are exactly 2 elements $y^{\pm i}$ in $\langle y \rangle$ such that $n_y(y^{\pm i}) = |y^{\pm i}| = i$. Set $h(z,u) = F_{\langle y \rangle}(z,u)$.

Now we can partition $\mathbf{F}_X$ as follows:

$$\mathbf{F}_X = \{e\} \amalg [\mathbf{F}_{X'} - \{e\}] \coprod_{n\geq 0} I_n$$

where

$$I_n = \{\omega_0 y^{i_1} \omega_1 \dots y^{i_{n-1}} \omega_{n-1} y^{i_n} \omega_n \mid \omega_j \in \mathbf{F}_{X'},\ \omega_j \neq e \text{ for } j \neq 0 \text{ or } n, \text{ and } i_j \neq 0\}.$$

It is easy to check that $F_{I_n}(z,u) = (f(z,u)+1)^2 h(z,u)(h(z,u)f(z,u))^{n-1}$. So we obtain that

$$\begin{aligned} F_{\mathbf{F}_X}(z,u) &= 1 + f(z,u) + \sum_{n\geq 1} (f(z,u)+1)^2 h(z,u)(h(z,u)f(z,u))^{n-1} \\ &= (1+f(z,u))\Big(1 + \frac{h(z,u)(f(z,u)+1)}{1-h(z,u)f(z,u)}\Big) \\ &= \frac{(1+z)(1+uz)}{1-(2k-1)z-uz(1+(2k+1)z)}. \end{aligned}$$

Borrowing notation from [2], let $g(z,u) = (1+z)(1+zu)$ and $P(z,u) = 1-(2k-1)z-uz(1+(2k+1)z) = 1-(2k-1+u)z-(2k+1)uz^2$. Then

$$F_{\mathbf{F}_X}(z,u) = \frac{g(z,u)}{P(z,u)}$$

and let $r(s)$ be the root of smallest modulus of $P(r(s), e^s) = 0$ in a small neighborhood of $s = 0$. In particular $r(0) = \frac{1}{2k+1}$. According to [2, (3.1)], we obtain from [2, Theorem 1] that

$$\lim_{n\to\infty} \sup_x \Big| \sum_{k\leq \sigma_n x + \mu_n} p_n(k) - \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-t^2/2} dt \Big| = 0$$

with $\mu = \frac{r'(0)}{r(0)}$, $\mu_n = n\mu = n\frac{r'(0)}{r(0)}$ and $\sigma_n^2 = n\sigma^2 = n(\mu^2 - \frac{r''(0)}{r(0)})$.

Computing $r'(0)$ or easy combinatorial considerations gives $\mu_n = \frac{n}{k+1}$. The actual value of σ is here useless.

Now let $\epsilon < \frac{1}{k+1}$ and $\delta > 0$. Let x such that $\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{x} e^{-t^2/2} dt < \delta$. Then there exists N such that for $n > N$, $\epsilon n < \sigma\sqrt{n}x + \frac{n}{k+1}$ since $\epsilon < \frac{1}{k+1}$. Therefore, for $n > N$,

$$\sum_{k<\epsilon n} p_n(k) \leq \sum_{k<\sigma_n x + \mu_n} p_n(k)$$

and there exists N_1 such that for $n > N_1$,

$$\sum_{k<\epsilon n} p_n(k) \leq 2\delta. \qquad \square$$

COROLLARY 3.2. *For* $\#X = k$, $\#R = n$, $x_0 \in X$ *and* $0 < \epsilon < 1/k$ *fixed, being* (ϵ, x_0)*-balanced is generic for* $\Gamma = \langle X \,|\, R \rangle$.

Proof of corollary. We choose n relations at random; by Lemma 3.1, every $r \in R$ is generically (ϵ, x_0)-balanced, but the conjunction of finitely many generic properties is also generic. □

4. SOME SUFFICIENT CONDITIONS FOR THE EXISTENCE OF FREE SUBGROUPS

We first begin by a very easy proposition.

PROPOSITION 4.1. *Let* $\Gamma = \langle X \mid R \rangle$ *be a finite presentation, which has a Dehn algorithm and such that for some* $y \in X$ *every subword* u *of every* $r \in R^*$ *with* $|u| > |r|/2$ *contains either* y *or* y^{-1}, *then* $X - \{y\}$ *generates a free subgroup in* Γ.

The proof of this proposition will follow from Lemma 4.2 below.

LEMMA 4.2. *For* $\langle X \,|\, R \rangle$ *a finite presentation of a group* Γ *and* $y \in X$, *the following are equivalent:*

- $X - \{y\}$ *freely generates a free subgroup of* Γ;
- *every non trivial element* $\omega \in \mathbf{F}_X$, *which represents the identity in* Γ, *contains either* y *or* y^{-1}.

Proof. 1) $\Rightarrow$ 2) : By contraposition, suppose that there exists a non trivial reduced element $\omega \in \mathbf{F}_{X-\{y\}}$ such that $\overline{\omega} = e$ (where $\overline{\omega}$ is the canonical projection of ω in Γ), then $X - \{y\}$ does not freely generate a free subgroup in Γ.

2) $\Rightarrow$ 1) : Let $\omega_1, \omega_2 \in \mathbf{F}_{X-\{y\}}$ be two reduced elements such that $\overline{\omega}_1 = \overline{\omega}_2 \in \Gamma$. Then $\overline{\omega_1 \omega_2^{-1}} = e \in \Gamma$. So $\omega_1 \omega_2^{-1}$ is an element of $\mathbf{F}_{X-\{y\}}$ which represents the identity in Γ. By hypothesis, this implies $\omega_1 = \omega_2$ in $\mathbf{F}_X$. Hence $X - \{y\}$ freely generates a free subgroup in Γ. □

Proof of Proposition 4.1. By Lemma 4.2, it is sufficient to show that every non trivial reduced word on $\mathbf{F}_X$ which represents the identity in Γ contains either y or y^{-1}. By assumption, $\Gamma = \langle X \,|\, R \rangle$ satisfies a Dehn algorithm, so such a word contains at least one half of a relator r in R which contains at least one occurrence of y or y^{-1}. □