Zeitschrift: L'Enseignement Mathématique

Herausgeber: Commission Internationale de l'Enseignement Mathématique

Band: 44 (1998)

Heft: 1-2: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: AN ASYMPTOTIC FREIHEITSSATZ FOR FINITELY GENERATED

GROUPS

Autor: Cherix, Pierre-Alain / SCHAEFFER, Gilles

Kapitel: 4. SOME SUFFICIENT CONDITIONS FOR THE EXISTENCE OF

FREE SUBGROUPS

DOI: https://doi.org/10.5169/seals-63893

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. Mehr erfahren

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. En savoir plus

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. Find out more

Download PDF: 29.11.2025

ETH-Bibliothek Zürich, E-Periodica, https://www.e-periodica.ch

COROLLARY 3.2. For #X = k, #R = n, $x_0 \in X$ and $0 < \epsilon < 1/k$ fixed, being (ϵ, x_0) -balanced is generic for $\Gamma = \langle X | R \rangle$.

Proof of corollary. We choose n relations at random; by Lemma 3.1, every $r \in R$ is generically (ϵ, x_0) -balanced, but the conjunction of finitely many generic properties is also generic.

4. Some sufficient conditions for the existence of free subgroups

We first begin by a very easy proposition.

PROPOSITION 4.1. Let $\Gamma = \langle X | R \rangle$ be a finite presentation, which has a Dehn algorithm and such that for some $y \in X$ every subword u of every $r \in R^*$ with |u| > |r|/2 contains either y or y^{-1} , then $X - \{y\}$ generates a free subgroup in Γ .

The proof of this proposition will follow from Lemma 4.2 below.

LEMMA 4.2. For $\langle X | R \rangle$ a finite presentation of a group Γ and $y \in X$, the following are equivalent:

- $X \{y\}$ freely generates a free subgroup of Γ ;
- every non trivial element $\omega \in \mathbf{F}_X$, which represents the identity in Γ , contains either y or y^{-1} .

Proof. 1) \Rightarrow 2): By contraposition, suppose that there exists a non trivial reduced element $\omega \in \mathbf{F}_{X-\{y\}}$ such that $\overline{\omega} = e$ (where $\overline{\omega}$ is the canonical projection of ω in Γ), then $X - \{y\}$ does not freely generate a free subgroup in Γ .

2) \Rightarrow 1): Let $\underline{\omega_1, \omega_2} \in \mathbf{F}_{X-\{y\}}$ be two reduced elements such that $\overline{\omega}_1 = \overline{\omega}_2 \in \Gamma$. Then $\overline{\omega_1 \omega_2^{-1}} = e \in \Gamma$. So $\omega_1 \omega_2^{-1}$ is an element of $\mathbf{F}_{X-\{y\}}$ which represents the identity in Γ . By hypothesis, this implies $\omega_1 = \omega_2$ in \mathbf{F}_X . Hence $X - \{y\}$ freely generates a free subgroup in Γ .

Proof of Proposition 4.1. By Lemma 4.2, it is sufficient to show that every non trivial reduced word on \mathbf{F}_X which represents the identity in Γ contains either y or y^{-1} . By assumption, $\Gamma = \langle X | R \rangle$ satisfies a Dehn algorithm, so such a word contains at least one half of a relator r in R which contains at least one occurrence of y or y^{-1} . \square

The interest of this proposition appears when we replace "having a Dehn's algorithm" by "satisfying the small cancellation condition C'(1/6)", because C'(1/6) and the fact that every subword u of any relation r with |u| > |r|/2 contains at least one y or y^{-1} are easy to check on a given presentation.

Unfortunately, as explained before, it is not known if the small cancellation hypothesis is generic, so we need other sufficient conditions to ensure that $X - \{y\}$ generates a free subgroup in Γ .

PROPOSITION 4.3. Let $\Gamma = \langle X \mid R \rangle$ be a finite presentation with k generators and l relations, which is (ϵ, x_0) -balanced for some $0 < \epsilon < 1/k$ and some $x_0 \in X$, and which satisfies a θ -condition such that $\theta < \epsilon/(2 - \epsilon)$. Then $X - \{x_0\}$ freely generates a free group in Γ .

To prove the proposition we need the following lemma and the following notations. For a cell f_i of the diagram, we denote by $Int(f_i)$ (resp. by $Ext(f_i)$) the number of edges of f_i which are internal to the diagram (resp. which are on the border of the diagram). We denote also by $\#(f_i)$ the total number of edges of the cell f_i .

LEMMA 4.4. Let $\Gamma = \langle X | R \rangle$ be a finite presentation of a group Γ which satisfies a θ -condition for some $0 < \theta < 1$, then for every reduced diagram, there exists a 2-cell f of Δ satisfying

$$Int(f) \leq \frac{2\theta}{1+\theta} \#(f)$$
.

Proof. First we prove it for simple diagrams. Let $\epsilon = 2\theta/(1+\theta)$. Because the diagram is simple we have the following equalities:

I)
$$\sum_{i} Ext(f_i) = E(\Delta) = |\partial \Delta|,$$

II) $\sum_{i} Int(f_i) = 2I(\Delta)$, because every internal edge belongs to two different cells.

So we get:

$$\#(\Delta) = \frac{1}{2} \sum_{i} Int(f_i) + \sum_{i} Ext(f_i) = \sum_{i} \#(f_i) - \frac{1}{2} \sum_{i} Int(f_i)$$
.

To obtain a contradiction, we suppose that every cell f_i of one diagram Δ is such that $(1/\epsilon)Int(f_i) > \#(f_i)$. Then we have

$$\frac{1}{\epsilon} \sum_{i} Int(f_i) > \sum_{i} \#(f_i) = \#(\Delta) + \frac{1}{2} \sum_{i} Int(f_i),$$

whence $\frac{2-\epsilon}{2\epsilon} \sum_{i} Int(f_i) > \#(\Delta)$ or $\frac{2-\epsilon}{\epsilon} I(\Delta) > \#(\Delta)$. Since $\epsilon = 2\theta/(1+\theta)$, we obtain $I(\Delta) > \theta \#(\Delta)$, which contradicts the θ -condition.

In fact, if the reduced diagram Δ is not simple, it is a union of simple diagrams linked by bridges. So each of its parts, which is a simple diagram, defines another reduced diagram (relative to another word), so the inequality holds for every part of Δ which is a simple diagram. We conclude by saying that increasing the number of external edges does not affect the inequality. \square

Proof of 4.3. By Lemma 4.2, it is sufficient to prove that the (ϵ, x_0) -balanced and θ -conditions imply that every non trivial reduced word in \mathbf{F}_X which vanishes in Γ contains at least one $x_0^{\pm 1}$.

Let us choose such a word ω and Δ a reduced diagram of ω . By Lemma 4.4, there exists a cell f with border equal to one $r \in R^*$, such that

$$Int(f) \leq \frac{2\theta}{1+\theta} \#(f) = \frac{2\theta}{1+\theta} |r| < \epsilon |r| \leq n_{x_0}(r),$$

because $\theta < \epsilon/(2-\epsilon)$. As there are more occurrences of x_0 or x_0^{-1} than the number of internal edges, it means that some occurrences of x_0 or x_0^{-1} will be external edges, i.e. will be in the border of Δ which is ω .

We are now able to prove the main theorem.

Proof of theorem 1.1. By Proposition 4.3, for a finite presentation $\langle X|R\rangle$, we know that being (ϵ, x_0) -balanced and satisfying a θ -condition is sufficient to ensure that $X - \{x_0\}$ freely generates a free subgroup in Γ . But by Corollary 3.2 and [13, Theorem 2], these two conditions are generic and so is the conjunction of these two conditions. \square

5. SPECTRAL ESTIMATES FOR ADJACENCY OPERATORS ON CAYLEY GRAPHS

The existence of a free subgroup generated by $X - \{x_0\}$ gives an upper bound for the spectral value of the adjacency operator on the Cayley graph of $\Gamma = \langle X | R \rangle$ associated with the symmetric generating system $S = X \cup X^{-1}$.

We briefly recall some definitions and notations. The Cayley graph $G(\Gamma, X)$ of Γ associated with S has its set of vertices in bijection with Γ and two