

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 44 (1998)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: HOMFLY POLYNOMIAL VIA AN INVARIANT OF COLORED PLANE GRAPHS
Autor: Murakami, Hitoshi / Yamada, Shuji
DOI: <https://doi.org/10.5169/seals-63908>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

HOMFLY POLYNOMIAL VIA AN INVARIANT OF COLORED PLANE GRAPHS

by Hitoshi MURAKAMI, Tomotada OHTSUKI and Shuji YAMADA

ABSTRACT. After the first discovery of the quantum invariant associated with $SU(2)$ by V.F.R. Jones [3], the invariants associated with $SU(n)$ were found by several authors [1]. It was first proved by V.G. Turaev [16] that all these come from so-called “quantum groups”, especially from their R -matrices corresponding to the vector representations. There also exist various quantum invariants corresponding to other representations (see for example [7], [14], [11]).

The aim of this paper is to give a graphical way to define $SU(n)$ quantum invariants for links. To do this we first construct an invariant of colored, oriented, trivalent, plane graphs for each n (≥ 2). Then we show that the $SU(n)$ polynomial invariant corresponding to the vector representation (HOMFLY polynomial) can be defined by using our graph invariant.

We can also show that our invariant defines the $SU(n)$ polynomial invariant corresponding to the anti-symmetric tensors of the vector representation.

We note that our graph invariant for $SU(3)$ was first introduced by G. Kuperberg in [8]. The second and the third authors used it in [12] to construct magic elements and defined the quantum $SU(3)$ invariants for 3-manifolds. Now [12] and the present paper together give an elementary and self-contained proof of the existence of magic elements for $SU(3)$ and so that of the quantum $SU(3)$ invariants of 3-manifolds just like W.B.R. Lickorish did for $SU(2)$ in [9] using the Kauffman bracket [5]. See [17] for a similar approach to $SU(n)$ invariants of 3-manifolds.

We also note that our graph invariant may be obtained (not checked yet) by direct computations of the universal R -matrix. But the advantage of our definition is that it does not require any knowledge of quantum groups nor representation theory. On the contrary we can recover the R -matrix of the quantum group $U_q(\mathfrak{sl}(n, \mathbb{C}))$ corresponding at least to the vector representation.

This work was inspired after conversation with M. Kosuda and J. Murakami, to whom the authors express their gratitude.

The first named author was in the Department of Mathematics, Osaka City University when the work was carried out and partially supported by Grant-in-Aid for Scientific Research on Priority Areas 231 “Infinite Analysis”, the Ministry of Education, Science, Sports and Culture. The second named author was in the Department of Mathematical Sciences, University of Tokyo.

§ 1. INVARIANTS OF GRAPHS

In this section we define an invariant of colored, oriented, trivalent, plane graphs.

Fix an integer $n \geq 2$ throughout this paper and put

$$\mathcal{N} = \{-(n-1)/2, -(n-1)/2 + 1, \dots, (n-1)/2\}.$$

For disjoint subsets A_1 and A_2 of \mathcal{N} we put

$$\pi(A_1, A_2) = \#\{(a_1, a_2) \in A_1 \times A_2 \mid a_1 > a_2\}.$$

Let G be an oriented, trivalent, plane graph with “color” or “flow” on each of its edges. Here a *flow* f is a map from the edge set to positive integers less than or equal to n such that for every vertex v in G the sum of its values on the edges coming into v is equal to that on the edges going out from v (see Figure 1.1). So we may say that G is a network with infinite capacity without source or sink. We also note that at each vertex two edges are “in” and one edge is “out”, or two edges are “out” and one edge is “in”. We call these two in- or out-edges the *legs* and one out- or in-edge the *head* of the vertex.




FIGURE 1.1

A graph with flow

A state σ is an assignment of a subset A of \mathcal{N} to each edge e such that $\#(A) = f(e)$ and, moreover, at each vertex the union of subsets assigned to its legs coincides with that assigned to its head, where $\#(A)$ is the number of elements in A (see Figure 1.2). We denote by $\sigma(e)$ the subset of \mathcal{N} assigned to an edge e .




FIGURE 1.2

A state of the graph with flow in Figure 1.1

Given a state σ , we define the *weight* $\text{wt}(v; \sigma)$ of a vertex v to be

$$q^{f(e_1)f(e_2)/4 - \pi(\sigma(e_1), \sigma(e_2))/2},$$

where q is an indeterminate, and e_1 and e_2 are left and right legs respectively with respect to the orientation of G (Figure 1.3).




FIGURE 1.3

If we replace every edge e with $f(e)$ copies of parallel edges, assign each copy an element of the subset determined by σ , and connect at every vertex each pair of edges with the same element, we have a union of simple closed curves each of which equipped with the same element of \mathcal{N} (Figure 1.4).




FIGURE 1.4

Simple closed curves defined by Figure 1.2

Then we define the *rotation number* $\text{rot}(\sigma)$ to be

$$\sum_C \sigma(C) \text{rot}(C)$$

where the sum is over all simple closed curves C equipped with $\sigma(C) \in \mathcal{N}$ and $\text{rot}(C)$ is the rotation number of C (i.e., 1 if C is counter-clockwise and -1 otherwise). For example $\text{rot}(\sigma) = 2$ for the state described in Figure 1.2 (see Figure 1.4).

Now we define $\langle G \rangle_n$ as follows.

$$\langle G \rangle_n = \sum_{\sigma: \text{state}} \left\{ \prod_{v: \text{vertex}} \text{wt}(v; \sigma) \right\} q^{\text{rot}(\sigma)}.$$

We define $\langle \text{empty graph} \rangle_n = 1$. It is clear that this is invariant under ambient isotopy of \mathbf{R}^2 . Note that our invariant can be regarded as a colored graph invariant introduced by N.Yu. Reshetikhin and V.G. Turaev in [14] replacing each vertex by a “coupon”. The coupon with two legs in would correspond to a projection $V_i \otimes V_j \rightarrow V_{i+j}$ and that with two legs out to an inclusion $V_{i+j} \rightarrow V_i \otimes V_j$, where V_i is the irreducible representation of $SU(n)$ corresponding to the i -fold anti-symmetric tensor of the vector representation.

§2. LOCAL PROPERTIES OF $\langle G \rangle_n$

We will describe some local properties of $\langle G \rangle_n$. In what follows diagrams indicated in each equality are identical outside the angle brackets $\langle \rangle_n$ and each equality also holds if we reverse all the orientations of diagrams in both hand sides. A number near an edge indicates its flow. If a flow in a diagram exceeds n , we disregard the term where the diagram appears.

We put

$$[k] = \frac{q^{k/2} - q^{-k/2}}{q^{1/2} - q^{-1/2}},$$

$$[k]! = [1][2] \cdots [k],$$

and

$$\begin{bmatrix} i \\ j \end{bmatrix} = \frac{[i]!}{[j]![i-j]}.$$

In the following equations we mean that if we replace the graph appearing in the left hand side with the one in the right hand side, we obtain the same value.

LEMMA 2.1.

$$\left\langle 1 \bigcirc \right\rangle_n = [n] \langle \emptyset \rangle_n.$$

Proof. From the definition the left hand side is equal to

$$\sum_{i=-(n-1)/2}^{(n-1)/2} q^i,$$

which is $[n]$, completing the proof. \square

LEMMA 2.2.

$$(2.1) \quad \left\langle 1 \begin{array}{c} \uparrow 2 \\ \downarrow 2 \end{array} 1 \right\rangle_n = [2] \left\langle \begin{array}{c} \uparrow 2 \\ \downarrow 2 \end{array} \right\rangle_n.$$

Proof. Consider a state σ of the left hand side. First note that the top-most and the bottom-most edges are equipped with the same subset. So we may put it $\{\alpha, \beta\}$ ($\alpha < \beta$). Then there are two cases; (i) the left edge is equipped with $\{\alpha\}$ and the right one with $\{\beta\}$ and (ii) the left edge is equipped with $\{\beta\}$ and the right one with $\{\alpha\}$. In the first case the weights of the upper and lower vertex are the same and equal to $q^{1/4}$. In the second case they are also the same and equal to $q^{-1/4}$. Therefore the contribution of the two vertices is

$$q^{1/2} + q^{-1/2} = [2],$$

which is independent of σ and the conclusion follows. \square

LEMMA 2.3.

$$\left\langle \begin{array}{c} 1 \uparrow \\ 2 \uparrow \\ 1 \uparrow \end{array} \bigcirc 1 \right\rangle_n = [n-1] \left\langle \begin{array}{c} \uparrow 1 \\ \downarrow 1 \end{array} \right\rangle_n.$$

Proof. In this case the top-most and the bottom-most edges of the left hand side are equipped with the same subset for any state σ as the previous lemma. So we put it $\{\alpha\}$. If the right edge is equipped with $\{\beta\}$ ($\beta \neq \alpha$), then the weights of the two vertices are the same and equal to $q^{\text{sign}(\beta-\alpha)/4}$.

Since its contribution to the rotation number is $-\beta$, the contribution of the left hand side is

$$\begin{aligned} \sum_{\beta \neq \alpha} q^{\text{sign}(\beta-\alpha)/2-\beta} &= \sum_{\beta=-(n-1)/2}^{\alpha-1} q^{-1/2-\beta} + \sum_{\beta=\alpha+1}^{(n-1)/2} q^{1/2-\beta} \\ &= q^{(n-2)/2} + q^{(n-4)/2} + \dots + q^{-\alpha+1/2} \\ &\quad + q^{-\alpha-1/2} + \dots + q^{-(n-2)/2} = [n-1] \end{aligned}$$

and the proof is complete. \square

LEMMA 2.4.

$$\left\langle \begin{array}{c} \text{Square with loops 1 and edges 1, 2} \end{array} \right\rangle_n = [n-2] \left\langle \begin{array}{c} \text{Two parallel edges 1 with arrows} \end{array} \right\rangle_n + \left\langle \begin{array}{c} \text{Vertical edge 1 with arrow} \end{array} \right\rangle_n$$

Proof. There are three possibilities:

- (1) the two edges at the top are equipped with $\{\alpha\}$ and the two edges at the bottom are equipped with $\{\beta\}$ ($\alpha \neq \beta$);
 - (2) both the top left and the bottom left edges are equipped with $\{\alpha\}$ and both the top right and the bottom right edges are equipped with $\{\beta\}$ ($\alpha \neq \beta$);
- and
- (3) all the four edges at the corners are equipped with $\{\alpha\}$.

In the first case, the two horizontal edges are equipped with $\{\gamma\}$ ($\alpha \neq \gamma \neq \beta$). So the contribution of the left hand side is

$$\left(\sum_{\substack{\gamma \\ \alpha \neq \gamma \neq \beta}} q^{\text{sign}(\alpha-\gamma)/2+\text{sign}(\beta-\gamma)/2} q^{\gamma} \right) q^{-(\alpha+\beta)/2} = [n-2] q^{-(\alpha+\beta)/2}$$

which is equal to the contribution of the right hand side.

In the second case, the contribution of the first term of the right hand side is zero. It is easy to see that the left hand side and the second term of the right hand side coincide.

In the third case, the contribution of the left hand side is $[n-2] q^{-\alpha} + 1$, which is equal to that of the right hand side. This completes the proof. \square

LEMMA 2.5.

Proof. There are three possibilities:

- (1) both the top left and the bottom left edges are equipped with $\{\alpha\}$ and both the top right and the bottom right edges are equipped with $\{\beta, \gamma\}$ (α, β and γ are all distinct);
 - (2) the top left and the bottom left edges are equipped with $\{\alpha\}$ and $\{\beta\}$ respectively and the top right and the bottom right edges are equipped with $\{\beta, \gamma\}$ and $\{\alpha, \gamma\}$ respectively (α, β and γ are all distinct);
- and
- (3) both the top left and the bottom left edges are equipped with $\{\alpha\}$ and both the top right and the bottom right edges are equipped with $\{\alpha, \beta\}$ ($\alpha \neq \beta$).

In the first case both the upper and the lower horizontal edges in the left hand side are equipped with $\{\beta\}$ or both of them are equipped with $\{\gamma\}$. So the contribution of the left hand side is $q^{\text{sign}(\beta-\alpha)/2+\text{sign}(\gamma-\beta)/2} + q^{\text{sign}(\gamma-\alpha)/2+\text{sign}(\beta-\gamma)/2}$. On the other hand that of the right hand side is $q^{1-\pi(\{\alpha\}, \{\beta, \gamma\})} + 1$. It can be easily checked that these are the same.

The second and the third cases are easily checked and left to the reader. \square

LEMMA 2.6.

Proof. This follows from the fact that

$$\begin{aligned}
 & \#\{(a_1, a_2) \in A_1 \times A_2 \mid a_1 > a_2\} + \#\{(a, a_3) \in (A_1 \cup A_2) \times A_3 \mid a > a_3\} \\
 &= \#\{(a_1, a_2) \in A_1 \times A_2 \mid a_1 > a_2\} \\
 &\quad + \#\{(a_1, a_3) \in A_1 \times A_3 \mid a_1 > a_3\} + \#\{(a_2, a_3) \in A_2 \times A_3 \mid a_2 > a_3\} \\
 &= \#\{(a_1, a) \in A_1 \times (A_2 \cup A_3) \mid a_1 > a\} + \#\{(a_2, a_3) \in A_2 \times A_3 \mid a_2 > a_3\}.
 \end{aligned}$$

\square

§3. INVARIANTS OF LINKS

In this section, we give a graphical definition of the HOMFLY polynomial of oriented links. For an oriented link diagram D , we define $\langle D \rangle_n$ by the following.

$$\left\langle \begin{array}{c} \nearrow \\ \searrow \\ \nearrow \\ \searrow \end{array} \right\rangle_n = q^{1/2} \left\langle \begin{array}{c} \uparrow \\ \uparrow \end{array} \right\rangle_n - \left\langle \begin{array}{c} \nearrow \quad \nearrow \\ \searrow \quad \searrow \\ \nearrow \quad \nearrow \\ \searrow \quad \searrow \end{array} \right\rangle_n$$

and

$$\left\langle \begin{array}{c} \nearrow \\ \searrow \\ \searrow \\ \nearrow \end{array} \right\rangle_n = q^{-1/2} \left\langle \begin{array}{c} \uparrow \\ \uparrow \end{array} \right\rangle_n - \left\langle \begin{array}{c} \nearrow \quad \nearrow \\ \searrow \quad \searrow \\ \searrow \quad \searrow \\ \nearrow \quad \nearrow \end{array} \right\rangle_n.$$

Then we have

THEOREM 3.1. $\langle D \rangle_n$ is invariant under the Reidemeister moves II and III.

Proof. From Lemma 2.2, we have

$$\begin{aligned} \left\langle \begin{array}{c} \nearrow \\ \searrow \\ \searrow \\ \nearrow \end{array} \right\rangle_n &= \left\langle \begin{array}{c} \uparrow \\ \uparrow \end{array} \right\rangle_n - q^{1/2} \left\langle \begin{array}{c} \nearrow \quad \nearrow \\ \searrow \quad \searrow \\ \nearrow \quad \nearrow \\ \searrow \quad \searrow \end{array} \right\rangle_n \\ &\quad - q^{-1/2} \left\langle \begin{array}{c} \nearrow \quad \nearrow \\ \searrow \quad \searrow \\ \searrow \quad \searrow \\ \nearrow \quad \nearrow \end{array} \right\rangle_n + \left\langle \begin{array}{c} \nearrow \quad \nearrow \\ \searrow \quad \searrow \\ \nearrow \quad \nearrow \\ \searrow \quad \searrow \end{array} \right\rangle_n \\ &= \left\langle \begin{array}{c} \uparrow \\ \uparrow \end{array} \right\rangle_n + (-q^{1/2} - q^{-1/2} + [2]) \left\langle \begin{array}{c} \nearrow \quad \nearrow \\ \searrow \quad \searrow \\ \nearrow \quad \nearrow \\ \searrow \quad \searrow \end{array} \right\rangle_n = \left\langle \begin{array}{c} \uparrow \\ \uparrow \end{array} \right\rangle_n. \end{aligned}$$

We also have

$$\begin{aligned}
 \left\langle \begin{array}{c} \text{Diagram 1} \end{array} \right\rangle_n &= \left\langle \begin{array}{c} \text{Diagram 2} \end{array} \right\rangle_n - q^{1/2} \left\langle \begin{array}{c} \text{Diagram 3} \end{array} \right\rangle_n \\
 &\quad - q^{-1/2} \left\langle \begin{array}{c} \text{Diagram 4} \end{array} \right\rangle_n + \left\langle \begin{array}{c} \text{Diagram 5} \end{array} \right\rangle_n \\
 &= ([n] - q^{1/2}[n-1] - q^{-1/2}[n-1] + [n-2]) \left\langle \begin{array}{c} \text{Diagram 2} \end{array} \right\rangle_n + \left\langle \begin{array}{c} \text{Diagram 6} \end{array} \right\rangle_n \\
 &= \left\langle \begin{array}{c} \text{Diagram 7} \end{array} \right\rangle_n
 \end{aligned}$$

from Lemmas 2.1, 2.3 and 2.4 and so $\langle D \rangle_n$ is invariant under the Reidemeister move II. Next we prove the invariance under the Reidemeister move III. Since we have

$$\left\langle \begin{array}{c} \text{Diagram 1} \end{array} \right\rangle_n = q^{1/2} \left\langle \begin{array}{c} \text{Diagram 2} \end{array} \right\rangle_n - \left\langle \begin{array}{c} \text{Diagram 3} \end{array} \right\rangle_n$$

and

$$\left\langle \begin{array}{c} \text{Diagram 1} \end{array} \right\rangle_n = q^{1/2} \left\langle \begin{array}{c} \text{Diagram 2} \end{array} \right\rangle_n - \left\langle \begin{array}{c} \text{Diagram 3} \end{array} \right\rangle_n,$$

it suffices to show that

The diagram shows an equality between two configurations of strands. On the left, there are two vertical strands on the far left, each with an upward arrow labeled 1. These strands enter a central box. Inside the box, the strands cross and then exit at the top as two strands with upward arrows labeled 1. At the bottom of the box, two strands enter with upward arrows labeled 1, which then merge into a single strand with an upward arrow labeled 2. The entire configuration is labeled with a subscript n at the bottom right. On the right side of the equals sign, the configuration is similar but the crossing inside the box is different, and the bottom strands are labeled 1 and 1 instead of 1 and 2. It is also labeled with a subscript n .

Since

This block contains a large diagrammatic equation. The leftmost term is the same as the one in the first equation. It is followed by an equals sign and a series of terms. The first term is multiplied by q and shows two vertical strands on the left with upward arrows labeled 1, entering a box that contains two vertical strands with upward arrows labeled 1, and exiting as two strands with upward arrows labeled 1. The bottom of the box is labeled with a subscript n . The second term is subtracted and is multiplied by $-q^{1/2}$; it shows two vertical strands on the left with upward arrows labeled 1, entering a box that contains a single vertical strand with an upward arrow labeled 1, and exiting as two strands with upward arrows labeled 1. The bottom of the box is labeled with a subscript n . The third term is added and shows two vertical strands on the left with upward arrows labeled 1, entering a box that contains a more complex internal structure with multiple crossings and arrows labeled 1 and 2, and exiting as two strands with upward arrows labeled 1. The bottom of the box is labeled with a subscript n . The fourth term is added and shows two vertical strands on the left with upward arrows labeled 1, entering a box that contains a similar complex internal structure, and exiting as two strands with upward arrows labeled 1. The bottom of the box is labeled with a subscript n . The equation continues with a minus sign and a term multiplied by $-q^{1/2}$, followed by a plus sign and a term. The final term is a large, complex diagram with multiple crossings and arrows labeled 1 and 2, with a subscript n at the bottom right.

The figure consists of two diagrams representing the decomposition of a product of two elements in a Coxeter group. The left diagram shows a product of two elements, each with three strands labeled 1, 2, and 3. The right diagram shows a product of two elements, each with three strands labeled 1, 2, and 3, with a coefficient $q^{1/2}$.

from Lemmas 2.2 and 2.5 and

The figure shows a sequence of seven diagrams representing the reduction of a four-loop diagram. The diagrams are arranged in three rows. The first row shows the initial diagram on the left, followed by an equals sign, then two diagrams with coefficients q and $-q^{1/2}$. The second row shows two diagrams with coefficients $-q^{1/2}$ and $+$. The third row shows the final result as a sum of three terms: a negative term, a term with coefficient $-q^{1/2}$, and a positive term. All diagrams consist of lines with arrows and labels 1, 2, and n .

$$= \left\langle \begin{array}{c} \text{Diagram 1} \end{array} \right\rangle_n - q^{1/2} \left\langle \begin{array}{c} \text{Diagram 2} \end{array} \right\rangle_n,$$

we have the required formula from Lemma 2.6.

Since the Reidemeister move III with other types of orientations can be obtained from the Reidemeister moves II and III described above (see [12]), the proof is complete. \square

If we define $P_n(D) = q^{(n/2)(-w(D))} \langle D \rangle_n$ with $w(D)$ the writhe (the algebraic sum of the crossings) of D , then we have

THEOREM 3.2. $P_n(D)$ is invariant under the Reidemeister moves I, II, and III and satisfies the following skein relation.

$$q^{n/2} P_n(D_+) - q^{-n/2} P_n(D_-) = (q^{1/2} - q^{-1/2}) P_n(D_0),$$

where D_+ , D_- and D_0 are identical link diagrams except near a crossing as described in Figure 3.1.




FIGURE 3.1
skein triple

The proof of this theorem follows immediately from the following lemma.

LEMMA 3.3. $\left\langle \begin{array}{c} \text{Diagram 1} \end{array} \right\rangle_n = q^{n/2} \left\langle \begin{array}{c} \text{Diagram 2} \end{array} \right\rangle_n$

and

$$\left\langle \begin{array}{c} \text{Diagram 1} \end{array} \right\rangle_n = q^{-n/2} \left\langle \begin{array}{c} \text{Diagram 2} \end{array} \right\rangle_n.$$

Proof. From Lemmas 2.1 and 2.3, we have

$$(3.1) \quad \left\langle \begin{array}{c} \uparrow \\ \bigcirc \end{array} \right\rangle_n = q^{1/2} \left\langle \begin{array}{c} 1 \uparrow \\ \bigcirc \\ 1 \end{array} \right\rangle_n - \left\langle \begin{array}{c} 1 \uparrow \\ 2 \bigcirc 1 \\ 1 \end{array} \right\rangle_n$$

$$(3.2) \quad = (q^{1/2}[n] - [n-1]) \left\langle \begin{array}{c} \uparrow 1 \\ \end{array} \right\rangle_n.$$

Since

$$q^{1/2}[n] - [n-1] = q^{n/2},$$

the first equality follows. The second equality follows similarly and the proof is complete. \square

Therefore P_n defines a link invariant and so we can put $P_n(L) = P_n(D)$ for the link L presented by D . Then P_n is a version of the HOMFLY polynomial [1], [13].

REMARK 3.4. If we define $[D]_n$ to be

$$\left[\begin{array}{c} \uparrow \quad \uparrow \\ \diagdown \quad \diagup \\ \end{array} \right]_n = Aq^{1/2} \left[\begin{array}{c} 1 \uparrow \\ \end{array} \right]_n - A \left[\begin{array}{c} 1 \quad 1 \\ \diagdown \quad \diagup \\ 2 \\ \diagup \quad \diagdown \\ 1 \quad 1 \end{array} \right]_n$$

and

$$\left[\begin{array}{c} \uparrow \quad \uparrow \\ \diagup \quad \diagdown \\ \end{array} \right]_n = A^{-1}q^{-1/2} \left[\begin{array}{c} 1 \uparrow \\ \end{array} \right]_n - A^{-1} \left[\begin{array}{c} 1 \quad 1 \\ \diagup \quad \diagdown \\ 2 \\ \diagdown \quad \diagup \\ 1 \quad 1 \end{array} \right]_n$$

with A an indeterminate, we also have a framed link invariant.

When $n = 2$ and $A = q^{-1/4}$, we have a version of the Kauffman bracket naturally defined from representation theory of $U_q(\mathfrak{sl}(2, \mathbb{C}))$ [6, Theorem 4.3].

For $n = 3$ we have G. Kuperberg's recursive formula [8] as follows. Consider an oriented, trivalent, plane graph with flow less than or equal to three. If we reverse the orientations of all the edges with flow two, remove all the edges with flow three, and forget the flow, then we have a trivalent graph without flow such that at every vertex all the edges are in or out. Putting $A = q^{-1/6}$ and replacing q with q^{-1} , we have Kuperberg's formula. (This corresponds to the fact that the two-fold anti-symmetric tensor of the vector representation of $SU(3)$ is isomorphic to its dual.) See also [12].

§4. STATE MODEL OF TURAEV AND JONES

In this section we show that our definition of the HOMFLY polynomial gives the state model due to V. G. Turaev and V. F. R. Jones [16], [4]. Moreover we can recover the R -matrix for the q -deformation of the universal enveloping algebra $U_q(\mathfrak{sl}(n, \mathbb{C}))$ found by M. Jimbo [2].

For readers' convenience, we first review Turaev's state model. Let $R: V^{\otimes 2} \rightarrow V^{\otimes 2}$ be an isomorphism with V an n -dimensional vector space over \mathbb{C} . It is called an R -matrix if it satisfies the so-called Yang-Baxter equation:

$$(R \otimes \text{id}) \circ (\text{id} \otimes R) \circ (R \otimes \text{id}) = (\text{id} \otimes R) \circ (R \otimes \text{id}) \circ (\text{id} \otimes R).$$

Let $\{e_1, e_2, \dots, e_n\}$ be a basis of V and $R_{i,j}^{k,l}$ be the entry of R with respect to a fixed basis of V , which is the coefficient of $e_k \otimes e_l$ of $R(e_i \otimes e_j)$.

Given a link diagram D , we regard it as a 4-valent graph (a crossing corresponding to a 4-valent vertex) and denote it by \tilde{D} . See Figure 4.1. Here the $+$ ($-$, respectively) sign indicates that the vertex comes from a positive (negative, respectively) crossing.




FIGURE 4.1

Change crossings to 4-valent vertices

A state $\tilde{\sigma}$ of \tilde{D} is a mapping from the edge set of \tilde{D} to \mathcal{N} . Then we define the weight $\text{wt}_R(\tilde{v}, \tilde{\sigma})$ of a vertex \tilde{v} to be R_{ab}^{cd} if \tilde{v} comes from a positive crossing and $(R^{-1})_{ab}^{cd}$ if it comes from a negative one, where $a, b, c, d \in \mathcal{N}$ are the values of the four edges adjacent to \tilde{v} (see Figure 4.2).




FIGURE 4.2

State around a vertex \tilde{v}

If we can define a suitable “rotation number” $\widetilde{\text{rot}}(\tilde{\sigma})$ for each state, then the quantity

$$\sum_{\tilde{\sigma}:\text{state}} \left\{ \prod_{v:\text{vertex}} \text{wt}_R(\tilde{v}, \tilde{\sigma}) \right\} q^{\widetilde{\text{rot}}(\tilde{\sigma})}$$

becomes a framed link invariant.

Now we will proceed conversely and define an R -matrix from our framed link invariant $\langle D \rangle_n$.

First we define a weight $\widetilde{\text{wt}}(\tilde{v}, \tilde{\sigma})$ by using $\langle D \rangle_n$ as follows. We assume that for a state $\tilde{\sigma}$, $a, b, c, d \in \mathcal{N}$ appear around a vertex \tilde{v} . We define

$$\begin{aligned} \widetilde{\text{wt}} \left(\begin{array}{c} c \uparrow \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ a \uparrow \quad b \uparrow \\ + \end{array} \right) &= \left\langle \begin{array}{c} \{c\} \uparrow \\ \diagdown \quad \diagup \\ \{a\} \uparrow \end{array} \begin{array}{c} \{d\} \uparrow \\ \diagdown \quad \diagup \\ \{b\} \uparrow \end{array} \right\rangle_n \\ &= q^{1/2} \text{wt} \left(\begin{array}{c} \{c\} \uparrow \\ \{a\} \uparrow \end{array} \begin{array}{c} \{d\} \uparrow \\ \{b\} \uparrow \end{array} \right) - \text{wt} \left(\begin{array}{c} \{c\} \uparrow \\ \diagdown \quad \diagup \\ \{a\} \uparrow \quad \{b\} \uparrow \\ \{a,b\} \uparrow \end{array} \right) \end{aligned}$$

and

$$\begin{aligned} \widetilde{\text{wt}} \left(\begin{array}{c} c \uparrow \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ a \uparrow \quad b \uparrow \\ - \end{array} \right) &= \left\langle \begin{array}{c} \{c\} \uparrow \\ \diagdown \quad \diagup \\ \{a\} \uparrow \end{array} \begin{array}{c} \{d\} \uparrow \\ \diagdown \quad \diagup \\ \{b\} \uparrow \end{array} \right\rangle_n \\ &= q^{-1/2} \text{wt} \left(\begin{array}{c} \{c\} \uparrow \\ \{a\} \uparrow \end{array} \begin{array}{c} \{d\} \uparrow \\ \{b\} \uparrow \end{array} \right) - \text{wt} \left(\begin{array}{c} \{c\} \uparrow \\ \diagdown \quad \diagup \\ \{a\} \uparrow \quad \{b\} \uparrow \\ \{a,b\} \uparrow \end{array} \right). \end{aligned}$$

Here the first terms of the right hand sides are zero if $a \neq c$ or $b \neq d$ and the second terms are zero if $\{a, b\} \neq \{c, d\}$. Therefore, for $a, b \in \mathcal{N}$ ($a \neq b$) we have

$$\begin{aligned} \widetilde{\text{wt}} \left(\begin{array}{c} a \uparrow \\ \diagdown \quad \diagup \\ \bullet \\ \diagup \quad \diagdown \\ a \uparrow \quad b \uparrow \\ + \end{array} \right) &= q^{1/2} \text{wt} \left(\begin{array}{c} \{a\} \uparrow \\ \{a\} \uparrow \end{array} \begin{array}{c} \{b\} \uparrow \\ \{b\} \uparrow \end{array} \right) - \text{wt} \left(\begin{array}{c} \{a\} \uparrow \\ \diagdown \quad \diagup \\ \{a\} \uparrow \quad \{b\} \uparrow \\ \{a,b\} \uparrow \end{array} \right) \\ &= q^{1/2} - q^{\text{sign}(b-a)/2} \\ &= \begin{cases} q^{1/2} - q^{-1/2} & \text{if } a > b, \\ 0 & \text{if } a < b. \end{cases} \end{aligned}$$

Similarly we have

$$\widetilde{\text{wt}} \left(\begin{array}{c} a \uparrow \\ b \uparrow \\ \text{+} \\ b \downarrow \\ a \downarrow \end{array} \right) = -1 \quad \text{and} \quad \widetilde{\text{wt}} \left(\begin{array}{c} a \uparrow \\ a \uparrow \\ \text{+} \\ a \downarrow \\ a \downarrow \end{array} \right) = q^{1/2}.$$

We have a similar formula for a negative crossing.

Therefore we see that our graph invariant gives an R -matrix of the form

$$R_{ij}^{kl} = \begin{cases} q^{1/2} - q^{-1/2} & \text{if } i = k > j = l, \\ -1 & \text{if } i = l \neq j = k, \\ q^{1/2} & \text{if } i = j = k = l, \\ 0 & \text{otherwise;} \end{cases}$$

and

$$(R^{-1})_{ij}^{kl} = \begin{cases} -q^{1/2} + q^{-1/2} & \text{if } i = k < j = l, \\ -1 & \text{if } i = l \neq j = k, \\ q^{-1/2} & \text{if } i = j = k = l, \\ 0 & \text{otherwise;} \end{cases}$$

which coincides with $-R(q^{1/2})^{-1}$, where $R(q^{1/2})$ is the R -matrix given in [16], replacing q with $q^{1/2}$.

§5. INVARIANTS CORRESPONDING TO ANTI-SYMETRIC TENSORS

In this section, we will show briefly that our graph invariant gives the quantum link invariant each of its component equipped with an anti-symmetric tensor of the standard n -dimensional representation of $SU(n)$.

Let D be a link diagram each of its component colored with an integer i ($1 \leq i \leq n$). This corresponds to the i -fold anti-symmetric tensor of the standard representation of $SU(n)$.

Then $\langle D \rangle_n$ is defined by

$$\left\langle \begin{array}{c} i \uparrow \\ \text{crossing} \\ j \uparrow \end{array} \right\rangle_n = \sum_{k=0}^i (-1)^{k+(j+1)i} q^{(i-k)/2} \left\langle \begin{array}{c} i \uparrow \quad j+k-i \uparrow \quad j \uparrow \\ j+k \uparrow \quad i-k \uparrow \\ j \uparrow \quad k \uparrow \quad i \uparrow \end{array} \right\rangle_n, \quad \text{for } i \leq j$$

and

$$\left\langle \begin{array}{c} i \uparrow \\ \text{crossing} \\ j \uparrow \end{array} \right\rangle_n = \sum_{k=0}^j (-1)^{k+(i+1)j} q^{(j-k)/2} \left\langle \begin{array}{c} i \uparrow \quad i+k-j \uparrow \quad j \uparrow \\ j-k \uparrow \quad i+k \uparrow \\ j \uparrow \quad k \uparrow \quad i \uparrow \end{array} \right\rangle_n, \quad \text{for } i > j$$

For a negative crossing, replace q with q^{-1} .

Now we will show

THEOREM 5.1. *The quantity $\langle D \rangle_n$ with D a colored link diagram is invariant under the Reidemeister moves II and III. Thus it is a colored framed link invariant.*

To prove the theorem above, we prepare some lemmas:

LEMMA 5.2.

$$\left\langle \begin{array}{c} 1 \uparrow \quad i+1 \uparrow \quad i \uparrow \\ i \downarrow \quad i+1 \downarrow \\ 1 \downarrow \quad i \downarrow \end{array} \right\rangle_n = [n-i-1] \left\langle \begin{array}{c} 1 \uparrow \quad i \uparrow \\ 1 \downarrow \quad i \downarrow \end{array} \right\rangle_n + \left\langle \begin{array}{c} 1 \uparrow \\ i \downarrow \end{array} \right\rangle_n.$$

Proof. The proof of this lemma is similar to that of Lemma 2.4 and we leave it to the reader. \square

LEMMA 5.3.

$$(5.1) \quad \left\langle \begin{array}{c} i \nearrow \\ \searrow j \\ \nearrow k \\ \searrow i+j \end{array} \right\rangle_n = \left\langle \begin{array}{c} i \nearrow \\ \searrow j \\ \nearrow k \\ \searrow i+j \end{array} \right\rangle_n.$$

Proof. It suffices to prove the case $i = 1$ or $j = 1$ since we have

$$\begin{aligned} \left\langle \begin{array}{c} i \nearrow \\ \searrow j \\ \nearrow k \\ \searrow i+j \end{array} \right\rangle_n &= \frac{1}{[j]} \left\langle \begin{array}{c} i \nearrow \\ \searrow j-1 \\ \nearrow k \\ \searrow i+j \end{array} \right\rangle_n \\ &= \frac{1}{[j]} \left\langle \begin{array}{c} i \nearrow \\ \searrow j-1 \\ \nearrow k \\ \searrow i+j \end{array} \right\rangle_n = \frac{1}{[j]} \left\langle \begin{array}{c} i \nearrow \\ \searrow j-1 \\ \nearrow k \\ \searrow i+j \end{array} \right\rangle_n \end{aligned}$$

and the conclusion follows from the case $i = 1$ or $j = 1$ and induction. Here we use Lemma A.1 in the first equality.

We only prove the case $j = 1$ and $i < k$ since the remaining case is similar. From the definition, the left hand side of (5.1) with $j = 1$ equals

$$\sum_{l=0}^{i+1} (-1)^{l+(k+1)(i+1)} q^{(i-l+1)/2} \left\langle \begin{array}{c} i \nearrow \\ \searrow 1 \\ \nearrow k \\ \searrow i+1 \end{array} \right\rangle_n.$$

The right hand side becomes

$$\begin{aligned}
 & \sum_{l=0}^i (-1)^{l+(k+1)(i+1)} q^{(i-l+1)/2} \left(\text{Diagram 1} \right) \\
 & + \sum_{l=0}^i (-1)^{l+(k+1)i+k} q^{(i-l)/2} \left(\text{Diagram 2} \right) .
 \end{aligned}$$

Sliding the bar colored with l using Lemma 2.6, the first diagram becomes

$$\begin{aligned}
 & \left(\text{Diagram 1} \right) = [i-l] \left(\text{Diagram 2} \right) + \left(\text{Diagram 3} \right) \\
 & = [i-l] \left(\text{Diagram 4} \right) + \left(\text{Diagram 5} \right) ,
 \end{aligned}$$

where the first equality follows from Lemma A.7 below.

The second diagram turns out to be

$$\left(\text{Diagram 1} \right) = [i - l + 1] \left(\text{Diagram 2} \right) .$$

Diagram 1: A braid-like diagram with three vertical strands labeled i , 1 , and k at the top. The bottom strands are labeled k , l , and $i+1$. The right side of the diagram is enclosed in a large bracket labeled n . Internal labels include $k+l$, $k+l-i$, $i-l$, $i-l+1$, and 1 .

Diagram 2: A similar diagram to Diagram 1, but with a different internal structure. It also has top strands $i, 1, k$ and bottom strands $k, l, i+1$, with a right bracket labeled n . Internal labels include $k+l$, $k+l-i$, and $i-l+1$.

Therefore the right hand side of (5.1) becomes

$$\left\{ \sum_{l=0}^i (-1)^{l+(k+1)(i+1)} q^{(i-l+1)/2} [i-l] + \sum_{l=0}^i (-1)^{l+(k+1)i+k} q^{(i-l)/2} [i-l+1] \right\}$$

$$\times \left(\text{Diagram 3} \right)$$

Diagram 3: A diagram with top strands $i, 1, k$ and bottom strands $k, l, i+1$. The right side is enclosed in a large bracket labeled n . Internal labels include $k+l$, $k+l-i$, and $i-l+1$.

$$+ \sum_{l=0}^i (-1)^{l+(k+1)(i+1)} q^{(i-l+1)/2} \left(\text{Diagram 4} \right)$$

Diagram 4: A diagram with top strands $i, 1, k$ and bottom strands $k, l, i+1$. The right side is enclosed in a large bracket labeled n . Internal labels include $i+1$, $k+l-i-1$, $k+l$, and $i-l+1$.

$$= \sum_{l=0}^i (-1)^{l+(k+1)(i+1)+1} \left(\text{Diagram 5} \right)$$

Diagram 5: A diagram with top strands $i, 1, k$ and bottom strands $k, l, i+1$. The right side is enclosed in a large bracket labeled n . Internal labels include $k+l$, $k+l-i$, and $i-l+1$.

$$+ \sum_{l=0}^i (-1)^{l+(k+1)(i+1)} q^{(i-l+1)/2} \left\langle \begin{array}{c} i \nearrow \quad 1 \nearrow \quad k \nearrow \\ i+1 \nearrow \quad k+l-i-1 \nearrow \\ k+l \nearrow \quad i-l+1 \nearrow \\ k \nearrow \quad l \nearrow \quad i+1 \nearrow \end{array} \right\rangle_n.$$

We finally see that the right hand side of (5.1) minus the left hand side equals

$$\begin{aligned} & \sum_{l=0}^i (-1)^{l+(k+1)(i+1)+1} \left\langle \begin{array}{c} i \nearrow \quad 1 \nearrow \quad k \nearrow \\ k+l-i \nearrow \quad k+1 \nearrow \\ k+l \nearrow \quad i-l+1 \nearrow \\ k \nearrow \quad l \nearrow \quad i+1 \nearrow \end{array} \right\rangle_n - (-1)^{i+1+(k+1)(i+1)} \left\langle \begin{array}{c} i \nearrow \quad 1 \nearrow \quad k \nearrow \\ i+1 \nearrow \quad k+i+1 \nearrow \\ k \nearrow \quad i+1 \nearrow \end{array} \right\rangle_n \\ &= \sum_{l=0}^{i+1} (-1)^{l+(k+1)(i+1)+1} \left\langle \begin{array}{c} i \nearrow \quad 1 \nearrow \quad k \nearrow \\ k+l-i \nearrow \quad k+1 \nearrow \\ k+l \nearrow \quad i-l+1 \nearrow \\ k \nearrow \quad l \nearrow \quad i+1 \nearrow \end{array} \right\rangle_n = 0. \end{aligned}$$

Here the last equality follows from Lemma A.9, completing the proof. \square

PROOF OF THEOREM 5.1

The invariance under the Reidemeister move II. We will first show

$$\left\langle \begin{array}{c} i \nearrow \quad j \nearrow \\ \text{Reidemeister II move} \end{array} \right\rangle_n = \left\langle \begin{array}{c} i \nearrow \\ j \nearrow \end{array} \right\rangle_n.$$

It suffices to show the case $i = 1$ from Lemmas A.1 and 5.3 since

$$\begin{aligned}
 & \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \end{array} \right\rangle_n = \frac{1}{[i]} \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \\ 1 \leftarrow i-1 \end{array} \right\rangle_n = \frac{1}{[i]} \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \\ 1 \rightarrow i-1 \end{array} \right\rangle_n \\
 & = \frac{1}{[i]} \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \\ 1 \leftarrow i-1 \end{array} \right\rangle_n = \frac{1}{[i]} \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \\ 1 \leftarrow i-1 \end{array} \right\rangle_n \\
 & = \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \end{array} \right\rangle_n,
 \end{aligned}$$

where the second equality follows from Lemma 5.3 and the fourth by induction on i . Now we have

$$\begin{aligned}
 & \left\langle \begin{array}{c} 1 \uparrow \\ j \uparrow \end{array} \right\rangle_n \\
 & = \left\langle \begin{array}{c} 1 \uparrow \\ j-1 \rightarrow j \\ j \uparrow \\ 1 \leftarrow j-1 \end{array} \right\rangle_n - q^{1/2} \left\langle \begin{array}{c} 1 \uparrow \\ j-1 \rightarrow j \\ j \uparrow \\ j+1 \uparrow \end{array} \right\rangle_n - q^{-1/2} \left\langle \begin{array}{c} 1 \uparrow \\ j+1 \uparrow \\ j \uparrow \\ j-1 \leftarrow j \end{array} \right\rangle_n + \left\langle \begin{array}{c} 1 \uparrow \\ j+1 \uparrow \\ j \uparrow \\ j+1 \uparrow \end{array} \right\rangle_n
 \end{aligned}$$

$$\begin{aligned}
&= \left\langle \begin{array}{c} 1 \uparrow \\ j \uparrow \\ 1 \uparrow \end{array} \begin{array}{c} j-1 \rightarrow \\ j \rightarrow \\ j-1 \rightarrow \end{array} \right\rangle_n + \left(-q^{1/2}[j] - q^{-1/2}[j] + [j+1] \right) \left\langle \begin{array}{c} 1 \uparrow \\ j+1 \uparrow \\ 1 \uparrow \end{array} \begin{array}{c} j \rightarrow \\ j+1 \rightarrow \\ j \rightarrow \end{array} \right\rangle_n \\
&= \left\langle \begin{array}{c} 1 \uparrow \\ j \uparrow \\ 1 \uparrow \end{array} \begin{array}{c} j \rightarrow \\ j \rightarrow \\ j \rightarrow \end{array} \right\rangle_n + \left([j-1] - q^{1/2}[j] - q^{-1/2}[j] + [j+1] \right) \left\langle \begin{array}{c} 1 \uparrow \\ j+1 \uparrow \\ 1 \uparrow \end{array} \begin{array}{c} j \rightarrow \\ j+1 \rightarrow \\ j \rightarrow \end{array} \right\rangle_n \\
&= \left\langle \begin{array}{c} 1 \uparrow \\ j \uparrow \\ 1 \uparrow \end{array} \begin{array}{c} j \rightarrow \\ j \rightarrow \\ j \rightarrow \end{array} \right\rangle_n,
\end{aligned}$$

where we use Lemma A.4 in the third equality.

Next we will show

$$\left\langle \begin{array}{c} i \uparrow \\ j \uparrow \\ j \downarrow \end{array} \right\rangle_n = \left\langle \begin{array}{c} i \uparrow \\ j \downarrow \end{array} \right\rangle_n.$$

It also suffices to show the case $i = 1$ as above. We have

$$\begin{aligned}
&\left\langle \begin{array}{c} 1 \uparrow \\ j \uparrow \\ j \downarrow \end{array} \right\rangle_n \\
&= \left\langle \begin{array}{c} 1 \uparrow \\ j-1 \rightarrow \\ j \rightarrow \\ j-1 \rightarrow \end{array} \right\rangle_n - q^{1/2} \left\langle \begin{array}{c} 1 \uparrow \\ j+1 \rightarrow \\ j \rightarrow \\ j-1 \rightarrow \end{array} \right\rangle_n - q^{-1/2} \left\langle \begin{array}{c} 1 \uparrow \\ j-1 \rightarrow \\ j \rightarrow \\ j+1 \rightarrow \end{array} \right\rangle_n + \left\langle \begin{array}{c} 1 \uparrow \\ j+1 \rightarrow \\ j \rightarrow \\ j+1 \rightarrow \end{array} \right\rangle_n \\
&= \left([n-j+1] - q^{1/2}[n-j] - q^{-1/2}[n-j] \right) \left\langle \begin{array}{c} 1 \uparrow \\ j-1 \rightarrow \\ j \rightarrow \\ j \rightarrow \end{array} \right\rangle_n + \left\langle \begin{array}{c} 1 \uparrow \\ j+1 \rightarrow \\ j \rightarrow \\ j+1 \rightarrow \end{array} \right\rangle_n
\end{aligned}$$

$$\begin{aligned}
&= \left([n-j+1] - q^{1/2}[n-j] - q^{-1/2}[n-j] + [n-j-1] \right) \left\langle \begin{array}{c} 1 \uparrow \\ \text{---} j-1 \text{---} \\ 1 \downarrow \end{array} \begin{array}{c} j \\ \text{---} j \text{---} \\ j \downarrow \end{array} \right\rangle_n \\
&\quad + \left\langle \begin{array}{c} 1 \uparrow \\ \text{---} \\ 1 \downarrow \end{array} \begin{array}{c} j \\ \text{---} \\ j \downarrow \end{array} \right\rangle_n \\
&= \left\langle \begin{array}{c} 1 \uparrow \\ \text{---} \\ 1 \downarrow \end{array} \begin{array}{c} j \\ \text{---} \\ j \downarrow \end{array} \right\rangle_n,
\end{aligned}$$

where we use Lemma 5.2 in the third equality. Now the proof for the Reidemeister move II is complete.

The invariance under the Reidemeister move III. This is proved by repeated application of Lemma 5.3 and details are omitted. See the proof of Theorem 3.1.

A. APPENDIX

In this appendix, we give proofs of lemmas used in the previous section.

LEMMA A.1.

$$\left\langle \begin{array}{c} i \uparrow \\ \text{---} j \text{---} i-j \\ i \downarrow \end{array} \right\rangle_n = [j] \left\langle \begin{array}{c} i \uparrow \\ \text{---} \\ i \downarrow \end{array} \right\rangle_n$$

for $i \geq j \geq 0$.

Proof. The proof for $j = 1$ is similar to that of Lemma 2.2 and omitted. For $j > 1$ we have

$$\begin{aligned}
\left\langle \begin{array}{c} i \uparrow \\ \text{---} j \text{---} i-j \\ i \downarrow \end{array} \right\rangle_n &= \frac{1}{[j]} \left\langle \begin{array}{c} i \uparrow \\ \text{---} j-1 \text{---} 1 \text{---} i-j \\ j \uparrow \text{---} j \downarrow \end{array} \right\rangle_n = \frac{1}{[j]} \left\langle \begin{array}{c} i \uparrow \\ \text{---} j-1 \text{---} 1 \text{---} i-j+1 \\ j \uparrow \text{---} j \downarrow \end{array} \right\rangle_n \\
&= \frac{[i-j+1]}{[j]} \left\langle \begin{array}{c} i \uparrow \\ \text{---} j-1 \text{---} i-j+1 \\ i \downarrow \end{array} \right\rangle_n = \frac{[i-j+1]}{[j]} [j-1] \left\langle \begin{array}{c} i \uparrow \\ \text{---} \\ i \downarrow \end{array} \right\rangle_n = [j] \left\langle \begin{array}{c} i \uparrow \\ \text{---} \\ i \downarrow \end{array} \right\rangle_n,
\end{aligned}$$

where the second equality follows from Lemma 2.6 and the fourth by induction. The proof is complete. \square

LEMMA A.2.

$$\left\langle \begin{array}{c} i \\ j+k \\ j \end{array} \begin{array}{c} j+k-i \\ k \\ i \end{array} \right\rangle_n = \left\langle \begin{array}{c} i \\ i-k \\ j \end{array} \begin{array}{c} k \\ j+k-i \\ i \end{array} \right\rangle_n$$

for $j+k \geq i \geq k \geq 0$.

Proof. By a π -rotation and orientation reversing, we get the right hand side from the left hand side. So there is a one to one correspondence between the states of both hand sides and the equality follows from the definition. \square

LEMMA A.3.

$$\left\langle \begin{array}{c} 1 \\ 2 \\ 1 \end{array} \begin{array}{c} 1 \\ j-1 \\ j \end{array} \right\rangle_n = \left\langle \begin{array}{c} 1 \\ 1 \end{array} \begin{array}{c} j \\ j+1 \\ j \end{array} \right\rangle_n + [j-1] \left\langle \begin{array}{c} 1 \\ 1 \end{array} \begin{array}{c} j \\ j \end{array} \right\rangle_n$$

for $j \geq 1$.

Proof. We can prove the equality directly from the definition as in the proofs in the lemmas in §2. Details are omitted. \square

LEMMA A.4.

$$\left\langle \begin{array}{c} 1 \\ k+1 \\ 1 \end{array} \begin{array}{c} k \\ j-k \\ j \end{array} \right\rangle_n = \begin{bmatrix} j-1 \\ k-1 \end{bmatrix} \left\langle \begin{array}{c} 1 \\ 1 \end{array} \begin{array}{c} j \\ j+1 \\ j \end{array} \right\rangle_n + \begin{bmatrix} j-1 \\ k \end{bmatrix} \left\langle \begin{array}{c} 1 \\ 1 \end{array} \begin{array}{c} j \\ j \end{array} \right\rangle_n$$

for $j \geq k \geq 1$.

Proof. From Lemma A.3 (substituting j with k) the left hand side becomes

$$\begin{aligned}
 & \left\langle \begin{array}{c} 1 \uparrow \\ 2 \leftarrow 1 \rightarrow k \\ 1 \uparrow \end{array} \begin{array}{c} j \\ k \\ k-1 \\ j-k \\ k \\ j \end{array} \right\rangle_n - [k-1] \left\langle \begin{array}{c} 1 \uparrow \\ k \leftarrow j \rightarrow j-k \\ j \end{array} \right\rangle_n \\
 &= \left\langle \begin{array}{c} 1 \uparrow \\ 2 \leftarrow 1 \rightarrow j \\ 1 \uparrow \end{array} \begin{array}{c} j \\ j-1 \\ k-1 \\ j-k \\ j-1 \\ j \end{array} \right\rangle_n - [k-1] \left[\begin{array}{c} j \\ k \end{array} \right] \left\langle \begin{array}{c} 1 \uparrow \\ \uparrow j \end{array} \right\rangle_n \\
 &= \left[\begin{array}{c} j-1 \\ k-1 \end{array} \right] \left\langle \begin{array}{c} 1 \uparrow \\ 2 \leftarrow 1 \rightarrow j \\ 1 \uparrow \end{array} \begin{array}{c} j \\ j-1 \\ j-1 \\ j \end{array} \right\rangle_n - [k-1] \left[\begin{array}{c} j \\ k \end{array} \right] \left\langle \begin{array}{c} 1 \uparrow \\ \uparrow j \end{array} \right\rangle_n \\
 &= \left[\begin{array}{c} j-1 \\ k-1 \end{array} \right] \left\langle \begin{array}{c} 1 \uparrow \\ 1 \leftarrow j+1 \rightarrow j \\ 1 \uparrow \end{array} \right\rangle_n + \left(\left[\begin{array}{c} j-1 \\ k-1 \end{array} \right] [j-1] - [k-1] \left[\begin{array}{c} j \\ k \end{array} \right] \right) \left\langle \begin{array}{c} 1 \uparrow \\ \uparrow j \end{array} \right\rangle_n \\
 &= \left[\begin{array}{c} j-1 \\ k-1 \end{array} \right] \left\langle \begin{array}{c} 1 \uparrow \\ 1 \leftarrow j+1 \rightarrow j \\ 1 \uparrow \end{array} \right\rangle_n + \left[\begin{array}{c} j-1 \\ k \end{array} \right] \left\langle \begin{array}{c} 1 \uparrow \\ \uparrow j \end{array} \right\rangle_n,
 \end{aligned}$$

where the third equality follows from Lemma A.3. The proof is complete. \square

LEMMA A.5.

$$\left\langle \begin{array}{c} 2 \uparrow \\ 3 \leftarrow 1 \rightarrow j \\ 2 \uparrow \end{array} \begin{array}{c} j \\ j-1 \\ j \\ j \end{array} \right\rangle_n - \left\langle \begin{array}{c} 2 \uparrow \\ 1 \leftarrow 1 \rightarrow j+1 \\ 2 \uparrow \end{array} \begin{array}{c} j \\ j+1 \\ j \\ j \end{array} \right\rangle_n = [j-2] \left\langle \begin{array}{c} 2 \uparrow \\ \uparrow j \end{array} \right\rangle_n$$

for $j \geq 2$.

Proof. We have

$$\begin{aligned}
 \left\langle \begin{array}{c} 2 \uparrow \\ 3 \uparrow \\ 2 \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ j-1 \uparrow \\ j \uparrow \end{array} \right\rangle_n &= \frac{1}{[j-1]} \left\langle \begin{array}{c} 2 \uparrow \\ 3 \uparrow \\ 2 \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ j-1 \uparrow \\ j-2 \uparrow \\ j-1 \uparrow \\ j \uparrow \end{array} \right\rangle_n \\
 &= \frac{1}{[j-1]} \left\langle \begin{array}{c} 2 \uparrow \\ 3 \uparrow \\ 2 \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ 2 \uparrow \\ 1 \uparrow \\ 2 \uparrow \\ j \uparrow \end{array} \right\rangle_n \\
 &= \frac{1}{[j-1]} \left\langle \begin{array}{c} 2 \uparrow \\ 1 \uparrow \\ 2 \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ 2 \uparrow \\ 3 \uparrow \\ 2 \uparrow \\ j \uparrow \end{array} \right\rangle_n \\
 &= \left\langle \begin{array}{c} 2 \uparrow \\ 1 \uparrow \\ 2 \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ j+1 \uparrow \\ j \uparrow \end{array} \right\rangle_n + \frac{[j-1]}{[j-1]} \left\langle \begin{array}{c} 2 \uparrow \\ 1 \uparrow \\ 2 \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \end{array} \right\rangle_n \\
 &= \left\langle \begin{array}{c} 2 \uparrow \\ 1 \uparrow \\ 2 \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ j+1 \uparrow \\ j \uparrow \end{array} \right\rangle_n + [j-2] \left\langle \begin{array}{c} 2 \uparrow \\ 1 \uparrow \\ 2 \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \end{array} \right\rangle_n,
 \end{aligned}$$

where we use Lemma A.2 in the third equality and Lemma A.4 in the fourth equality. The proof is complete. \square

LEMMA A.6.

$$\left\langle \begin{array}{c} i \uparrow \\ i+1 \uparrow \\ i \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ j-1 \uparrow \\ j \uparrow \end{array} \right\rangle_n - \left\langle \begin{array}{c} i \uparrow \\ i-1 \uparrow \\ i \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ j+1 \uparrow \\ j \uparrow \end{array} \right\rangle_n = [j-i] \left\langle \begin{array}{c} i \uparrow \\ \\ \end{array} \begin{array}{c} \phantom{\xrightarrow{1}} \\ \phantom{\xleftarrow{1}} \end{array} \begin{array}{c} j \uparrow \\ \\ \end{array} \right\rangle_n.$$

Proof. We have

$$\begin{aligned} \left\langle \begin{array}{c} i \uparrow \\ i+1 \uparrow \\ i \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ j-1 \uparrow \\ j \uparrow \end{array} \right\rangle_n &= \frac{1}{[j-1]} \left\langle \begin{array}{c} i \uparrow \\ i+1 \uparrow \\ i \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ j-1 \uparrow \\ \text{diamond} \\ j-1 \uparrow \\ j \uparrow \end{array} \right\rangle_n \\ &= \frac{1}{[j-1]} \left\langle \begin{array}{c} i \uparrow \\ i+1 \uparrow \\ i \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} \text{rectangle} \\ j \uparrow \end{array} \right\rangle_n \\ &= \frac{1}{[j-1]} \left\langle \begin{array}{c} i \uparrow \\ i-1 \uparrow \\ i \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} \text{rectangle} \\ j \uparrow \end{array} \right\rangle_n - \frac{[i-2]}{[j-1]} \left\langle \begin{array}{c} i \uparrow \\ \\ \end{array} \begin{array}{c} \text{diamond} \\ \end{array} \right\rangle_n \\ &= \left\langle \begin{array}{c} i \uparrow \\ i-1 \uparrow \\ i \uparrow \end{array} \begin{array}{c} \xrightarrow{1} \\ \xleftarrow{1} \end{array} \begin{array}{c} j \uparrow \\ j+1 \uparrow \\ j \uparrow \end{array} \right\rangle_n + \left(\frac{[j-2][i]}{[2]} - \frac{[i-2][j]}{[j-1]} \right) \left\langle \begin{array}{c} i \uparrow \\ \\ \end{array} \begin{array}{c} \phantom{\xrightarrow{1}} \\ \phantom{\xleftarrow{1}} \end{array} \begin{array}{c} j \uparrow \\ \\ \end{array} \right\rangle_n \end{aligned}$$

$$= \left\langle \begin{array}{c} i \uparrow \\ i-1 \uparrow \\ i \uparrow \\ j \uparrow \\ j+1 \uparrow \\ j \uparrow \end{array} \right\rangle_n + [j-i] \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \end{array} \right\rangle_n,$$

where we use Lemmas A.3 and A.4 in the third and the fourth equalities respectively. The proof is complete. \square

LEMMA A.7.

$$\left\langle \begin{array}{c} i \uparrow \\ i+k \uparrow \\ 1 \uparrow \\ j \uparrow \\ j-k \uparrow \\ i+j-1 \uparrow \end{array} \right\rangle_n = \begin{bmatrix} j-1 \\ k-1 \end{bmatrix} \left\langle \begin{array}{c} i \uparrow \\ 1 \uparrow \\ j \uparrow \\ i+j \uparrow \\ i+j-1 \uparrow \end{array} \right\rangle_n + \begin{bmatrix} j-1 \\ k \end{bmatrix} \left\langle \begin{array}{c} i \uparrow \\ 1 \uparrow \\ j \uparrow \\ i-1 \uparrow \\ i+j-1 \uparrow \end{array} \right\rangle_n$$

for $j > k \geq 1$.

Proof. We first prove the case $k = 1$. We will show

$$\left\langle \begin{array}{c} i \uparrow \\ i+1 \uparrow \\ 1 \uparrow \\ j \uparrow \\ j-1 \uparrow \\ i+j-1 \uparrow \end{array} \right\rangle_n = \left\langle \begin{array}{c} i \uparrow \\ 1 \uparrow \\ j \uparrow \\ i+j \uparrow \\ i+j-1 \uparrow \end{array} \right\rangle_n + [j-1] \left\langle \begin{array}{c} i \uparrow \\ 1 \uparrow \\ j \uparrow \\ i-1 \uparrow \\ i+j-1 \uparrow \end{array} \right\rangle_n$$

by induction on j . Note that Lemma A.3 is the case $i = 1$.

The left hand side becomes

$$\frac{1}{[i]} \left\langle \begin{array}{c} i \uparrow \\ i+1 \uparrow \\ i \uparrow \\ 1 \uparrow \\ j \uparrow \\ j-1 \uparrow \\ i+j-1 \uparrow \end{array} \right\rangle_n = \frac{1}{[i]} \left\langle \begin{array}{c} i \uparrow \\ i-1 \uparrow \\ i \uparrow \\ 1 \uparrow \\ j \uparrow \\ j+1 \uparrow \\ i+j-1 \uparrow \end{array} \right\rangle_n + \frac{[j-i]}{[i]} \left\langle \begin{array}{c} i \uparrow \\ 1 \uparrow \\ j \uparrow \\ i-1 \uparrow \\ i+j-1 \uparrow \end{array} \right\rangle_n$$

$$\begin{aligned}
&= \frac{1}{[i]} \left\langle \begin{array}{c} i \uparrow \\ \downarrow i-1 \\ \uparrow j+1 \\ \downarrow 1 \\ \uparrow i+j \\ \downarrow i+j-1 \end{array} \right\rangle_n + \frac{[j]}{[i]} \left\langle \begin{array}{c} i \uparrow \\ \downarrow i-1 \\ \uparrow j+1 \\ \downarrow 1 \\ \uparrow i-2 \\ \downarrow i+j-1 \end{array} \right\rangle_n + \frac{[j-i]}{[i]} \left\langle \begin{array}{c} i \uparrow \\ \downarrow 1 \\ \uparrow i-1 \\ \downarrow i+j-1 \end{array} \right\rangle_n \\
&= \left\langle \begin{array}{c} i \uparrow \\ \downarrow 1 \\ \uparrow j \\ \downarrow i+j \\ \uparrow i+j-1 \end{array} \right\rangle_n + \frac{[j][i-1] + [j-i]}{[i]} \left\langle \begin{array}{c} i \uparrow \\ \downarrow 1 \\ \uparrow i-1 \\ \downarrow i+j-1 \end{array} \right\rangle_n,
\end{aligned}$$

which is equal to the right hand side. Here we use Lemma A.6 in the first equality and the inductive hypothesis in the second equality.

The proof for $k > 1$ is similar and omitted. \square

LEMMA A.8.

$$\left\langle \begin{array}{c} i \uparrow \\ \downarrow j+k \\ \uparrow j+1 \\ \downarrow j \\ \uparrow i-k+1 \\ \downarrow i+1 \end{array} \right\rangle_n = \left\langle \begin{array}{c} i \uparrow \\ \downarrow i-k \\ \uparrow j+1 \\ \downarrow j \\ \uparrow j+k-i \\ \downarrow i+1 \end{array} \right\rangle_n + \left\langle \begin{array}{c} i \uparrow \\ \downarrow i-k+1 \\ \uparrow j+1 \\ \downarrow j \\ \uparrow j+k-i-1 \\ \downarrow i+1 \end{array} \right\rangle_n$$

for $j \geq i \geq k \geq 1$.

Proof. For $k = 1$ we have

$$\begin{aligned}
\left\langle \begin{array}{c} i \uparrow \\ \downarrow j+1 \\ \uparrow j+1 \\ \downarrow j \\ \uparrow i \\ \downarrow i+1 \end{array} \right\rangle_n &= \frac{1}{[i]} \left\langle \begin{array}{c} i \uparrow \\ \downarrow j+1 \\ \uparrow j \\ \downarrow j+1 \\ \uparrow i-1 \\ \downarrow i \\ \uparrow i+1 \end{array} \right\rangle_n = \frac{1}{[i]} \left\langle \begin{array}{c} i \uparrow \\ \downarrow i-1 \\ \uparrow j \\ \downarrow j+1 \\ \uparrow j-i+1 \\ \downarrow i \\ \uparrow i+1 \end{array} \right\rangle_n \\
&= \left\langle \begin{array}{c} i \uparrow \\ \downarrow i-1 \\ \uparrow j+1 \\ \downarrow j \\ \uparrow j-i+1 \\ \downarrow i+1 \end{array} \right\rangle_n + \left\langle \begin{array}{c} i \uparrow \\ \downarrow j \\ \uparrow j+1 \\ \downarrow i+1 \end{array} \right\rangle_n,
\end{aligned}$$

where we use Lemma A.7 in the last equality.

The proof for $k > 1$ is similar and left to the reader. \square

PROPOSITION A.9.

$$\sum_{k=0}^{i+1} (-1)^k \left\langle \begin{array}{c} i \uparrow \\ j+k \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xrightarrow{j+k-i} \\ \xleftarrow{k} \end{array} \begin{array}{c} j+1 \uparrow \\ i-k+1 \uparrow \\ i+1 \uparrow \end{array} \right\rangle_n = 0.$$

Proof. From Lemma A.8, we see that the left hand side equals

$$\begin{aligned} & \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xrightarrow{j-i} \end{array} \begin{array}{c} j+1 \uparrow \\ i+1 \uparrow \end{array} \right\rangle_n + \sum_{k=1}^i (-1)^k \left\langle \begin{array}{c} i \uparrow \\ i-k \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xleftarrow{k} \\ \xrightarrow{j+k-i} \end{array} \begin{array}{c} j+1 \uparrow \\ j+k+1 \uparrow \\ i+1 \uparrow \end{array} \right\rangle_n \\ & + \sum_{k=1}^i (-1)^k \left\langle \begin{array}{c} i \uparrow \\ i-k+1 \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xleftarrow{k-1} \\ \xrightarrow{j+k-i-1} \end{array} \begin{array}{c} j+1 \uparrow \\ j+k \uparrow \\ i+1 \uparrow \end{array} \right\rangle_n + (-1)^{i+1} \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xrightarrow{i+j+1} \end{array} \begin{array}{c} j+1 \uparrow \\ i+1 \uparrow \end{array} \right\rangle_n \\ & = \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xrightarrow{j-i} \end{array} \begin{array}{c} j+1 \uparrow \\ i+1 \uparrow \end{array} \right\rangle_n + \sum_{k=1}^i (-1)^k \left\langle \begin{array}{c} i \uparrow \\ i-k \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xleftarrow{k} \\ \xrightarrow{j+k-i} \end{array} \begin{array}{c} j+1 \uparrow \\ j+k+1 \uparrow \\ i+1 \uparrow \end{array} \right\rangle_n \\ & + \sum_{k=0}^{i-1} (-1)^{k+1} \left\langle \begin{array}{c} i \uparrow \\ i-k \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xleftarrow{k} \\ \xrightarrow{j+k-i} \end{array} \begin{array}{c} j+1 \uparrow \\ j+k+1 \uparrow \\ i+1 \uparrow \end{array} \right\rangle_n + (-1)^{i+1} \left\langle \begin{array}{c} i \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xrightarrow{i+j+1} \end{array} \begin{array}{c} j+1 \uparrow \\ i+1 \uparrow \end{array} \right\rangle_n \\ & = 0, \quad \text{and the proof is complete.} \quad \square \end{aligned}$$

More generally we have the following formula, which was suggested by J. Murakami. The proof is left to the reader.

PROPOSITION A.10.

$$\left\langle \begin{array}{c} i \uparrow \\ j+k \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xrightarrow{j+k-i} \\ \xleftarrow{k} \end{array} \begin{array}{c} j+l \uparrow \\ i-k+l \uparrow \\ i+l \uparrow \end{array} \right\rangle_n = \sum_{m=0}^i \begin{bmatrix} l \\ k-m \end{bmatrix} \left\langle \begin{array}{c} i \uparrow \\ i-m \uparrow \\ j \uparrow \end{array} \begin{array}{c} \xrightarrow{m} \\ \xrightarrow{j+m-i} \end{array} \begin{array}{c} j+l \uparrow \\ j+m+l \uparrow \\ i+l \uparrow \end{array} \right\rangle_n$$

for $j \geq i \geq k \geq 1$. Here $\begin{bmatrix} x \\ y \end{bmatrix} = 0$ if $y < 0$ or $y > x$.

B. TABLE OF THE INVARIANT

The following is the table of $P_n/[n]$ for knots with crossings less than or equal to eight. (Note that P_n is always divisible by $[n]$.) We refer the reader to [15] for the notations of knots. We used MAPLE V to translate the Homfly polynomial table in [10] to our invariants.

3 ₁	$-q^{2n} + q^{n+1} + q^{n-1}$
----------------	-------------------------------

4 ₁	$q^n - q + 1 - q^{-1} + q^{-n}$
----------------	---------------------------------

5 ₁	$-q^{3n+1} - q^{3n-1} + q^{2n+2} + q^{2n} + q^{2n-2}$
5 ₂	$-q^{3n} + q^{2n+1} - q^{2n} + q^{2n-1} + q^{n+1} - q^n + q^{n-1}$

6 ₁	$q^{2n} - q^{n+1} + q^n - q^{n-1} - q + 2 - q^{-1} + q^{-n}$
6 ₂	$q^{2n+1} - q^{2n} + q^{2n-1} - q^{n+2} + q^{n+1} - 2q^n + q^{n-1} - q^{n-2} + q + q^{-1}$
6 ₃	$-q^{n+1} + q^n - q^{n-1} + q^2 - q + 3 - q^{-1} + q^{-2} - q^{-n+1} + q^{-n} - q^{-n-1}$

7 ₁	$-q^{4n+2} - q^{4n} - q^{4n-2} + q^{3n+3} + q^{3n+1} + q^{3n-1} + q^{3n-3}$
7 ₂	$-q^{4n} + q^{3n+1} - q^{3n} + q^{3n-1} + q^{2n+1} - 2q^{2n} + q^{2n-1} + q^{n+1} - q^n + q^{n-1}$
7 ₃	$q^{-2n+2} - q^{-2n+1} + q^{-2n} - q^{-2n-1} + q^{-2n-2} + q^{-3n+2} - q^{-3n+1} + 2q^{-3n} - q^{-3n-1} + q^{-3n-2} - q^{-4n+1} - q^{-4n-1}$
7 ₄	$q^{-n+1} - 2q^{-n} + q^{-n-1} + 2q^{-2n+1} - 2q^{-2n} + 2q^{-2n-1} + q^{-3n+1} - 2q^{-3n} + q^{-3n-1} - q^{-4n}$
7 ₅	$-q^{4n+1} + q^{4n} - q^{4n-1} + q^{3n+2} - 2q^{3n+1} + 2q^{3n} - 2q^{3n-1} + q^{3n-2} + q^{2n+2} - q^{2n+1} + 2q^{2n} - q^{2n-1} + q^{2n-2}$
7 ₆	$-q^{3n} + 2q^{2n+1} - 2q^{2n} + 2q^{2n-1} - q^{n+2} + 2q^{n+1} - 3q^n + 2q^{n-1} - q^{n-2} + q - 1 + q^{-1}$
7 ₇	$-q^{n+1} + 2q^n - q^{n-1} + q^2 - 2q + 4 - 2q^{-1} + q^{-2} - 2q^{-n+1} + 2q^{-n} - 2q^{-n-1} + q^{-2n}$

8 ₁	$q^{3n} - q^{2n+1} + q^{2n} - q^{2n-1} - q^{n+1} + 2q^n - q^{n-1} - q + 2 - q^{-1} + q^{-n}$
8 ₂	$q^{3n+2} - q^{3n+1} + q^{3n} - q^{3n-1} + q^{3n-2} - q^{2n+3} + q^{2n+2} - 2q^{2n+1} + q^{2n} - 2q^{2n-1} + q^{2n-2} - q^{2n-3} + q^{n+2} + q^n + q^{n-2}$
8 ₃	$q^{2n} - 2q + 3 - 2q^{-1} - q^{n+1} + 2q^n - q^{n-1} - q^{-n+1} + 2q^{-n} - q^{-n-1} + q^{-2n}$
8 ₄	$q^{2n+1} - q^{2n} + q^{2n-1} - q^{n+2} + 2q^{n+1} - 2q^n + 2q^{n-1} - q^{n-2} - q^2 + q - 2 + q^{-1} - q^{-2} + q^{-n+1} + q^{-n-1}$
8 ₅	$q^{-n+2} + 2q^{-n} + q^{-n-2} - q^{-2n+3} + q^{-2n+2} - 3q^{-2n+1} + q^{-2n} - 3q^{-2n-1} + q^{-2n-2} - q^{-2n-3} + q^{-3n+2} - q^{-3n+1} + 2q^{-3n} - q^{-3n-1} + q^{-3n-2}$
8 ₆	$q^{3n+1} - q^{3n} + q^{3n-1} - q^{2n+2} + 2q^{2n+1} - 3q^{2n} + 2q^{2n-1} - q^{2n-2} - q^{n+2} + 2q^{n+1} - 3q^n + 2q^{n-1} - q^{n-2} + q + q^{-1}$
8 ₇	$-q^2 + q - 1 + q^{-1} - q^{-2} + q^{-n+3} - q^{-n+2} + 3q^{-n+1} - 2q^{-n} + 3q^{-n-1} - q^{-n-2} + q^{-n-3} - q^{-2n+2} + q^{-2n+1} - 2q^{-2n} + q^{-2n-1} - q^{-2n-2}$
8 ₈	$-q^{n+1} + q^n - q^{n-1} + q^2 - 2q + 4 - 2q^{-1} + q^{-2} + q^{-n+2} - 2q^{-n+1} + 3q^{-n} - 2q^{-n-1} + q^{-n-2} - q^{-2n+1} + q^{-2n} - q^{-2n-1}$
8 ₉	$q^{n+2} - q^{n+1} + 2q^n - q^{n-1} + q^{n-2} - q^3 + q^2 - 3q + 3 - 3q^{-1} + q^{-2} - q^{-3} + q^{-n+2} - q^{-n+1} + 2q^{-n} - q^{-n-1} + q^{-n-2}$
8 ₁₀	$-q^2 + q - 2 + q^{-1} - q^{-2} + q^{-n+3} - q^{-n+2} + 4q^{-n+1} - 2q^{-n} + 4q^{-n-1} - q^{-n-2} + q^{-n-3} - q^{-2n+2} + q^{-2n+1} - 3q^{-2n} + q^{-2n-1} - q^{-2n-2}$
8 ₁₁	$q^{3n+1} - q^{3n} + q^{3n-1} - q^{2n+2} + 2q^{2n+1} - 4q^{2n} + 2q^{2n-1} - q^{2n-2} + q - 1 + q^{-1} - q^{n+2} + 3q^{n+1} - 3q^n + 3q^{n-1} - q^{n-2}$
8 ₁₂	$-2q^{n+1} + q^{2n} - 2q^{n-1} + 3q^n + q^2 - 3q + 5 - 3q^{-1} + q^{-2} - 2q^{-n+1} + 3q^{-n} - 2q^{-n-1} + q^{-2n}$
8 ₁₃	$-q^{2n+1} + q^{2n} - q^{2n-1} + q^{n+2} - 2q^{n+1} + 4q^n - 2q^{n-1} + q^{n-2} - q^{-n+1} + 2q^{-n} - q^{-n-1} + q^2 - 3q + 4 - 3q^{-1} + q^{-2}$
8 ₁₄	$q^{3n+1} - 2q^{3n} + q^{3n-1} - q^{2n+2} + 3q^{2n+1} - 4q^{2n} + 3q^{2n-1} - q^{2n-2} - q^{n+2} + 3q^{n+1} - 4q^n + 3q^{n-1} - q^{n-2} + q - 1 + q^{-1}$
8 ₁₅	$q^{5n} - 3q^{4n+1} + 2q^{4n} - 3q^{4n-1} + 2q^{3n+2} - 3q^{3n+1} + 5q^{3n} - 3q^{3n-1} + 2q^{3n-2} + q^{2n+2} - 2q^{2n+1} + 3q^{2n} - 2q^{2n-1} + q^{2n-2}$
8 ₁₆	$-q^{2n+2} + 2q^{2n+1} - 3q^{2n} + 2q^{2n-1} - q^{2n-2} + q^{n+3} - 2q^{n+2} + 4q^{n+1} - 4q^n + 4q^{n-1} - 2q^{n-2} + q^{n-3} - q^2 + 2q - 2 + 2q^{-1} - q^{-2}$

8 ₁₇	$q^{n+2} - 2q^{n+1} + 3q^n - 2q^{n-1} + q^{n-2} - q^3 + 2q^2 - 4q + 5 - 4q^{-1} + 2q^{-2} - q^{-3} + q^{-n+2} - 2q^{-n+1} + 3q^{-n} - 2q^{-n-1} + q^{-n-2}$
8 ₁₈	$q^{n+2} - 3q^{n+1} + 3q^n - 3q^{n-1} + q^{n-2} - q^3 + 3q^2 - 4q + 7 - 4q^{-1} + 3q^{-2} - q^{-3} + q^{-n+2} - 3q^{-n+1} + 3q^{-n} - 3q^{-n-1} + q^{-n-2}$
8 ₁₉	$q^{-3n+3} + q^{-3n+1} + q^{-3n} + q^{-3n-1} + q^{-3n-3} - q^{-4n+2} - q^{-4n+1} - q^{-4n} - q^{-4n-1} - q^{-4n-2} + q^{-5n}$
8 ₂₀	$-q^{2n+1} - q^{2n-1} + q^{n+2} + 2q^n + q^{n-2} - q + 1 - q^{-1}$
8 ₂₁	$q^{3n+1} - q^{3n} + q^{3n-1} - q^{2n+2} + q^{2n+1} - 3q^{2n} + q^{2n-1} - q^{2n-2} + 2q^{n+1} - q^n + 2q^{n-1}$

REFERENCES

- [1] FREYD, P., D. YETTER, J. HOSTE, W.B.R. LICKORISH, K. MILLETT and A. OCNEANU. A new polynomial invariant of knots and links. *Bull. Amer. Math. Soc. (N.S.)* 12 (1985), 239–246.
- [2] JIMBO, M. Quantum R matrix for the generalized Toda system. *Comm. Math. Phys.* 102 (1986), 537–547.
- [3] JONES, V.F.R. A polynomial invariant for knots via von Neumann algebras. *Bull. Amer. Math. Soc. (N.S.)* 12 (1985), 103–111.
- [4] — On knot invariants related to some statistical mechanical models. *Pacific J. Math.* 137 (1989), 311–336.
- [5] KAUFFMAN, L.H. State models and the Jones polynomial. *Topology* 26 (1987), 395–407.
- [6] KIRBY, R. and P. MELVIN. The 3-manifold invariants of Witten and Reshetikhin–Turaev for $\mathfrak{sl}(2, \mathbb{C})$. *Invent. Math.* 105 (1991), 473–545.
- [7] KIRILLOV, A.N. and N.YU. RESHETIKHIN. Representations of the algebra $U_q(\mathfrak{sl}(2))$, q -orthogonal polynomials and invariants of links. *Infinite Dimensional Lie Algebras and Groups*. V.G. Kac, ed., Advanced Series in Mathematical Physics, vol. 7, World Scientific, Singapore, 1989.
- [8] KUPERBERG, G. The quantum G_2 link invariant. *Internat. J. Math.* 5 (1994), 61–85.
- [9] LICKORISH, W.B.R. The skein method for three-manifold invariants. *J. Knot Theory Ramifications* 2 (1993), 171–194.
- [10] — *An Introduction to Knot Theory*. Graduate Texts in Mathematics, vol. 175, Springer-Verlag, 1997.
- [11] MORTON, H.R. Invariants of links and 3-manifolds from skein theory and from quantum groups. *Topics in Knot Theory*. M.E. Bozhüyük, ed., Kluwer, 1993.

- [12] OHTSUKI, T. and S. YAMADA. Quantum $SU(3)$ invariants of 3-manifolds via linear skein theory. *J. Knot Theory Ramifications* 6 (1997), 373–404.
- [13] PRZYTICKI, J.H. and P. TRACZYK. Invariant of links of Conway type. *Kobe J. Math.* 4 (1988), 115–139.
- [14] RESHETIKHIN, N.YU. and V.G. TURAEV. Ribbon graphs and their invariants derived from quantum groups. *Comm. Math. Phys.* 127 (1990), 1–26.
- [15] ROLFSEN, D. *Knots and Links*. Mathematics Lecture Series, vol. 7, Publish or Perish, Inc., Houston, Texas, 1990. Second printing, with corrections.
- [16] TURAEV, V.G. The Yang-Baxter equation and invariants of links. *Invent. Math.* 92 (1988), 527–553.
- [17] YOKOTA, Y. Skeins and quantum $SU(N)$ invariants of 3-manifolds. *Math. Ann.* 307 (1997), 109–138.

(Reçu le 10 mars 1998)

Hitoshi Murakami

Department of Mathematics
School of Science and Engineering
Waseda University
Ohkubo, Shinjuku-ku
Tokyo 169
Japan
email : hitoshi@uguisu.co.jp

Tomotada Ohtsuki

Department of Mathematical and Computing Sciences
Tokyo Institute of Technology
Oh-Okayama, Meguro-ku
Tokyo 152
Japan
email : tomotada@is.titech.ac.jp

Shuji Yamada

Department of Computer Science
Kyoto Sangyo University
Kamigamo Motoyama, Kita-ku
Kyoto 603
Japan
email : yamada@cc.kyoto-su.ac.jp