

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	44 (1998)
Heft:	3-4: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	HOMFLY POLYNOMIAL VIA AN INVARIANT OF COLORED PLANE GRAPHS
Autor:	Murakami, Hitoshi / Yamada, Shuji
Kapitel:	§3. Invariants of links
DOI:	https://doi.org/10.5169/seals-63908

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

§3. INVARIANTS OF LINKS

In this section, we give a graphical definition of the HOMFLY polynomial of oriented links. For an oriented link diagram D , we define $\langle D \rangle_n$ by the following.

$$\left\langle \begin{array}{c} \nearrow \\ \curvearrowright \\ \curvearrowright \end{array} \right\rangle_n = q^{1/2} \left\langle \begin{array}{c} 1 \uparrow \\ \uparrow \end{array} \right\rangle_n - \left\langle \begin{array}{c} 1 \uparrow \\ 1 \curvearrowright \\ 1 \curvearrowright \end{array} \right\rangle_n$$

and

$$\left\langle \begin{array}{c} \nearrow \\ \curvearrowright \\ \curvearrowright \end{array} \right\rangle_n = q^{-1/2} \left\langle \begin{array}{c} 1 \uparrow \\ \uparrow \end{array} \right\rangle_n - \left\langle \begin{array}{c} 1 \uparrow \\ 1 \curvearrowright \\ 1 \curvearrowright \end{array} \right\rangle_n.$$

Then we have

THEOREM 3.1. $\langle D \rangle_n$ is invariant under the Reidemeister moves II and III.

Proof. From Lemma 2.2, we have

$$\begin{aligned} \left\langle \begin{array}{c} \nearrow \\ \curvearrowright \\ \curvearrowright \end{array} \right\rangle_n &= \left\langle \begin{array}{c} 1 \uparrow \\ \uparrow \end{array} \right\rangle_n - q^{1/2} \left\langle \begin{array}{c} 1 \uparrow & 1 \uparrow \\ \hline 1 \curvearrowright & 1 \curvearrowright \end{array} \right\rangle_n \\ &\quad - q^{-1/2} \left\langle \begin{array}{c} 1 \uparrow & 1 \uparrow \\ \hline 1 \curvearrowright & 1 \curvearrowright \end{array} \right\rangle_n + \left\langle \begin{array}{c} 1 \uparrow & 1 \uparrow \\ \hline 1 \curvearrowright & 1 \curvearrowright \end{array} \right\rangle_n \\ &= \left\langle \begin{array}{c} 1 \uparrow \\ \uparrow \end{array} \right\rangle_n + (-q^{1/2} - q^{-1/2} + [2]) \left\langle \begin{array}{c} 1 \uparrow & 1 \uparrow \\ \hline 1 \curvearrowright & 1 \curvearrowright \end{array} \right\rangle_n = \left\langle \begin{array}{c} 1 \uparrow \\ \uparrow \end{array} \right\rangle_n. \end{aligned}$$

We also have

$$\begin{aligned}
 & \left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle = \left\langle \begin{array}{c} \text{Diagram with one strand} \\ n \end{array} \right\rangle - q^{1/2} \left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle \\
 & \quad - q^{-1/2} \left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle + \left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle \\
 & = ([n] - q^{1/2}[n-1] - q^{-1/2}[n-1] + [n-2]) \left\langle \begin{array}{c} \text{Diagram with one strand} \\ n \end{array} \right\rangle + \left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle \\
 & = \left\langle \begin{array}{cc} 1 \uparrow & 1 \downarrow \\ & \end{array} \right\rangle_n
 \end{aligned}$$

from Lemmas 2.1, 2.3 and 2.4 and so $\langle D \rangle_n$ is invariant under the Reidemeister move II. Next we prove the invariance under the Reidemeister move III. Since we have

$$\left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle = q^{1/2} \left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle - \left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle$$

and

$$\left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle = q^{1/2} \left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle - \left\langle \begin{array}{c} \text{Diagram with two strands crossing} \\ n \end{array} \right\rangle,$$

it suffices to show that

$$\text{Diagram 1} = \text{Diagram 2}.$$

Since

$$\text{Diagram 1} = q \left(\text{Diagram 3} - q^{1/2} \text{Diagram 4} \right)$$

$$-q^{1/2} \left(\text{Diagram 5} + \text{Diagram 6} \right)$$

$$= - \left\langle \text{Diagram 7} \right\rangle - q^{1/2} \left(\text{Diagram 8} + \text{Diagram 9} \right)$$

$$= \left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & 3 \nearrow & 1 \downarrow \\ & 2 \nearrow & \\ & 1 \downarrow & \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n - q^{1/2} \left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & 1 \nearrow & 1 \downarrow \\ & 2 \nearrow & \\ & 1 \downarrow & \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n$$

from Lemmas 2.2 and 2.5 and

$$\left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n = q \left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n - q^{1/2} \left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & 1 \nearrow & 1 \downarrow \\ & 2 \nearrow & \\ & 1 \downarrow & \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n$$

$$- q^{1/2} \left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & 1 \nearrow & 1 \downarrow \\ & 2 \nearrow & \\ & 1 \downarrow & \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n + \left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & 1 \nearrow & 1 \downarrow \\ & 2 \nearrow & \\ & 1 \downarrow & \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n$$

$$= - \left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n - q^{1/2} \left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & 1 \nearrow & 1 \downarrow \\ & 2 \nearrow & \\ & 1 \downarrow & \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n + \left\langle \begin{array}{c} 1 \uparrow & & 1 \uparrow \\ & 2 \nearrow & \swarrow 1 \\ 1 \downarrow & 1 \nearrow & 1 \downarrow \\ & 2 \nearrow & \\ & 1 \downarrow & \\ 1 \downarrow & & 1 \downarrow \end{array} \right\rangle_n$$

$$= \left\{ \begin{array}{c} \text{Diagram 1: } \text{A link diagram with } n \text{ strands. The strands are labeled 1, 2, 3, ..., } n \text{ from bottom to top. There are two crossing points. The first crossing has strands 1, 2, 3, 1. The second crossing has strands 1, 2, 1, 1. Arrows indicate orientation: strand 1 goes up, strand 2 goes down, strand 3 goes up, and so on.} \\ \text{Diagram 2: } \text{A link diagram with } n \text{ strands. The strands are labeled 1, 2, 1, 1, ..., 1 from bottom to top. There are two crossing points. The first crossing has strands 1, 2, 1. The second crossing has strands 1, 2, 1, 1. Arrows indicate orientation: strand 1 goes up, strand 2 goes down, strand 1 goes up, and so on.} \end{array} \right\} - q^{1/2} \left\{ \begin{array}{c} \text{Diagram 3: } \text{A link diagram with } n \text{ strands. The strands are labeled 1, 1, 2, 1, ..., 1 from bottom to top. There are two crossing points. The first crossing has strands 1, 1, 2. The second crossing has strands 1, 1, 2, 1. Arrows indicate orientation: strand 1 goes up, strand 1 goes up, strand 2 goes down, and so on.} \end{array} \right\},$$

we have the required formula from Lemma 2.6.

Since the Reidemeister move III with other types of orientations can be obtained from the Reidemeister moves II and III described above (see [12]), the proof is complete. \square

If we define $P_n(D) = q^{(n/2)(-\text{w}(D))} \langle D \rangle_n$ with $\text{w}(D)$ the writhe (the algebraic sum of the crossings) of D , then we have

THEOREM 3.2. $P_n(D)$ is invariant under the Reidemeister moves I, II, and III and satisfies the following skein relation.

$$q^{n/2} P_n(D_+) - q^{-n/2} P_n(D_-) = (q^{1/2} - q^{-1/2}) P_n(D_0),$$

where D_+ , D_- and D_0 are identical link diagrams except near a crossing as described in Figure 3.1.




FIGURE 3.1
skein triple

The proof of this theorem follows immediately from the following lemma.

LEMMA 3.3.

$$\left\langle \begin{array}{c} \uparrow \\ \circlearrowleft \\ \uparrow \end{array} \right\rangle_n = q^{n/2} \left\langle \begin{array}{c} \uparrow^1 \\ \uparrow \end{array} \right\rangle_n$$

and

$$\left\langle \begin{array}{c} \uparrow \\ \circlearrowright \\ \uparrow \end{array} \right\rangle_n = q^{-n/2} \left\langle \begin{array}{c} \uparrow^1 \\ \uparrow \end{array} \right\rangle_n.$$

Proof. From Lemmas 2.1 and 2.3, we have

$$(3.1) \quad \left\langle \begin{array}{c} \uparrow \\ \text{---} \end{array} \right\rangle_n = q^{1/2} \left\langle \begin{array}{c} 1 \uparrow \\ | \\ \text{---} \end{array} \right\rangle_n - \left\langle \begin{array}{c} 1 \uparrow \\ 2 \uparrow \\ | \\ \text{---} \end{array} \right\rangle_n$$

$$(3.2) \quad = (q^{1/2}[n] - [n-1]) \left\langle \begin{array}{c} \uparrow \\ 1 \\ | \\ \text{---} \end{array} \right\rangle_n.$$

Since

$$q^{1/2}[n] - [n-1] = q^{n/2},$$

the first equality follows. The second equality follows similarly and the proof is complete. \square

Therefore P_n defines a link invariant and so we can put $P_n(L) = P_n(D)$ for the link L presented by D . Then P_n is a version of the HOMFLY polynomial [1], [13].

REMARK 3.4. If we define $[D]_n$ to be

$$\left[\begin{array}{c} \uparrow \\ \text{---} \end{array} \right]_n = Aq^{1/2} \left[\begin{array}{c} 1 \uparrow \\ | \\ \text{---} \end{array} \right]_n - A \left[\begin{array}{c} 1 \uparrow \\ 2 \uparrow \\ | \\ \text{---} \end{array} \right]_n$$

and

$$\left[\begin{array}{c} \uparrow \\ \text{---} \end{array} \right]_n = A^{-1}q^{-1/2} \left[\begin{array}{c} 1 \uparrow \\ | \\ \text{---} \end{array} \right]_n - A^{-1} \left[\begin{array}{c} 1 \uparrow \\ 2 \uparrow \\ | \\ \text{---} \end{array} \right]_n$$

with A an indeterminate, we also have a framed link invariant.

When $n = 2$ and $A = q^{-1/4}$, we have a version of the Kauffman bracket naturally defined from representation theory of $U_q(\mathfrak{sl}(2, \mathbb{C}))$ [6, Theorem 4.3].

For $n = 3$ we have G. Kuperberg's recursive formula [8] as follows. Consider an oriented, trivalent, plane graph with flow less than or equal to three. If we reverse the orientations of all the edges with flow two, remove all the edges with flow three, and forget the flow, then we have a trivalent graph without flow such that at every vertex all the edges are in or out. Putting $A = q^{-1/6}$ and replacing q with q^{-1} , we have Kuperberg's formula. (This corresponds to the fact that the two-fold anti-symmetric tensor of the vector representation of $SU(3)$ is isomorphic to its dual.) See also [12].