

Zeitschrift: L'Enseignement Mathématique
Herausgeber: Commission Internationale de l'Enseignement Mathématique
Band: 44 (1998)
Heft: 3-4: L'ENSEIGNEMENT MATHÉMATIQUE

Artikel: THE ORIGIN OF REPRESENTATION THEORY
Autor: CONRAD, Keith
Kapitel: 7. RECENT RESULTS
DOI: <https://doi.org/10.5169/seals-63909>

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 07.08.2025

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

irreducible representations of a finite group, a theorem of Brauer says this number equals the number of conjugacy classes in the group consisting of elements with order prime to p .

7. RECENT RESULTS

Character tables do not provide a way to distinguish any two finite groups, in general. For example, for any prime p the two nonisomorphic nonabelian groups of order p^3 have the same character table. Can we find a computational tool extending the character table which will distinguish any two non-isomorphic finite groups? In 1991, Formanek and Sibley [19] showed that if there is a bijection between two groups G and H which converts $\Theta(G)$ to $\Theta(H)$, then G and H are isomorphic. Since the irreducible characters can be read off (in principle) from the factors of $\Theta(G)$, we see $\Theta(G)$ is one answer to the question. However, if $\#G$ is large then $\Theta(G)$ is too hard to compute. Is there something closer to the character table which works? Yes. See the articles of Hoehnke and Johnson [28, 29] and Johnson and Sehgal [31].

ACKNOWLEDGEMENTS. I thank T. Geisser, K. Khuri-Makdisi, T. Klenke, T. Y. Lam, A. Ofer, R. Pollack, J.-P. Serre, and R. Vakil for their assistance, and MSRI for its hospitality, where part of this was written.

REFERENCES

- [1] ARTIN, E. Über eine neue Art von L -Reihen. *Math. Ann.* 89 (1923), 89–108.
- [2] BOREVICH, Z. I. and I. R. SHAFAREVICH. *Number Theory*. Academic Press, New York, 1966.
- [3] BURNSIDE, W. On a property of certain determinants. *Mess. of Math. (N.S.)* 23 (1894), 112–114.
- [4] —— On the condition of reducibility of any group of linear substitutions. *Proc. London Math. Soc. (2)* 3 (1905), 430–434.
- [5] CATALAN, E. Recherches sur les déterminants. *Bulletins de l'Académie Royale des sciences, des lettres et des beaux-arts de Belgique* 13 (1846), 534–555.
- [6] CREMONA, L. Intorno ad un teorema di Abel. *Annali di Scienze matematiche e fisiche* 7 (1856), 99–105.
- [7] CURTIS, C. Representation theory of finite groups: from Frobenius to Brauer. *Math. Intelligencer* 14 (1992), 48–57.
- [8] CURTIS, C. and I. REINER. *Methods of Representation Theory, Vol. 1*. John Wiley & Sons, New York, 1981.