

Zeitschrift:	L'Enseignement Mathématique
Herausgeber:	Commission Internationale de l'Enseignement Mathématique
Band:	44 (1998)
Heft:	1-2: L'ENSEIGNEMENT MATHÉMATIQUE
 Artikel:	THREE DISTANCE THEOREMS AND COMBINATORICS ON WORDS
Autor:	Alessandri, Pascal / Berthé, Valérie
Kapitel:	3. The three distance theorem
DOI:	https://doi.org/10.5169/seals-63900

Nutzungsbedingungen

Die ETH-Bibliothek ist die Anbieterin der digitalisierten Zeitschriften auf E-Periodica. Sie besitzt keine Urheberrechte an den Zeitschriften und ist nicht verantwortlich für deren Inhalte. Die Rechte liegen in der Regel bei den Herausgebern beziehungsweise den externen Rechteinhabern. Das Veröffentlichen von Bildern in Print- und Online-Publikationen sowie auf Social Media-Kanälen oder Webseiten ist nur mit vorheriger Genehmigung der Rechteinhaber erlaubt. [Mehr erfahren](#)

Conditions d'utilisation

L'ETH Library est le fournisseur des revues numérisées. Elle ne détient aucun droit d'auteur sur les revues et n'est pas responsable de leur contenu. En règle générale, les droits sont détenus par les éditeurs ou les détenteurs de droits externes. La reproduction d'images dans des publications imprimées ou en ligne ainsi que sur des canaux de médias sociaux ou des sites web n'est autorisée qu'avec l'accord préalable des détenteurs des droits. [En savoir plus](#)

Terms of use

The ETH Library is the provider of the digitised journals. It does not own any copyrights to the journals and is not responsible for their content. The rights usually lie with the publishers or the external rights holders. Publishing images in print and online publications, as well as on social media channels or websites, is only permitted with the prior consent of the rights holders. [Find out more](#)

Download PDF: 14.01.2026

ETH-Bibliothek Zürich, E-Periodica, <https://www.e-periodica.ch>

We deduce from Theorem 8 that the lengths of the intervals $I(w_1, \dots, w_n)$, and thus the lengths of the intervals obtained by placing the points $0, \{1 - \alpha\}, \dots, \{n(1 - \alpha)\}$ on the unit circle, take at most three values. Hence Theorem 8 is equivalent to the three distance theorem and provides a combinatorial proof of this result.

REMARK. In fact this point of view, and more precisely the study of the evolution of the graphs of words with respect to the length n of the factors, allows us to give a proof of the most complete version of the three distance theorem as given in [53] (for more details, the reader is referred to [3]).

3. THE THREE DISTANCE THEOREM

The three distance theorem was initially conjectured by Steinhaus, first proved by V.T. Sós (see [53] and also [52]), and then by Świerczkowski [56], Surányi [55], Slater [51], Halton [31]. More recent proofs have also been given by van Ravenstein [44] and Langevin [35]. A survey of the different approaches used by these authors is to be found in [44, 51, 35]. In the literature this theorem is called *the Steinhaus theorem*, *the three length*, *three gap* or *the three step theorem*. In order to avoid any ambiguity, we will always call it the three distance theorem, reserving the name *three gap* for the theorem introduced in the next section.

THREE DISTANCE THEOREM. *Let $0 < \alpha < 1$ be an irrational number and n a positive integer. The points $\{i\alpha\}$, for $0 \leq i \leq n$, partition the unit circle into $n+1$ intervals, the lengths of which take at most three values, one being the sum of the other two.*

More precisely, let $(\frac{p_k}{q_k})_{k \in \mathbb{N}}$ and $(c_k)_{k \in \mathbb{N}}$ be the sequences of convergents and partial quotients associated to α in its continued fraction expansion (if $\alpha = [0, c_1, c_2, \dots]$, then $\frac{p_n}{q_n} = [0, c_1, \dots, c_n]$). Let $\eta_k = (-1)^k(q_k\alpha - p_k)$. Let n be a positive integer. There exists a unique expression for n of the form

$$n = mq_k + q_{k-1} + r,$$

with $1 \leq m \leq c_{k+1}$ and $0 \leq r < q_k$. Then the circle is divided by the points $0, \{\alpha\}, \{2\alpha\}, \dots, \{n\alpha\}$ into $n+1$ intervals which satisfy:

- $n + 1 - q_k$ of them have length η_k (which is the largest of the three lengths),
- $r + 1$ have length $\eta_{k-1} - m\eta_k$,
- $q_k - (r + 1)$ have length $\eta_{k-1} - (m - 1)\eta_k$.

REMARKS.

- One can reformulate this result in terms of n -Farey points. Let us recall that an n -Farey point is a rational element $\frac{p}{q}$ of $[0, 1]$ such that $p \geq 0$, $1 \leq q \leq n$ and p, q are coprime (see [32] for instance). Note that the two successive n -Farey points, say $\frac{p^{(1)}}{q^{(1)}}$ and $\frac{p^{(2)}}{q^{(2)}}$, satisfying $\frac{p^{(1)}}{q^{(1)}} < \alpha < \frac{p^{(2)}}{q^{(2)}}$ are $\frac{p_k}{q_k}$ and $\frac{mp_k + p_{k-1}}{mq_k + q_{k-1}}$, with the above notation. The three distance theorem states that the lengths of the intervals belong to the set

$$\{p^{(2)} - \alpha q^{(2)}, \alpha q^{(1)} - p^{(1)}, \alpha(q^{(1)} - q^{(2)}) + p^{(2)} - p^{(1)}\}.$$

- As α is irrational, the three lengths are distinct. The third length in the above theorem, which is the largest since it is the sum of the two others, appears if and only if

$$n \neq q^{(1)} + q^{(2)} - 1 = (m + 1)q_k + q_{k-1} - 1.$$

Thus there are infinitely many integers n for which there are only two lengths. The other two lengths do always appear.

- The structure and the transformation rules for the partitioning in two-length intervals are studied in details in [44]. Furthermore, in [45] van Ravenstein, Winley and Tognetti prove the following: for α having as sequence of partial quotients the constant sequence $aaaa\cdots$, label by large and small the lengths of intervals of the partition $\{i\alpha\}$, for $0 \leq i \leq q_n + q_{n-1} - 1$, where q_n is the denominator of a reduced convergent of α (there are only two lengths in this case); this binary finite sequence of lengths is a prefix after a permutation of the characteristic sequence of α (i.e., the Sturmian coding of the orbit of α). For a precise study of the limit points of these finite binary sequences (corresponding to the two-length case), see [48].

- In the two-length case, it is easily seen that the largest length is less than or equal to twice the second one. In [14] (see also [15, 16]) Chevallier extends this result to the two-dimensional torus \mathbf{T}^2 , by studying the notion of best approximation.

- The point $\{(n + 1)\alpha\}$ belongs to an interval of largest length in the partition of the unit circle by the points $\{i\alpha\}$, for $0 \leq i \leq n$.

- The three distance theorem is a geometric illustration of the properties of good approximation of the n -Farey points. Indeed, the two intervals containing 0 are of distinct lengths and their lengths are the two smallest. We thus have

$$\alpha q^{(1)} - p^{(1)} = \inf \{k\alpha, \text{ for } 0 \leq k \leq n\}$$

and

$$p^{(2)} - \alpha q^{(2)} = 1 - \sup \{k\alpha, \text{ for } 0 \leq k \leq n\}.$$

- For a deeper study of the rational case, the reader is referred for instance to [51].

4. THE THREE GAP THEOREM

The following theorem, called the *three gap theorem*, is in some sense the dual of the three distance theorem. This theorem was first proved by Slater (see [49] and see also [50, 51]), in the early fifties, whereas the first proofs of the three distance theorem date back to the late fifties. For other proofs of the three-gap theorem, see also [25], and more recently, [58] and [35].

The formulation of the three gap theorem quoted below is due to Slater. Following the notation of [51], let k_i be the sequence of integers k satisfying $k\alpha < \beta$. Then any difference $k_{i+1} - k_i$ is called a *gap*. Moreover, the *frequency* of a gap is defined as its frequency in the sequence of the successive gaps $(k_{i+1} - k_i)_{i \in \mathbb{N}}$.

THREE GAP THEOREM. *Let α be an irrational number in $]0, 1[$ and let $\beta \in]0, 1/2[$. The gaps between the successive integers j such that $\{j\alpha\} < \beta$ take at most three values, one being the sum of the other two.*

More precisely, let $(\frac{p_k}{q_k})_{k \in \mathbb{N}}$ and $(c_k)_{k \in \mathbb{N}}$ be the sequences of the convergents and partial quotients associated to α in its continued fraction expansion. Let $\eta_k = (-1)^k(q_k\alpha - p_k)$. There exists a unique expression for β of the form

$$\beta = m\eta_k + \eta_{k+1} + \psi,$$

with $k \geq 0$, $0 < \psi \leq \eta_k$, and if $k = 0$ then $1 \leq m \leq c_1 - 1$; otherwise, $1 \leq m \leq c_{k+1}$. Then the gaps between two successive j such that $\{j\alpha\} \in [0, \beta[$ satisfy the following:

- *the gap q_k has frequency $(m - 1)\eta_k + \eta_{k+1} + \psi$,*
- *the gap $q_{k+1} - mq_k$ has frequency ψ ,*
- *the gap $q_{k+1} - (m - 1)q_k$ has frequency $\eta_k - \psi$.*